

Research Article

Open Access

David Bryant, André Nies, and Paul Tupper*

A Universal Separable Diversity

<https://doi.org/10.1515/agms-2017-0008>

Received September 6, 2017; accepted November 10, 2017

Abstract: The Urysohn space is a separable complete metric space with two fundamental properties: (a) *universality*: every separable metric space can be isometrically embedded in it; (b) *ultrahomogeneity*: every finite isometry between two finite subspaces can be extended to an auto-isometry of the whole space. The Urysohn space is uniquely determined up to isometry within separable metric spaces by these two properties. We introduce an analogue of the Urysohn space for diversities, a recently developed variant of the concept of a metric space. In a diversity any finite set of points is assigned a non-negative value, extending the notion of a metric which only applies to unordered pairs of points. We construct the unique separable complete diversity that it is ultrahomogeneous and universal with respect to separable diversities.

Keywords: Diversities, Urysohn space, Katětov functions, universality, ultrahomogeneity

MSC: 51F99, 54E50, 54E99

1 Introduction

Urysohn [10] in 1927 constructed a remarkable metric space which is now named after him. The Urysohn space is the unique (up to isometry) separable complete metric space with the following two properties: (a) *universality*: all separable metric spaces can be isometrically embedded within it; (b) *ultrahomogeneity*: any isometry between two finite subspaces of the Urysohn space can be extended to an auto-isometry of the whole space.

The property of universality is straightforward to grasp, and holds for several separable complete metric spaces, such as $C[0, 1]$. The property of ultrahomogeneity is less known. Recall that homogeneity of a metric space means that given any two points x, y in the space, there is an automorphism (or self-isometry) of the space that maps x to y . Likewise, a space is 2-homogeneous if for every pair of pairs (x_1, x_2) and (y_1, y_2) such that $d(x_1, x_2) = d(y_1, y_2)$, there is an automorphism of the space taking x_1 to y_1 and x_2 to y_2 . For any $k \geq 1$, k -homogeneity is defined similarly. Ultrahomogeneity just extends this property to any pair of isometric finite subsets of the space. An example of a complete separable ultrahomogeneous space is the separable infinite-dimensional Hilbert space ℓ_2 ; see Melleray [8]. Urysohn established that the Urysohn space is the unique (up to isometry) separable metric space satisfying both universality and ultrahomogeneity [8].

Here we construct the analogue of the Urysohn space for *diversities*, a generalization of the concept of metric spaces wherein all finite subsets, and not just pairs of points, are assigned a non-negative value. A diversity is a pair (X, δ) where X is a set and δ is a function from the finite subsets of X to \mathbb{R} satisfying

- (D1) $\delta(A) \geq 0$, and $\delta(A) = 0$ if and only if $|A| \leq 1$.
- (D2) If $B \neq \emptyset$ then $\delta(A \cup B) + \delta(B \cup C) \geq \delta(A \cup C)$

for all finite $A, B, C \subseteq X$. Diversities were introduced in [2]. They form an extension of the concept of a metric space. Indeed, every diversity has an *induced metric*, given by $d(a, b) = \delta(\{a, b\})$ for all $a, b \in X$. Note also

David Bryant: Dept. of Mathematics and Statistics, University of Otago, New Zealand, E-mail: david.bryant@otago.ac.nz

André Nies: Dept. of Computer Science, University of Auckland, New Zealand, E-mail: andre@cs.auckland.ac.nz

***Corresponding Author: Paul Tupper:** Dept. of Mathematics, Simon Fraser University, Canada, E-mail: pft3@sfu.ca

that δ is *monotonic*: $A \subseteq B$ implies $\delta(A) \leq \delta(B)$. Also δ is *subadditive on sets with nonempty intersection*: $\delta(A \cup B) \leq \delta(A) + \delta(B)$ when $A \cap B \neq \emptyset$ [2, Prop. 2.1]. We say that a diversity (X, δ) is complete if its induced metric (X, d) is complete [9], and that a diversity is separable if its induced metric is separable.

Our main goal is to construct the diversity analog $(\mathbb{U}, \delta_{\mathbb{U}})$ of the Urysohn metric space. It is determined uniquely by being universal for separable diversities, and ultrahomogeneous in the sense that isometric finite subdiversities are automorphic.

The construction follows the same approach as Katětov's construction of the Urysohn universal metric space [7]. Starting with any diversity (X, δ) , we denote by $E(X)$ the set of one-point extensions of X . Since $E(X)$ turns out to not be separable under the natural metric, we instead consider extensions with *finite support*, which provides a separable diversity $E(X, \omega)$ in which (X, δ) is naturally embedded. Repeating this procedure we obtain a nested sequence of separable diversities. The analogue of the Urysohn metric space is constructed as the completion of the direct limit of all these diversities. Finally we show that this complete separable diversity has the diversity analogue of Urysohn's extension property, and hence is universal and ultrahomogeneous.

Another aspect of metric space theory that has been successfully carried over to diversities is tight span theory [5]. The tight span of a metric space (X, d) can be viewed as the set of all *minimal* members of $E(X)$. The connections between the tight span of a metric space, the space of all one-point extensions $E(X)$, and the Urysohn space are explored in [1]. It is shown, for example, that the tight span of the Urysohn space is not separable. Similar themes have been developed for diversities. In [2] tight span theory was studied for diversities, where the tight span of a diversity (X, δ) consists of all minimal one-point extensions with an appropriate diversity, and therefore can be seen as a subset of $E(X)$.

2 Background and Preliminaries

Recall from above that any diversity (X, δ) has an induced metric (X, d) where $d(a, b) = \delta(\{a, b\})$ for all $a, b \in X$. Conversely, given any metric space (X, d) , consider the diversities that have (X, d) as an induced metric. Lower and upper bounds on the possible diversities that have (X, d) as the induced metric are provided by the *diameter diversity* and the *Steiner diversity*, which we now introduce.

For any metric space (X, d) , the corresponding diameter diversity $(X, \delta_{\text{diam}})$ is defined by

$$\delta_{\text{diam}}(A) = \max_{a, b \in A} d(a, b)$$

for all finite $A \subseteq X$.

On the other hand, given a metric space (X, d) , consider the weighted complete graph (X, E, w) where X is the set of vertices, E is the set of all unordered pairs of vertices, and w assigns weight $d(a, b)$ to the edge (a, b) . A tree T with vertices in X covers a finite set $A \subseteq X$ if A is a subset of the vertices of T . The Steiner diversity $(X, \delta_{\text{Steiner}})$ is defined by letting $\delta_{\text{Steiner}}(A)$ be the infimum, over all trees that cover A , of the total weight of the tree.

The diameter diversity and the Steiner diversity of a metric space (X, d) are important in that for any other diversity (X, δ) that has (X, d) as an induced metric space we have

$$\delta_{\text{diam}}(A) \leq \delta(A) \leq \delta_{\text{Steiner}}(A),$$

for all finite $A \subseteq X$ [3].

The two diversities δ_{diam} and δ_{Steiner} are determined purely by their values on pairs of points. We will show in the last section of this paper that the diversity analogue of the Urysohn metric space is neither a diameter diversity nor a Steiner diversity of any metric space. In particular, it is neither the diameter diversity nor the the Steiner diversity of the Urysohn metric space, even though it has the Urysohn metric space as its induced metric space.

3 Analogue of Katětov functions

For a metric space (X, d) , a Katětov function $f: X \rightarrow \mathbb{R}$ describes a potential one-point extension of X by a point z : a metric \widehat{d} on $X \cup \{z\}$ extending d is given by defining $\widehat{d}(x, z) = f(x)$ for each $x \in X$. By [7] we have

$$f \in E(X) \Leftrightarrow \forall x \forall y |f(x) - f(y)| \leq d(x, y) \leq f(x) + f(y). \quad (3.1)$$

$E(X)$ is the set of Katětov functions, which form a metric space with the sup distance $d_\infty(f, g) = \sup_x |f(x) - g(x)|$. Identifying $x \in X$ with the function $y \mapsto d(x, y)$ isometrically embeds X into $E(X)$.

Let $\mathcal{P}_{\text{fin}}(X)$ denote the collection of all finite subsets of X . Let (X, δ) be a diversity. We will define its extension $E(X)$ by adapting Katětov's approach [7].

Definition 3.1. A function $f: \mathcal{P}_{\text{fin}}(X) \rightarrow \mathbb{R}$ is called admissible if for some point z , $(X \cup \{z\}, \widehat{\delta})$ is a diversity, where

$$\widehat{\delta}(A) = \delta(A), \quad \widehat{\delta}(A \cup \{z\}) = f(A)$$

for all finite $A \subseteq X$. The point z may be in X .

As before, each admissible function on (X, δ) corresponds to a way of extending (X, δ) by one point z . We let $E(X)$ be the set of all admissible functions on (X, δ) . We provide the analogue of (3.1).

Lemma 3.2. A function $f: \mathcal{P}_{\text{fin}}(X) \rightarrow \mathbb{R}$ is in $E(X)$ if and only if f satisfies the following:

- (i) $f(\emptyset) = 0$,
- (ii) $f(A) \geq \delta(A)$, for all A ,
- (iii) $f(A \cup C) + \delta(B \cup C) \geq f(A \cup B)$, for all A, B , and C with $C \neq \emptyset$
- (iv) $f(A) + f(B) \geq f(A \cup B)$.

Proof. \Rightarrow : Suppose f is admissible, so $\widehat{\delta}$ is a diversity on $X \cup \{z\}$, and $f(A) = \widehat{\delta}(A \cup \{z\})$ for all $A \in \mathcal{P}_{\text{fin}}(X)$. Then $\widehat{\delta}(\{z\}) = 0$ implies property (i). Monotonicity of $\widehat{\delta}(A)$ implies $f(A) = \widehat{\delta}(A \cup \{z\}) \geq \widehat{\delta}(A) = \delta(A)$, which is property (ii). The triangle inequality (D2) for $\widehat{\delta}$ gives, for all $C \neq \emptyset$,

$$f(A \cup C) + \delta(B \cup C) = \widehat{\delta}(A \cup C \cup \{z\}) + \widehat{\delta}(B \cup C) \geq \widehat{\delta}(A \cup B \cup \{z\}) = f(A \cup B),$$

which is property (iii). Finally, using the triangle inequality for $\widehat{\delta}$ again gives

$$f(A) + f(B) = \widehat{\delta}(A \cup \{z\}) + \widehat{\delta}(B \cup \{z\}) \geq \widehat{\delta}(A \cup B \cup \{z\}) = f(A \cup B),$$

which is property (iv).

\Leftarrow : Suppose now that f satisfies the properties (i) through (iv). If $f(\{x\}) = 0$ for some $x \in X$, let $z = x$. Otherwise let $z \notin X$. Define $\widehat{\delta}$ on $X \cup \{z\}$ by

$$\widehat{\delta}(A) = \delta(A), \quad \widehat{\delta}(A \cup \{z\}) = f(A)$$

for all finite $A \subseteq X$. Note that if $z \in A$, we have $\delta(A) \leq f(A) \leq \delta(A) + f(\{z\}) = \delta(A)$ using properties (ii) and (iii), so that the definition is consistent. We need to show that $\widehat{\delta}$ is a diversity function. Since δ is a diversity, $\widehat{\delta}(\{x\}) = 0$ for all $x \in X$, and Property (i) gives that $\widehat{\delta}(\{z\}) = 0$. For monotonicity, we need to show $\widehat{\delta}(A \cup \{y\}) \geq \widehat{\delta}(A)$ for four different cases. First, if z is not equal to y and not in A , then it follows from monotonicity of δ . Secondly, if $z = y$ and $z \notin A$ then

$$\widehat{\delta}(A \cup \{z\}) = f(A) \geq \delta(A) = \widehat{\delta}(A)$$

by property (ii). Thirdly, if $z \neq y$ and $z \in A$ then

$$\widehat{\delta}(A \cup \{y\}) = f(A \setminus \{z\} \cup y) \geq f(A \setminus \{z\}) - \delta(\{y\}) = \widehat{\delta}(A) - 0$$

by property (iii). Fourthly, if $z = y$ and $z \in A$ we have $A \cup \{y\} = A$ and hence $\widehat{\delta}(A \cup \{y\}) = \widehat{\delta}(A)$. To show that $\widehat{\delta}$ is subadditive on intersecting sets we again have several cases. If $z \notin A$ and $z \notin B$ then

$$\widehat{\delta}(A) + \widehat{\delta}(B) = \delta(A) + \delta(B) \geq \delta(A \cup B) = \widehat{\delta}(A \cup B).$$

If z is in A but not in B then

$$\widehat{\delta}(A) + \widehat{\delta}(B) = f(A \setminus \{z\}) + \delta(B) \geq f(A \setminus \{z\} \cup B) = \widehat{\delta}(A \cup B).$$

using property (iii). Likewise if z is in B but not in A . Finally, suppose $z \in A \cap B$. Then

$$\widehat{\delta}(A) + \widehat{\delta}(B) = f(A \setminus \{z\}) + f(B \setminus \{z\}) \geq f((A \cup B) \setminus \{z\}) = \widehat{\delta}(A \cup B).$$

Hence $\widehat{\delta}$ is subadditive on intersecting sets. Together with monotonicity this gives the triangle inequality for diversities. \square

Analogous to the metric d_∞ in Katětov’s construction, we define a diversity function $\widehat{\delta}$ on $E(X)$. The motivating idea for our choice of function is that since every admissible function f corresponds to extending a diversity by an additional point z , considering admissible functions f_1, \dots, f_k should require us to extend the diversity by points z_1, \dots, z_k simultaneously, giving a new diversity δ_E defined on $X \cup \{z_1, \dots, z_k\}$. This diversity must coincide with δ on X , and also satisfy that $f_i(A) = \delta_E(A \cup \{z_i\})$ for $i = 1, \dots, k$. Once we have fixed a choice of δ_E given these constraints, we let

$$\widehat{\delta}(\{f_1, \dots, f_k\}) = \delta_E(\{z_1, \dots, z_k\}).$$

One choice for $\widehat{\delta}$ that turns out to generalize from the metric case nicely is to let $\widehat{\delta}$ to be the minimum diversity satisfying the constraints

$$\widehat{\delta}(A) = \delta(A), \quad \widehat{\delta}(A \cup \{z_i\}) = f_i(A), \quad i = 1, \dots, k, \tag{3.2}$$

for all finite $A \subseteq X$. We now describe how to obtain an explicit expression for $\widehat{\delta}$.

We say that a collection of finite subsets E_1, \dots, E_k is *connected* if, when we partition E_1, \dots, E_k into two non-empty collections of sets, there is an E_i on one side of the partition and an E_j on the other side of the partition such that $E_i \cap E_j \neq \emptyset$. Equivalently, define a graph with v_1, \dots, v_k corresponding to E_1, \dots, E_k and there is an edge between v_i and v_j if and only if $E_i \cap E_j \neq \emptyset$. Then the collection of sets is *connected* if and only if the graph is connected.

To determine $\widehat{\delta}$, we first obtain some lower bounds on $\widehat{\delta}(\{z_1, \dots, z_k\})$. Choose any j from $1, \dots, k$. For $i \neq j$ choose finite subsets A_i of X . The sets $A_i \cup \{z_i\}$, $i \neq j$ together with $\{z_1, \dots, z_k\}$ form a connected cover of the set $\{z_j\} \cup \bigcup_{i \neq j} A_i$. So by the triangle inequality for diversities, we should have that

$$\widehat{\delta} \left(\{z_j\} \cup \bigcup_{i \neq j} A_i \right) \leq \widehat{\delta}(\{z_1, \dots, z_k\}) + \sum_{i \neq j} \widehat{\delta}(A_i \cup \{z_i\}).$$

Putting this into terms of admissible functions we get

$$f_j \left(\bigcup_{i \neq j} A_i \right) \leq \widehat{\delta}(\{f_1, \dots, f_k\}) + \sum_{i \neq j} f_i(A_i).$$

This gives the following lower bound on $\widehat{\delta}$:

$$\widehat{\delta}(\{f_1, \dots, f_k\}) \geq f_j \left(\bigcup_{i \neq j} A_i \right) - \sum_{i \neq j} f_i(A_i).$$

Now this bound must hold for each choice of j and A_i for $i \neq j$, which suggests the following definition of $\widehat{\delta}$ on $E(X)$:

$$\widehat{\delta}(\{f_1, \dots, f_k\}) = \max_{j=1, \dots, k} \sup_{A_1, \dots, A_k} \left\{ f_j \left(\bigcup_{i \neq j} A_i \right) - \sum_{i \neq j} f_i(A_i) \right\} \tag{3.3}$$

where all A_i are finite subsets of X . We define $\widehat{\delta}(\emptyset)$ and $\widehat{\delta}(\{f\})$ to be zero, for all $f \in E(X)$. Theorem 3.3 below shows that this is a diversity on $E(X)$ which extends (X, δ) naturally. The considerations above show that it is the minimal diversity satisfying conditions (3.2). Note that if $k = 2$ we simply have

$$\widehat{\delta}(\{f_1, f_2\}) = \sup_{B \text{ finite}} |f_1(B) - f_2(B)|.$$

We now make some comparisons between $(E(X), \widehat{\delta})$ and the tightspan diversity of (X, δ) defined in [2]. Points in $E(X)$ correspond to one-point extensions of the diversity (X, δ) ; points in the tightspan $T(X)$ of X correspond to *minimal* one-point extensions of (X, δ) . Thus $T(X) \subseteq E(X)$. By Lemma 2.6 of [2], the tightspan diversity δ_T equals the restriction of $\widehat{\delta}$ to $T(X)$, noting that on $T(X)$ the k different suprema we are taking the maxima over in (3.3) are all identical, and hence the expression simplifies.

Theorem 3.3. *$(E(X), \widehat{\delta})$ is a diversity, and (X, δ) is embedded in $(E(X), \widehat{\delta})$ via the map $x \rightarrow \kappa_x$ where $\kappa_x(A) = \delta(A \cup \{x\})$.*

Proof. First note that by construction we get $\widehat{\delta}(\emptyset) = 0$ and $\widehat{\delta}(\{f\}) = 0$ for any single admissible function f . If f and g are distinct members of $E(X)$ then $|f(B) - g(B)| > 0$ for some finite B , so $\widehat{\delta}(\{f, g\}) > 0$.

To show monotonicity of $\widehat{\delta}$, note that restricting the size of the set of elements of $E(X)$ restricts the number of functions that can take the first position in the supremum and restricts that the corresponding A_i must be the empty set. So $\widehat{\delta}$ can only decrease when removing elements from a set.

To show that $\widehat{\delta}$ satisfies the triangle inequality, let F and G be two finite sets of functions in $E(X)$ and let h be another admissible function. Let arbitrary $\epsilon > 0$ be given. By the definition of $\widehat{\delta}$ there is a collection of sets A_i and B_j as well an index j such that

$$\widehat{\delta}(F \cup G) - \epsilon \leq f_j \left(\bigcup_{i \neq j} A_i \cup \bigcup_k B_k \right) - \sum_{i \neq j} f_i(A_i) - \sum_k g_k(B_k).$$

We can assume, without loss of generality, that f_j belongs to one of the admissible functions in F . Adding and subtracting $h(\cup_k B_k)$ gives

$$\begin{aligned} \widehat{\delta}(F \cup G) - \epsilon &\leq f_j(\bigcup_{i \neq j} A_i \cup \bigcup_k B_k) - \sum_{i \neq j} f_i(A_i) - h(\cup_k B_k) + h(\cup_k B_k) - \sum_k g_k(B_k) \\ &\leq \widehat{\delta}(F \cup \{h\}) + \widehat{\delta}(G \cup \{h\}). \end{aligned}$$

This is true for all $\epsilon > 0$ so the triangle inequality holds.

Finally, since $\widehat{\delta}$ restricted to $T(X)$ is the tight span diversity δ_T , and, by Theorem 2.8 of [2], $\delta_T(\kappa(A)) = \delta(A)$ for all $A \in \mathcal{P}_{\text{fin}}(X)$. We conclude that $\widehat{\delta}(\kappa(A)) = \delta(A)$ for all $A \in \mathcal{P}_{\text{fin}}(X)$. □

4 Extensions, supports, and $E(X, \omega)$

Recall that a diversity (X, δ) is separable if the underlying induced metric is separable. Analogous to the metric case, the diversity $(E(X), \widehat{\delta})$ need not be separable, even when (X, δ) is. To get a separable but sufficiently rich subspace of $E(X)$ we develop the concepts of support for admissible functions of diversities.

Definition 4.1. *Let (X, δ) be a diversity, let $S \subseteq X$, and let $f \in E(S)$. We define the extension of f to X as*

$$f_S^X(A) = \inf \left\{ f(B) + \sum_{b \in B} \delta(A_b \cup \{b\}) : \text{finite } B \subseteq S, \bigcup_{b \in B} A_b = A \right\}. \tag{4.1}$$

for finite $A \subseteq X$.

The definition of f_S^X can be viewed as a *one-point amalgamation*. Amalgamation is a concept from algebra that also occurs in model theory. Two structures that share a common substructure are simultaneously embedded into a larger structure. Here the two structures are diversities. One is (X, δ) , and the other is the diversity on $S \cup \{z\}$ corresponding to the function f , where z is a single point that may or may not be in S . Since $S \subseteq X$, the two diversities overlap (have a common substructure) on S . In Lemma 4.3 below, we show that f_S^X is an admissible function, and hence it corresponds to a diversity on $X \cup \{z\}$ that extends both (X, δ) and the diversity on $S \cup \{z\}$ corresponding to f . Furthermore, it is the maximal such extension.

Definition 4.2. Let g be an admissible function on (X, δ) and $S \subseteq X$ be nonempty. If $g = f_S^X$ for some $f \in E(S)$ we say that g has support S . We say that f is finitely supported if it has some finite support S .

In the following we use $g \upharpoonright S$ to denote the restriction of g to S .

Lemma 4.3. Let (X, δ) be a diversity, let $S \subseteq X$, and let $f \in E(S)$. Then f_S^X is an admissible function on X such that $f_S^X(A) = f(A)$ for all finite $A \subseteq S$. Furthermore, it is the unique maximal such extension, in that for any admissible function g such that $g \upharpoonright S = f$, we have $g(A) \leq f_S^X(A)$ for any finite set $A \subseteq X$.

Proof. We first show that f_S^X is an admissible function on X by checking conditions (i) through (iv). (i) follows from the non-negativity of f and δ and setting B to be the empty set. To show (ii) we use property (ii) for f to see that the expression inside the infimum for $f_S^X(A)$ satisfies

$$f(B) + \sum_{b \in B} \delta(A_b \cup \{b\}) \geq \delta(B) + \sum_{i=1}^k \delta(A_{b_i} \cup \{b_i\}) \geq \delta\left(\bigcup_{b \in B} A_b\right) = \delta(A),$$

where we have used condition (ii) for f and then the triangle inequality for diversities. For condition (iii), let C be an arbitrary nonempty set. Then

$$f_S^X(A \cup C) + \delta(B \cup C) = \inf_{D \subseteq S} \inf_{\bigcup_{d \in D} A_d = A \cup C} \left\{ f(D) + \sum_{d \in D} \delta(A_d \cup \{d\}) \right\} + \delta(B \cup C).$$

For each such choice of $\{A_d\}$, $d \in D$, let e be an element of D such that A_e and C intersect. Then from the triangle inequality $\delta(A_e \cup \{e\}) + \delta(B \cup C) \geq \delta(A_e \cup B \cup C \cup \{e\})$. So

$$f_S^X(A \cup C) + \delta(B \cup C) \geq \inf_{D \subseteq S} \inf_{\bigcup_{d \in D} A_d = A \cup C} \left\{ f(D) + \sum_{d \neq e} \delta(A_d \cup \{d\}) + \delta(A_e \cup B \cup C \cup \{e\}) \right\}$$

Now the union of the sets A_d for $d \neq e$ together with $A_e \cup B \cup C$ is $A \cup B \cup C$. So from the definition of $f_S^X(A \cup B \cup C)$ we get

$$f_S^X(A \cup C) + \delta(B \cup C) \geq f_S^X(A \cup B \cup C) \geq f_S^X(A \cup B)$$

the last step following from monotonicity of f_S^X .

For condition (iv) note that for all finite $A, B \subseteq X$

$$\begin{aligned} f_S^X(A \cup B) &= \inf_{D \subseteq S} \inf_{\bigcup_{d \in D} G_d = A \cup B} \left\{ f(D) + \sum_{d \in D} \delta(G_d \cup \{d\}) \right\} \\ &\leq \inf_{E, F \subseteq S} \inf_{\bigcup_{e \in E} A_e = A} \inf_{\bigcup_{f \in F} B_f = B} \left\{ f(E \cup F) + \sum_{e \in E} \delta(A_e \cup \{e\}) + \sum_{f \in F} \delta(B_f \cup \{f\}) \right\} \end{aligned}$$

where we have used the fact that the infimum increases because we restricted it to the case when D is a union of two sets, one of which indexes a cover of A and the other indexes a cover of B (and we've allowed some double counting of indices). Now since $f(E \cup F) \leq f(E) + f(F)$, we can decompose the infimum to get $f_S^X(A \cup B) \leq f_S^X(A) + f_S^X(B)$, as required.

Next we show that f_S^X is an extension of f in that $f_S^X(A) = f(A)$ for all finite $A \subseteq S$. First note that taking $B = A$ and $A_b = \{b\}$ for all $b \in B$ in the definition of f_S^X gives that $f_S^X(A) \leq f(A)$. Secondly, if we use condition (iii) of admissible functions repeatedly in the expression in the infimum we get $f_S^X(A) \geq f(A)$, giving the result.

To show that f_S^X has S as a support, just replace the f with f_S^X in the definition of f_S^X and see that it does not change the result, which you can do since f and f_S^X agree on all subsets of S . \square

Let f be any admissible function on (X, δ) and let $S = X$. Repeated use of property (iii) of admissible functions shows

$$f(B) + \sum_{b \in B} \delta(A_b \cup \{b\}) \geq f(A)$$

in equation (4.1), so equality holds for all A . Hence all admissible functions on (X, δ) have X as a support.

We define

$$E(X, \omega) = \{f \in E(X) : f \text{ is finitely supported}\}$$

Note that $E(X, \omega)$ is a subspace of $E(X)$, and that κ_x is finitely supported for each $x \in X$ since it has support $\{x\}$. So $E(X, \omega)$ with diversity $\widehat{\delta}$ is still an extension of the given diversity (X, δ) .

We now show that $(E(X, \omega), \widehat{\delta})$ is separable whenever (X, δ) is. Recall that separability of a diversity just means separability of the induced metric space.

Lemma 4.4. *Let (X, δ) be a diversity with $|X| = n < \infty$. Then $E(X) = E(X, \omega)$ is homeomorphic to a closed subspace of $\mathbb{R}^{\mathcal{P}_{\text{fin}}(X)}$.*

Proof. Every function $f \in E(X)$ can be naturally identified as an element of $\mathbb{R}^{\mathcal{P}_{\text{fin}}(X)}$. $E(X)$ corresponds to those elements of $\mathbb{R}^{\mathcal{P}_{\text{fin}}(X)}$ with the element satisfying the conditions of an admissible function. Since these conditions consist of a linear equality and some non-strict linear inequalities, the subset of $E(X)$ is closed in $\mathbb{R}^{\mathcal{P}_{\text{fin}}(X)}$. We just need to show that the metric induced by $\widehat{\delta}$ is homeomorphic to the Euclidean metric.

Since $\widehat{\delta}(\{f, g\}) = \sup_{B \in \mathcal{P}_{\text{fin}}(X)} |f(B) - g(B)|$, which is the ℓ_∞ norm, this gives the same topology as the Euclidean norm in $\mathbb{R}^{\mathcal{P}_{\text{fin}}(X)}$. \square

Lemma 4.5. *Let f be an admissible function on the diversity (X, δ) . Let $A = \{a_1, \dots, a_n\}$ and $B = \{b_1, \dots, b_n\}$ be subsets of X , where $\delta(\{a_i, b_i\}) \leq \epsilon$ for $i = 1, \dots, n$. Then*

$$|f(A) - f(B)| \leq n\epsilon.$$

Proof. Using property (iii) of admissible functions

$$\begin{aligned} f(A) &= f(\{a_1, \dots, a_n\}) \\ &\leq f(\{b_1, a_2, \dots, a_n\}) + \delta(\{a_1, b_1\}) \\ &\leq \dots \\ &\leq f(\{b_1, \dots, b_n\}) + \sum_{i=1}^n \delta(\{a_i, b_i\}) \\ &= f(B) + n\epsilon. \end{aligned}$$

Applying the same argument with B and A reversed gives $f(B) \leq f(A) + n\epsilon$. \square

Theorem 4.6. *Let (X, δ) be a separable diversity. Then $(E(X, \omega), \widehat{\delta})$ is a separable diversity.*

Proof. Let D be a countable dense set in (X, δ) . We will show that $(E(D, \omega), \widehat{\delta}_D)$ is separable, and that $(E(D, \omega), \widehat{\delta}_D)$ can be densely embedded in $(E(X, \omega), \widehat{\delta}_X)$.

To show that $(E(D, \omega), \widehat{\delta}_D)$ is separable, note that it is the union, over all finite subsets $S \subseteq D$, of the extensions of (D, δ) with support S . Since each set of extensions is separable (being homeomorphic to a closed subset of a finite-dimensional Euclidean space by Lemma 4.4), and there are only countably many of them, $(E(D, \omega), \widehat{\delta}_D)$ is separable.

To show that $(E(D, \omega), \widehat{\delta}_D)$ is densely embeddable in $(E(X, \omega), \widehat{\delta}_X)$, we define the embedding γ . For $f \in E(D, \omega)$ we will define $\gamma f = \widehat{f}: \mathcal{P}_{\text{fin}}(X) \rightarrow \mathbb{R}$ via $\widehat{f} = f_D^X$. From Lemma 4.3 we have that \widehat{f} is an admissible function on X , \widehat{f} is an extension of f , and D is a support of \widehat{f} .

Next we need to show that for any finite set F of admissible functions on D

$$\widehat{\delta}_X(\gamma F) = \widehat{\delta}_X(F).$$

First note that

$$\begin{aligned} \widehat{\delta}_X(\gamma F) &= \max_j \sup_{A_1, \dots, A_k \subseteq X} \left\{ \gamma f_j(\cup_{i \neq j} A_i) - \sum_{i \neq j} \gamma f_i(A_i) \right\} \\ &\geq \max_j \sup_{A_1, \dots, A_k \subseteq D} \left\{ \gamma f_j(\cup_{i \neq j} A_i) - \sum_{i \neq j} \gamma f_i(A_i) \right\} \\ &= \max_j \sup_{A_1, \dots, A_k \subseteq D} \left\{ f_j(\cup_{i \neq j} A_i) - \sum_{i \neq j} f_i(A_i) \right\} \\ &= \widehat{\delta}_D(F), \end{aligned}$$

where we have used that D is a subset of X and that γf agrees with f on D . To show conversely that $\widehat{\delta}_X(\gamma F) \leq \widehat{\delta}_D(F)$, we need to show that for any choice of j and finite $A_1, \dots, A_k \subseteq X$, we can find finite B_1, \dots, B_k so that $\gamma f_j(\cup_{i \neq j} B_i)$ is arbitrarily close to $\gamma f_j(\cup_{i \neq j} A_i)$ and $\gamma f_i(B_i)$ is arbitrarily close to $\gamma f_i(A_i)$ for all $i \neq j$. That such B_i exist follows from the density of D in X and Lemma 4.5.

We have shown that the map $\gamma: E(D, \omega) \rightarrow E(X, \omega)$ is an embedding. We still need to show that it is a dense embedding. Let $f \in E(X, \omega)$. Suppose f has finite support S , with $|S| = n$ and elements s_1, \dots, s_n . For any $\epsilon > 0$, find a $T \subseteq D$ with $|T| = n$ elements t_1, \dots, t_n such that for any subindices i_1, \dots, i_m of $1, \dots, n$ we have

$$|f(\{t_{i_1}, \dots, t_{i_m}\}) - f(\{s_{i_1}, \dots, s_{i_m}\})| < \epsilon.$$

(This is possible by Lemma 4.5.) Now f restricted to T is still an admissible function. We want to extend it to all of D . For any finite subset A of D , define $g = (f \upharpoonright T)_T^D$. By Lemma 4.3, g is an admissible function on (D, δ) , it is an extension of $f \upharpoonright T$, and it has support T . Now we let $\widehat{g} = \gamma g$ be the image of g under the embedding. We need to show that \widehat{g} is close to f .

The functions $\widehat{g}, f: \mathcal{P}_{\text{fin}}(X) \rightarrow \mathbb{R}$ agree on subsets of T , but \widehat{g} is supported on T and f is supported on S . Let A be an arbitrary finite subset of X . Since T is finite, we have for some $B \subseteq T, B = \{t_{i_1}, \dots, t_{i_m}\}$ and $\{A_b\}_{b \in B}$ with $\cup_{b \in B} A_b = A$

$$\begin{aligned} \widehat{g}(A) &\geq f(B) + \sum_{b \in B} \delta(A_b \cup \{b\}) - \epsilon \\ &\geq f(C) - \epsilon + \sum_{c \in C} \delta(A_c \cup \{c\}) - \epsilon \\ &\geq f(A) - 2\epsilon \end{aligned}$$

where $C \subset S$ and $C = \{s_{i_1}, \dots, s_{i_m}\}$. A similar argument starting with $f(A)$ gives $f(A) \leq \widehat{g}(A) + 2\epsilon$. Together we have $|\widehat{g}(A) - f(A)| \leq 2\epsilon$ for all finite $A \subseteq X$ and so $\delta_X(\{\widehat{g}, f\}) \leq 2\epsilon$ can be made arbitrarily small as required. \square

5 Construction of the diversity analogue of the Urysohn metric space.

Here we define the diversity analogue of the Urysohn metric space. We show that it is the unique universal Polish diversity. We also show that it is ultrahomogeneous.

In what follows we will need the following lemma. For each $k \geq 1$, let δ_k be the function that sends (a_1, \dots, a_k) to $\delta(\{a_1, \dots, a_k\})$.

Proposition 5.1. *Let (X, δ) be a diversity. For each k , the function δ_k is 1-Lipschitz in each argument.*

Proof. Consider varying the i th argument of δ_k from x_i to x'_i . We know from the triangle inequality that

$$\begin{aligned} \delta_k(x_1, \dots, x_i, \dots, x_k) &= \delta(\{x_1, \dots, x_i, \dots, x_k\}) \\ &\leq \delta(\{x_1, \dots, x'_i, \dots, x_k\}) + \delta(\{x_i, x'_i\}) \\ &= \delta_k(x_1, \dots, x'_i, \dots, x_k) + d(x_i, x'_i). \end{aligned}$$

Similarly, $\delta_k(x_1, \dots, x'_i, \dots, x_k) \leq \delta_k(x_1, \dots, x_i, \dots, x_k) + d(x_i, x'_i)$. So

$$|\delta_k(x_1, \dots, x_i, \dots, x_k) - \delta_k(x_1, \dots, x'_i, \dots, x_k)| \leq d(x_i, x'_i)$$

as required. \square

Definition 5.2. *A diversity (X, δ) has the extension property if for any finite subset F of X and any admissible function f on F , there is $x \in X$ such that $f(A) = \delta(A \cup \{x\})$ for any finite $A \subseteq F$.*

The extension property on metric spaces is also known as the Urysohn property [6].

Definition 5.3. *We say a diversity is Polish if its induced metric space is Polish, i.e. it is separable and complete.*

Lemma 5.4. *Let (X, δ_X) and (Y, δ_Y) be diversities where X is separable with a dense subset D_X and Y is complete. Any isomorphism from D_X into Y can be extended to an isomorphism from X into Y .*

Proof. Let ϕ be an isomorphism from D_X into Y . Since ϕ preserves the diversity it also preserves the induced metrics between the two sets and is hence a uniformly continuous map. This means we can extend it to a uniformly continuous function $\bar{\phi}$ between X and Y . To show that $\bar{\phi}$ is an isomorphism, let $A \subset X$ be an arbitrary finite set, with $A = \{a_1, \dots, a_n\}$. For each $k = 1, \dots, n$, let $a_k^1, a_k^2, a_k^3, \dots$ be a sequence in D_X such that with $a_k^i \rightarrow a_k$ as $i \rightarrow \infty$. We define $A^i = \{a_1^i, a_2^i, \dots, a_n^i\}$.

$$\begin{aligned} \delta_Y(\bar{\phi}(A)) &= \delta_Y(\bar{\phi}(\lim_i A^i)) \\ &= \delta_Y(\lim_i \bar{\phi}(A^i)) \\ &= \lim_i \delta_Y(\bar{\phi}(A^i)) \\ &= \lim_i \delta_X(A^i) \\ &= \delta_X(\lim_i A^i) = \delta_X(A). \end{aligned}$$

where we have used the uniform continuity of δ_X and δ_Y , by Proposition 5.1. \square

Theorem 5.5. *Let (X, δ_X) and (Y, δ_Y) be Polish diversities with the extension property. Then (X, δ_X) and (Y, δ_Y) are isomorphic.*

Proof. We mostly follow the proof of [6, Thm. 1.2.5]. Let $\{x_0, x_1, x_2, \dots\}$ be a dense set in X and let $\{y_0, y_1, y_2, \dots\}$ be a dense set in Y . We will define a diversity isomorphism between these dense sets and then extend it to the whole space.

We will construct a sequence of partial diversity isomorphisms $\phi_0, \phi_1, \phi_2, \dots$. Let ϕ_0 be defined on the single point x_0 so that $\phi_0(x_0) = y_0$.

At stage $n > 0$, suppose that ϕ_{n-1} has been defined so that $\{x_0, \dots, x_{n-1}\} \subseteq \text{dom}(\phi_{n-1})$ and $\{y_0, \dots, y_{n-1}\} \subseteq \text{range}(\phi_{n-1})$. If $x_n \in \text{dom}(\phi_{n-1})$ then we let $\phi' = \phi_{n-1}$. Otherwise, let $F = \text{range}(\phi_{n-1})$ and consider the admissible function on F defined by

$$f(A) = \delta_X(\phi_{n-1}^{-1}(A) \cup \{x_n\})$$

for finite $A \subseteq F$. By the extension property of Y there is $y \in Y$ so that

$$\delta_Y(A \cup \{y\}) = f(A) = \delta_X(\phi_{n-1}^{-1}(A) \cup \{x_n\}).$$

We extend ϕ_{n-1} to ϕ' by defining $\phi'(x_n) = y$. Now if $y_n \in \text{range}(\phi')$ then we let $\phi_n = \phi'$ and go on to the next stage. Otherwise apply the above argument to ϕ'^{-1} and use the extension property of X to obtain an extension of ϕ' . Define ϕ_n to be this extension. We have thus finished the definition of ϕ_n . Let ϕ be the union of all ϕ_n we have defined. Then it has the required properties.

Finally we use Lemma 5.4 to extend ϕ between $\{x_0, x_1, x_2, \dots\}$ and $\{y_0, y_1, y_2, \dots\}$ to an isomorphism between X and Y . \square

The completion of a diversity is defined in [9]: we take the completion of the diversity's induced metric space, and then extend the original diversity function to this larger set using continuity.

Following [8] we define the following weakened version of the extension property.

Definition 5.6. *A diversity (X, δ) has the approximate extension property if for any finite subset F of X , any admissible function f on F , and any $\epsilon > 0$, there is an $x \in X$ such that $|\delta(A \cup \{x\}) - f(A)| \leq \epsilon$ for any $A \subseteq F$.*

Lemma 5.7. *If a separable diversity has the approximate extension property, then its completion has the approximate extension property.*

Proof. Let (X, δ) be a diversity that is the completion of dense subset D , where (D, δ) has the approximate extension property. Let F be a finite subset of X , $f \in E(F)$, and $\epsilon > 0$ be given. We need to find a point $y \in X$ such that $|\delta(A \cup \{y\}) - f(A)| \leq \epsilon$ for all $A \subseteq F$.

Order all non-empty subsets of F , A_1, \dots, A_{2^n-1} so that if $A_j \subseteq A_i$ then $j \geq i$. Let $\epsilon_0 = \epsilon/2(2^n + n)$. Define a bijective map γ from F to $\gamma F \subseteq D$ so that for all nonempty $A \subseteq F$, $|\delta(\gamma A) - \delta(A)| < \epsilon_0$, which is possible by Proposition 5.1.

Define $g: \mathcal{P}_{\text{fin}}(\gamma F) \rightarrow \mathbb{R}$ by $g(\emptyset) = 0$ and

$$g(\gamma A_i) = f(A_i) + \epsilon_{A_i}, \quad \text{for } i = 1, \dots, 2^n - 1,$$

where $\epsilon_{A_i} = i\epsilon_0$. Note that g is monotonic by construction. We claim that $g \in E(\gamma F)$.

To show $g \in E(\gamma F)$ we need to verify the four conditions of Lemma 3.2. Condition (i) ($g(\emptyset) = 0$) follows by definition. To obtain condition (ii), note that for non-empty A , $g(\gamma A) = f(A) + \epsilon_A \geq \delta(A) + \epsilon_A \geq \delta(\gamma A) - \epsilon_0 + \epsilon_A \geq \delta(\gamma A)$. For condition (iii), we first observe that for any admissible function f on F and $C \neq \emptyset$ we have from the triangle inequality

$$f(A \cup C) + \delta(B \cup C) = f((A \cup C) \cup C) + \delta(B \cup C) \geq f(A \cup B \cup C).$$

So, given $A, B, C \subseteq F$, with $C \neq \emptyset$,

$$\begin{aligned} g(\gamma A \cup \gamma C) + \delta(\gamma B \cup \gamma C) &\geq f(A \cup C) + \epsilon_{A \cup C} + \delta(B \cup C) - \epsilon_0 \\ &\geq f(A \cup B \cup C) + \epsilon_{A \cup C} - \epsilon_0 \\ &= g(\gamma A \cup \gamma B \cup \gamma C) - \epsilon_{A \cup B \cup C} + \epsilon_{A \cup C} - \epsilon_0 \\ &\geq g(\gamma A \cup \gamma B) \end{aligned}$$

where we use the fact that g is monotonic and that $A \cup B \cup C$ is later than $A \cup C$ on the list of subsets, and so $\epsilon_{A \cup B \cup C} + \epsilon_0 \leq \epsilon_{A \cup C}$. Now for condition (iv) we have

$$\begin{aligned} g(\gamma A) + g(\gamma B) &\geq f(A) + \epsilon_A + f(B) + \epsilon_B \\ &\geq f(A \cup B) + \epsilon_A + \epsilon_B \\ &\geq g(\gamma A \cup \gamma B) - \epsilon_{A \cup B} + \epsilon_A + \epsilon_B \\ &\geq g(\gamma A \cup \gamma B) \end{aligned}$$

since $\epsilon_{A \cup B} \leq \epsilon_A$.

So g is admissible on γF . By the approximate extension property of (D, δ) , there is a point y such that $|g(\gamma A) - \delta(\gamma A \cup \{y\})| \leq \epsilon/2$ for all $A \subseteq F$.

Now for any $A \subseteq F$

$$\begin{aligned} |f(A) - \delta(A \cup \{y\})| &\leq |f(A) - g(\gamma A)| + |g(\gamma A) - \delta(\gamma A \cup \{y\})| + |\delta(\gamma A \cup \{y\}) - \delta(A \cup \{y\})| \\ &\leq \epsilon_A + \epsilon/2 + n\epsilon_0 \leq 2^n \epsilon_0 + \epsilon/2 + n\epsilon_0 \leq \epsilon \end{aligned}$$

as required. \square

Lemma 5.8. *Any complete diversity with the approximate extension property has the extension property.*

Proof. Our proof follows that of the metric case in Theorem 3.4 of [8] and Theorem 1.2.7 of [6].

Let (X, δ) be a complete diversity with the approximate extension property. Let finite $F \subseteq X$ be given, and let $f \in E(F)$. It suffices to show there is a sequence z_0, z_1, \dots in X such that for all p , $|\delta(A \cup \{z_p\}) - f(A)| \leq 2^{-p}$ for all $A \subseteq F$ and $\delta(\{z_p, z_{p+1}\}) \leq 2^{1-p}$. Since X is complete and f is continuous, the sequence will have a limit $z \in X$ such that $\delta(A \cup \{z\}) = f(A)$ for all $A \subseteq F$.

By the approximate extension property of (X, δ) we can define z_0 . To use induction, suppose we have z_0, z_1, \dots, z_p satisfying the conditions and we need to specify z_{p+1} . Let $f_p \in E(F)$ be defined by $f_p(A) = \delta(A \cup \{z_p\})$ for $A \subseteq F$. Note that for all A

$$|f_p(A) - f(A)| = |\delta(A \cup \{z_p\}) - f(A)| \leq 2^{-p}.$$

So $\widehat{\delta}(\{f_p, f\}) \leq 2^{-p}$.

Now let g_p be defined on $F \cup \{z_p\}$ by $g_p(A) = f(A)$, $g_p(A \cup \{z_p\}) = \widehat{\delta}(A \cup \{f_p, f\})$. This is in an admissible function on $F \cup \{z_p\}$ because it is realized by the points $F \cup \{f_p, f\}$ in $E(F)$. So by the approximate extension property there is a $z \in X$ that realizes g_p with error at most $2^{-(p+1)}$. In other words

$$|\delta(A \cup \{z\}) - g_p(A)| \leq 2^{-(p+1)}, \quad |\delta(A \cup \{z, z_p\}) - g_p(A \cup \{z_p\})| \leq 2^{-(p+1)}.$$

The first inequality shows that $|\delta(A \cup \{z\}) - f(A)| \leq 2^{-(p+1)}$ and the second inequality shows that, choosing $A = \emptyset$

$$\delta(\{z_p, z\}) \leq g_p(\{z_p\}) + 2^{-(p+1)} = \widehat{\delta}(\{f_p, f\}) + 2^{-(p+1)} \leq 2^{-p} + 2^{-(p+1)} \leq 2^{-p+1}.$$

Now let $z_{p+1} = z$. \square

Theorem 5.9. *If (X, δ) is a separable diversity with the extension property then its completion also has the extension property.*

Proof. Since (X, δ) has the extension property, it certainly has the approximate extension property. By Lemma 5.7 the completion of (X, δ) has the approximate extension property. Then by Lemma 5.8 the completion of (X, δ) has the extension property, being complete. \square

We now work towards defining a complete separable diversity with the extension property. We start with a given diversity (X, δ) . We let $X_0 = X$, $\delta_0 = \delta$. Now, for $n > 0$ we inductively define (X_n, δ_n) by letting $X_n = E(X_{n-1}, \omega)$ with the diversity $\delta_n = \widehat{\delta}_{n-1}$. We define $(X_\omega, \delta_\omega)$ to be the union of all these diversities, which is well-defined because each (X_n, δ_n) is embedded in (X_{n+1}, δ_{n+1}) .

Theorem 5.10. *For any diversity (X, δ) the diversity $(X_\omega, \delta_\omega)$ has the extension property.*

Proof. Let F be a finite subset of X_ω , and let f be an admissible function on F . F must be contained in X_n for some n . By construction, there is some $x \in X_{n+1}$ such that $f(A) = \delta(A \cup \{x\})$ for all $A \subseteq F$. So there is such an x in X_ω . \square

We define the diversity $(\mathbb{U}, \delta_{\mathbb{U}})$ to be the completion of $(X_{\omega}, \delta_{\omega})$ when (X, δ) is the trivial diversity of a single point. By Theorem 5.9, $(\mathbb{U}, \delta_{\mathbb{U}})$ also has the extension property.

We say that a Polish diversity is universal if any separable diversity is isomorphic to a subset it.

Theorem 5.11. $(\mathbb{U}, \delta_{\mathbb{U}})$ is a universal Polish diversity.

Proof. Let (X, δ) be an arbitrary separable diversity. We construct a sequence of partial isomorphisms whose union is the desired isomorphism. Let x_0, x_1, x_2, \dots be a dense sequence in X . Let y be an arbitrary point in \mathbb{U} . Let ϕ_0 be defined on $\{x_0\}$ by $\phi_0(x_0) = y$. Now suppose that we have an isomorphism ϕ_n from $\{x_0, x_1, \dots, x_n\}$ into \mathbb{U} , with $\phi(x_i) = y_i$ for $i = 1, \dots, n$. Define the admissible function on $\{y_0, \dots, y_n\}$ for finite subset A by $f(A) = \delta(\phi_n^{-1}(A) \cup x_{n+1})$. By the extension property, there is a point y_{n+1} in \mathbb{U} such that $\delta(\phi_n^{-1}(A) \cup x_{n+1}) = f(A) = \delta_{\mathbb{U}}(A \cup y_{n+1})$. Define ϕ_{n+1} by extending ϕ_n with one point with $\phi_{n+1}(x_{n+1}) = y_{n+1}$. Now take the union of all of the ϕ_n to obtain an isomorphism between $\{x_0, x_1, x_2, \dots\}$ and a subset of \mathbb{U} . By Lemma 5.4 this isomorphism can be extended to all of X . \square

A Polish diversity (X, δ) is ultrahomogeneous if given any two isomorphic finite subsets $A, A' \subseteq X$, and any isomorphism $\phi: A \rightarrow A'$, there is an isomorphism of (X, δ) to itself that extends ϕ .

Theorem 5.12. $(\mathbb{U}, \delta_{\mathbb{U}})$ is ultrahomogeneous.

Proof. This proof follows the same plan as Theorem 5.5. Let A, A' be two isomorphic subsets of \mathbb{U} , with isomorphism ϕ between them. Let $\{x_1, x_2, \dots\}$ be a dense subset of $\mathbb{U} \setminus A$ and let $\{y_1, y_2, \dots\}$ be a dense subset of $\mathbb{U} \setminus A'$. Let $\phi_0 = \phi$. Suppose we have defined ϕ_{n-1} so that it is an isomorphism and $A \cup \{x_1, \dots, x_{n-1}\} \subseteq \text{dom}(\phi_{n-1})$ and $A' \cup \{y_1, \dots, y_{n-1}\} \subseteq \text{range}(\phi_{n-1})$. Following the proof of Theorem 5.5 yields a suitable ϕ_n . Taking the union of these ϕ_n and applying Lemma 5.4 gives the desired isomorphism from \mathbb{U} to itself that is an extension of ϕ . \square

Theorem 5.13. Any ultrahomogeneous, universal Polish diversity has the extension property, and is thus isomorphic to $(\mathbb{U}, \delta_{\mathbb{U}})$.

Proof. Let (X, δ) be an ultrahomogeneous, universal Polish diversity. Let F be a finite subset of X and let f be an admissible function on F . So we can define a diversity on $F \cup \{z\}$ for some z such that $f(A) = \delta(A \cup \{z\})$ for $A \subseteq F$. Since (X, δ) is universal, there is an embedding ϕ taking $F \cup \{z\}$ into X . Let $F' = \phi(F)$. Since ϕ is an isomorphism from F to F' , there is an isomorphism ϕ' of the whole space that extends ϕ . Consider the point $\phi'^{-1}(z)$. It satisfies the property that $\delta(A \cup \phi'^{-1}(z)) = f(A)$ for all $A \subseteq F$. \square

6 The relationship between $(\mathbb{U}, \delta_{\mathbb{U}})$ and the Urysohn metric space

We denote the Urysohn metric space by (\mathbb{U}_m, d) . The induced metric space of the diversity $(\mathbb{U}, \delta_{\mathbb{U}})$ in, in fact, isometric to (\mathbb{U}_m, d) . We will show that $(\mathbb{U}, \delta_{\mathbb{U}})$ is neither a diameter diversity nor a Steiner diversity. These notions were recalled at the beginning of Section 2.

Proposition 6.1. 1. The metric space induced by $(\mathbb{U}, \delta_{\mathbb{U}})$ is isometric to (\mathbb{U}_m, d) .

2. $(\mathbb{U}, \delta_{\mathbb{U}})$ is not a diameter diversity.

3. $(\mathbb{U}, \delta_{\mathbb{U}})$ is not a Steiner diversity.

Proof. 1. Recall that (\mathbb{U}_m, d) is up to isometry the unique separable complete metric space with the metric extension property. Since $(\mathbb{U}, \delta_{\mathbb{U}})$ is a separable complete diversity, its induced metric space is also separable and complete.

It remains to show that the induced metric space has the metric extension property, which for \mathbb{U} states that for any finite $A \subseteq \mathbb{U}$ and any $f: \mathbb{U} \rightarrow \mathbb{R}$ satisfying (3.1) at the beginning of Section 3 there is a $z \in \mathbb{U}$

such that $d(z, a) = f(a)$ for all $a \in A$. Such a function f corresponds to a metric space $(A \cup \{z\}, \widehat{d})$, in that $f(a) = \widehat{d}(a, z)$ for all $a \in A$, where \widehat{d} restricted to A coincides with the induced metric restricted to A . Define a diversity $\widehat{\delta}$ on $A \cup \{z\}$ by letting $\widehat{\delta}(B) = \delta_{\mathbb{U}}(B)$ and

$$\widehat{\delta}(B \cup \{z\}) = \delta_{\mathbb{U}}(B) + \min_{b \in B} \widehat{d}(b, z)$$

for $B \subseteq A$. Then $(A \cup \{z\}, \widehat{\delta})$ is a one-point extension of $(A, \delta_{\mathbb{U}})$. Since $(\mathbb{U}, \delta_{\mathbb{U}})$ has the (diversity) extension property, z can be identified with a point in \mathbb{U} . For any $a \in A$,

$$d(a, z) = \delta_{\mathbb{U}}(\{a, z\}) = \widehat{\delta}(\{a, z\}) = \widehat{d}(a, z) = f(a)$$

as required. So the metric induced on \mathbb{U} has the extension property, and therefore is isometric to the Urysohn metric space.

2. Consider the diversity on three points given by $X = \{a, b, c\}$, $\delta(a, b) = \delta(a, c) = \delta(b, c) = 1$ and $\delta(a, b, c) = 2$. (X, δ) is not a diameter diversity, since in that case we would have $\delta(a, b, c) = 1$. Since $(\mathbb{U}, \delta_{\mathbb{U}})$ is universal, and (X, δ) is separable and complete, there is a subset of \mathbb{U} that is isometric to (X, δ) . Subsets of diameter diversities are still diameter diversities, so $(\mathbb{U}, \delta_{\mathbb{U}})$ is not a diameter diversity.

3. Consider the diversity on three points given by $X = \{a, b, c\}$, $\delta(a, b) = \delta(a, c) = \delta(b, c) = 1$ and $\delta(a, b, c) = 1$. Since $(\mathbb{U}, \delta_{\mathbb{U}})$ is universal and (X, δ) is separable and complete, we can identify (X, δ) with a subset of \mathbb{U} . Suppose that $(\mathbb{U}, \delta_{\mathbb{U}})$ is a Steiner diversity. Then we can find trees in \mathbb{U} that cover (a, b, c) and have total weight arbitrarily close to $\delta(a, b, c) = 1$. Suppose we have a tree with total weight less than 1.25, covering $\{a, b, c\}$. We can assume that the tree has leaves a, b, c and a single internal node z , which is possibly not distinct from a, b, c . Let the branches of the tree have lengths α, β, γ , corresponding to the leaves a, b, c respectively. Now we have

$$\alpha + \beta \geq 1, \quad \beta + \gamma \geq 1, \quad \alpha + \gamma \geq 1, \quad \alpha + \beta + \gamma < 1.25.$$

Summing the first three inequalities and dividing by 2 gives $\alpha + \beta + \gamma \geq 3/2$ which contradicts the final inequality. Therefore, $(\mathbb{U}, \delta_{\mathbb{U}})$ is not a Steiner diversity. \square

7 Questions

Many questions that have been considered for the Urysohn metric space also make sense for the Urysohn diversity. For instance, it is easy to show that $(\mathbb{U}, \delta_{\mathbb{U}})$ is compact homogeneous, namely, any isomorphic compact subdiversities are automorphic. This follows the construction in Melleray [8, Section 4.5] for the metric case. It would be worthwhile to study the isometry group of $(\mathbb{U}, \delta_{\mathbb{U}})$ along the lines of the results surveyed in [8, Section 4.5].

Another interesting avenue to explore is determining universal and ultrahomogeneous structures for restricted classes of diversities, as is described for metric spaces in [11]. For example, for $S \subseteq [0, \infty)$ let \mathcal{M}_S be the class of all finite metric spaces with distances taking values in S . One can ask for each S whether there is a metric space that is universal and ultrahomogeneous with respect to metrics in \mathcal{M}_S . For example, if $S = [0, \infty)$ then the corresponding structure is the Urysohn space. Delhommé, Laflamme, Pouzet, and Sauer [4] give a complete characterization of which sets S admit such a structure. These questions have natural analogues for diversities.

Acknowledgement: DB was supported by a University of Otago Strategic Grant. AN was partially supported by the Marsden fund of New Zealand. Some of this work was carried out while AN visited the Institute for Mathematical Sciences, Singapore. PT was supported by an NSERC Discovery Grant, an NSERC Discovery Grant and a Tier 2 Canada Research Chair.

References

- [1] Asuman Guven Aksoy and Zair Ibragimov. Convexity of the Urysohn universal space. *Journal of Nonlinear and Convex Analysis*, 17(6):1239–1247, 2016.
- [2] David Bryant and Paul F. Tupper. Hyperconvexity and tight-span theory for diversities. *Advances in Mathematics*, 231(6):3172–3198, 2012.
- [3] David Bryant and Paul F. Tupper. Diversities and the geometry of hypergraphs. *Discrete Math. Theor. Comput. Sci.*, 16(2):1–20, 2014.
- [4] Christian Delhommé, Claude Laflamme, Maurice Pouzet, and Norbert Sauer. Divisibility of countable metric spaces. *European Journal of Combinatorics*, 28(6):1746–1769, 2007.
- [5] Andreas W. M. Dress. Trees, tight extensions of metric spaces, and the cohomological dimension of certain groups: a note on combinatorial properties of metric spaces. *Adv. Math.*, 53(3):321–402, 1984.
- [6] Su Gao. *Invariant descriptive set theory*, volume 293 of *Pure and Applied Mathematics (Boca Raton)*. CRC Press, Boca Raton, FL, 2009.
- [7] Miroslav Katětov. On universal metric spaces. In *General topology and its relations to modern analysis and algebra, VI (Prague, 1986)*, volume 16 of *Res. Exp. Math.*, pages 323–330. Heldermann, Berlin, 1988.
- [8] Julien Melleray. Some geometric and dynamical properties of the Urysohn space. *Topology and its Applications*, 155(14):1531–1560, 2008.
- [9] Andrew Poelstra. On the topological and uniform structure of diversities. *Journal of Function Spaces and Applications*, 2013:9 pages, 2013.
- [10] Paul Urysohn. Sur un espace métrique universel. *Bull. Sci. Math*, 51(2):43–64, 1927.
- [11] Lionel Nguyen Van Thé. *Structural Ramsey theory of metric spaces and topological dynamics of isometry groups*. American Mathematical Soc., 2010.