

## Nests of the black stork *Ciconia nigra* as a habitat for mesostigmatid mites (Acari: Mesostigmata)

Jerzy BŁOSZYK<sup>1,2</sup>, Dariusz J. GWIAZDOWICZ<sup>3</sup>, Bruce HALLIDAY<sup>4</sup>, Paweł T. DOLATA<sup>5</sup>  
 & Bartłomiej GOŁDYN<sup>1</sup>

<sup>1</sup>Department of General Zoology, Faculty of Biology, Adam Mickiewicz University, Umultowska 89, 61-614 Poznań, Poland; e-mail: bloszyk@amu.edu.pl

<sup>2</sup>Natural Science Collection, Faculty of Biology, Adam Mickiewicz University, Umultowska 89, 61-614 Poznań, Poland; e-mail: bloszyk@amu.edu.pl

<sup>3</sup>Department of Forest and Environment Protection, August Cieszkowski Agricultural University, Wojska Polskiego 71C, 60-625 Poznań, Poland; e-mail: dagwiazd@au.poznan.pl

<sup>4</sup>CSIRO Entomology, GPO Box 1700, Canberra ACT 2601, Australia; e-mail: Bruce.Halliday@csiro.au

<sup>5</sup>South Wielkopolska Group of Polish Society for the Protection of Birds, Wrocławska 60A/7, 63-400 Ostrów Wielkopolski, Poland; e-mail: p.dolata@op.pl

**Abstract:** We surveyed the Mesostigmatid mite fauna of nests of the black stork *Ciconia nigra*, to determine the role of these mites in the biology of their hosts. We present preliminary results obtained on the basis of material collected from 31 nests. A total of 1,615 mite specimens was recorded, belonging to 39 species. The most abundant species were *Dendrolaelaps strenzkei*, *Apionoseius infirmus*, *Macrocheles merdarius* and *Macrocheles ancyleus*, which constituted more than 65% of all the specimens recorded. The presence of large numbers of predatory mites could be beneficial to the birds, if they feed on the eggs and larvae of the bird's parasites. It is likely that many of the mite species found in these nests were carried there by phoresy on insects, mainly Coleoptera.

**Key words:** mites; Acari; Mesostigmata; *Ciconia nigra*; nest fauna; Poland

### Introduction

Birds' nests constitute a specific, impermanent type of microhabitat that is inhabited by a unique assemblage of invertebrate fauna from a range of taxonomic groups. The most numerous among these are the arthropods, especially mites (Acari). The majority of studies on mites in this type of habitat have concentrated on the annual nests of small, mainly passerine birds (Błoszyk & Olszanowski 1985, 1986; Maśán & Krištofik 1995; Krištofik et al. 2001; Tryjanowski et al. 2001; Gwiazdowicz 2003a). The limited data available on the fauna of multi-annual nests of large species like birds of prey or wading birds, show that their mite assemblages are more uniform, characterized by specific species composition and structure (Philips 1981, 2000; Philips et al. 1983; Gwiazdowicz et al. 1999, 2000, 2005, 2006; Gwiazdowicz 2003b; Błoszyk et al. 2005, 2006; Bajerlein et al. 2006). The present paper adds to the available data by documenting the mites found in nests of the black stork *Ciconia nigra* (L., 1758) in Poland.

The black stork is a widely distributed species, breeding in at least 50 countries on three continents (Tamás et al. 2006). The breeding area covers central Europe and Asia, from the Atlantic to the Pacific, and a resident breeding population also exists in

South Africa. The total world population is estimated at 10,000–25,000 breeding pairs (Janssen et al. 2004; Tamás et al. 2006). The Polish population was recently estimated at 1,100–1,200 breeding pairs (Profus & Wójciak 2007). These numbers represent a significant recovery from estimates of 500–530 breeding pairs in 1966 (Bednorz 1974, probably underestimated) and 800–900 pairs in 1981–1982 (Keller & Profus 1992), as a result of legal and active protection.

In Poland, the black stork occupies its nest from the end of March or April to July or August, sometimes September (Zawadzka et al. 1990; Pugacewicz 1994, 1995; Stój 1995). It nests mainly in old deciduous or mixed forest, close to wet areas such as rivers, old river beds, streams, moors, swamps and fishponds (Bednorz 1974; Keller & Profus 1992; Profus & Wójciak 2007), where it obtains its main food, consisting of fish and amphibians (Zawadzka et al. 1990; Keller & Profus 1992; Kurowski et al. 1995; Hampl et al. 2007). In lowland areas of Poland, *C. nigra* nests mainly in oak trees, *Quercus robur* L. and *Q. sessilis* Ehrh., and in Scots Pine, *Pinus sylvestris* L. (Bednorz 1974; Cieślak 1988; Zawadzka et al. 1990; Keller & Profus 1992; Pugacewicz 1994, 1995; Kurowski et al. 1995; Kuźniak et al. 1999; Kołodzki et al. 2003). In the Polish mountains the main nesting trees are fir *Abies alba* Mill. and beech *Fagus*

*sylvatica* L. (Bednorz 1974; Stój 1995). It prefers old trees, older than 100 years (Pugacewicz 1994; Kołodzki et al. 2003), in which nests are built from 3 to 25 m above ground (Keller & Profus 1992; Pugacewicz 1994, 1995; Kurowski et al. 1995; Kołodzki et al. 2003, P. T. Dolata unpublished data).

Nests are built from sticks and twigs (Gotzman & Jabłoński 1972; Makatsch 1974; Snow et al. 1998; Janssen et al. 2004), which may be up to 3 cm thick (Janssen et al. 2004), brought from the ground or broken from the nest tree, and reinforced by earth and grass (Snow et al. 1998). Nests of this species differ from those of other species of large birds in Poland, because black stork nests are composed of successive layers of sticks, soil and wheat-grass, sticks, and so on (Gotzman & Jabłoński 1972). The shallow nesting niche inside the nest is typically lined with moss, but also sometimes with grass, leaves, animal hair, dung, hay, bark, reeds, and also sometimes artificial materials, including cloth, paper, string and plastic bags (Gotzman & Jabłoński 1972; Makatsch 1974; Snow et al. 1998; Janssen et al. 2004). Nests are large. In Latvia Strazds (2003) reported average nest dimensions of 115 × 111 × 49 cm. The oldest nest was 170 × 155 × 115 cm, and was estimated to exceed one tonne in weight. Nests may be occupied for 40–50 seasons, sometimes with breaks (Bednorz 1974; Pugacewicz 1995). In Latvia Strazds (2003) reported nests lasting 11–40 years, and the longest uninterrupted period of successful breeding in one nest was 11 seasons.

The nests of the black stork form a favourable microhabitat for mites, but their mite fauna has never been studied. Microenvironmental conditions in these nests are subject to cyclic annual changes determined by the biology of the birds. We can identify two periods during the annual cycle of events in these nests during which the nest microclimate is determined by different factors. The first is the breeding season in which the birds are present, and the microclimate is strongly influenced by the birds' behaviour, and the second is the time when the nest is empty and the microclimate is influenced mainly by ambient atmospheric conditions. During the breeding season the presence of the birds makes the nutritional and microclimatic conditions in the nest more favourable for the mite fauna, so the species richness during this period should be the greatest. The results presented here refer to the mite fauna of black stork nests during the birds' breeding season.

The mites occurring in birds' nests may be either harmful, neutral, or beneficial to their host and its clutch. Some of the mite species may be parasites or vectors of pathogenic microorganisms. On the other hand, other species feed on the eggs and larvae of these parasites, and could be beneficial to the birds. The structure of the mite community might therefore provide new insights into the biology and breeding ecology of the bird. Furthermore, comparison of mite assemblages from multi-annual nests of different, closely related bird species can show the differences in environmental conditions occurring in those microhabitats.

The purpose of this study was to determine the species composition of the Mesostigmata fauna in black stork nests, and specifically, to find which species are dominant, most frequent and characteristic of this type of microhabitat. We then compared these results with those recorded from the nests of the white stork *Ciconia ciconia* (L., 1758), which is a member of the same genus as the black stork, but which shows important differences in biology and behaviour, including in nest building.

## Material and methods

Thirty-one samples of nest material were collected from 27 nests of *Ciconia nigra* in June to July in 2004 and 2005, in the south Wielkopolska Region of Poland (51°25'–52°10' N, 17°10'–18°05' E), mainly in Antonin, Jarocin, Krotoszyn and Taczanów Forest Inspectorates (Jarocin, Ostrów Wielkopolski, Ostrzeszów and Środa Wielkopolska districts). Each sample consisted of 250–300 grams of nest padding material, collected from the middle of nests during the ringing of fledglings. Living mites were extracted from samples of nest material in Tullgren funnels for 48 hours, and collected into 75% ethanol. Temporary slide preparations were made in lactophenol for microscopic examination. Permanent mounts were made in PVA for selected specimens, or when identification of species was difficult. Mite specimens are deposited at the Invertebrate Databank Collection in the Department of Animal Taxonomy and Ecology of Adam Mickiewicz University, Poznań and the Department of Forest Protection in the University of Life Sciences in Poznań. The zoocenological analysis was conducted using two indices: occurrence (C%) and domination (D%) coefficients, as for previous papers in this series.

## Results

A total of 1,614 specimens of Mesostigmata was found in the collected material, belonging to 29 species in the suborder Gamasina and 10 species in the suborder Uropodina (Table 1). Below we present a short account of the biology of the species that were most frequent and most abundant in the studied nests, with special consideration of their microhabitat preference and geographical distribution. The seven species listed below comprise 87% of all the mites collected.

*Dendrolaelaps strenzkei* Hirschmann, 1960 (Digamasellidae). This predatory species has been found in rotting wood, under bark, in the nests of *Formica rufa* L., 1761 ants, and also in soil, forest litter and compost (Hirschmann & Wiśniewski 1982; Karg 1993). It also inhabits nests of the white-tailed eagle *Haliaeetus albicilla* (L., 1758) (Gwiazdowicz et al. 2006). It occurs all over Europe, from Italy in the South, to Finland in the North, from Spain in the West, to Russia in the East.

*Macrocheles merdarius* (Berlese, 1889) (Macrocheilidae) is a cosmopolitan predatory species found in manure and decomposing plant material, such as compost, silage and hay, and is phoretic on dung beetles of the family Scarabaeidae (Bregotova & Koroleva 1960; Krauss 1970; Bregotova 1977b; Maśán 2003; Błoszyk et al. 2005). It is often found in dung, either fresh and wet

Table 1. Mite species occurring in the nests of *Ciconia nigra*.

Species	Total	F	M	DN	PN	LV	D%	C%
<b>Gamasina</b>								
<i>Dendrolaelaps strenzkei</i> Hirschmann, 1960	507	177	28	263	37	2	31.41	45.16
<i>Macrocheles merdarius</i> (Berlese, 1889)	151	146	5	0	0	0	9.36	29.03
<i>Macrocheles ancyleus</i> Krauss, 1970	126	89	35	2	0	0	7.81	51.61
<i>Dendrolaelaps longiusculus</i> (Leitner, 1949)	97	61	21	15	0	0	6.01	22.58
<i>Macrocheles glaber</i> (Müller, 1860)	92	41	51	0	0	0	5.70	19.35
<i>Proctolaelaps pygmaeus</i> (Müller, 1860)	81	78	3	0	0	0	5.02	32.26
<i>Androlaelaps casalis</i> (Berlese, 1887)	57	49	8	0	0	0	3.53	19.35
<i>Parasitus fimetorum</i> (Berlese, 1904)	37	11	5	21	0	0	2.29	12.90
<i>Macrocheles penicilliger</i> (Berlese, 1904)	21	19	2	0	0	0	1.30	6.45
<i>Hypoaspis brevipilis</i> Hirschmann, 1969	18	17	1	0	0	0	1.12	19.35
<i>Alliphis halleri</i> (G. & R. Canestrini, 1881)	11	10	1	0	0	0	0.68	6.45
<i>Hypoaspis praesternalis</i> Willmann, 1949	10	8	2	0	0	0	0.62	6.45
<i>Halolaelaps</i> sp.	9	5	2	2	0	0	0.56	6.45
<i>Dendrolaelaps fallax</i> (Leitner, 1949)	7	5	2	0	0	0	0.43	12.90
<i>Zercon triangularis</i> C.L. Koch, 1836	7	3	2	1	0	1	0.38	6.45
<i>Ameroseius macrochelae</i> Westerboer, 1963	6	5	0	1	0	0	0.37	9.68
<i>Parasitus coleoptratorum</i> (L., 1758)	4	0	0	4	0	0	0.25	3.23
<i>Hypoaspis aculeifer</i> (Canestrini, 1883)	3	0	0	3	0	0	0.19	6.45
<i>Asca aphidioides</i> (L., 1758)	3	3	0	0	0	0	0.19	3.23
<i>Dendrolaelaps arvicolus</i> (Leitner, 1949)	3	3	0	0	0	0	0.19	3.23
<i>Hypoaspis giffordi</i> Evans & Till, 1966	3	3	0	0	0	0	0.19	3.23
<i>Gamasodes spiniger</i> (Trägårdh, 1910)	2	1	0	1	0	0	0.12	3.23
<i>Veigaia nemorensis</i> (C.L. Koch, 1839)	2	1	0	1	0	0	0.12	3.23
<i>Amblyseius</i> sp.	1	1	0	0	0	0	0.06	3.23
<i>Gamasellodes bicolor</i> (Berlese, 1918)	1	1	0	0	0	0	0.06	3.23
<i>Pergamasus mediocris</i> Berlese, 1904	1	1	0	0	0	0	0.06	3.23
<i>Dendrolaelaps cornutus</i> (Kramer, 1886)	1	1	0	0	0	0	0.06	3.23
<i>Hypoaspis heselhausi</i> Oudemans, 1912	1	1	0	0	0	0	0.06	3.23
<i>Zercon peltatus peltatus</i> C.L. Koch, 1836	1	0	1	0	0	0	0.06	3.23
Gamasina total	1263	740	169	314	37	3	78.25	
<b>Uropodina</b>								
<i>Apionoseius infirmus</i> (Berlese, 1887)	289	78	151	69	0	0	18.19	41.94
<i>Leiodinychus orbicularis</i> (C.L. Koch, 1839)	36	13	4	10	9	0	2.27	3.23
<i>Oodinychus ovalis</i> (C.L. Koch, 1839)	10	3	6	1	0	0	0.63	22.58
<i>Trachytes aegrota</i> (C.L. Koch, 1841)	7	2	2	3	0	0	0.44	6.45
<i>Uroobovella pyriformis</i> (Berlese, 1920)	3	1	1	1	0	0	0.19	6.45
<i>Nenteria breviunguiculata</i> (Willmann, 1949)	2	0	2	0	0	0	0.13	6.45
<i>Janetiella pulchella</i> (Berlese, 1904)	1	1	0	0	0	0	0.06	3.23
<i>Uroobovella marginata</i> (C.L. Koch, 1839)	1	0	0	1	0	0	0.06	3.23
<i>Uroobovella fimicola</i> (Berlese, 1903)	1	0	1	0	0	0	0.06	3.23
<i>Uropoda orbicularis</i> (Müller, 1776)	1	0	0	1	0	0	0.06	3.23
Uropodina total	351	98	167	86	9	0	21.75	
Mesostigmata total	1614	838	336	400	46	3	100.00	

Explanations: F – female; M – male; DN – deutonymph; PN – protonymph; LV – larva; D% – dominance coefficient; C% – occurrence coefficient.

or old and dry, and is also found in bird nests, including those of the common blackbird *Turdus merula* L., 1758 (Mašán 2003), and the white stork (Błoszyk et al. 2005; Bajerlein et al. 2006).

*Macrocheles ancyleus* Krauss, 1970 (Macrochelidae) was described from a rotting log, but appears to have a strong affinity for bird nests. It has often been collected in the nests of birds of prey (Gwiazdowicz et al. 1999, 2000, 2005, 2006; Mašán 2003), and also in guano in a cave (Cicolani 1983; Mašán & Zubáčová 2001). It is a predatory species known only from Europe.

*Dendrolaelaps longiusculus* (Leitner, 1949) (Digamasellidae). This predatory European mite species has been recorded in manure, compost and decomposing litter (Hirschmann & Wiśniewski 1982) and is also found

in bird nests, e.g., white stork (Błoszyk et al. 2005; Bajerlein et al. 2006) and the white-tailed eagle (Gwiazdowicz et al. 2005, 2006).

*Macrocheles glaber* (Müller, 1860) (Macrochelidae). This predatory species is found in decomposing organic matter, especially compost and dung (Krauss 1970). It has been found in the nests of birds, e.g., *Accipiter gentilis* (L., 1758), *Acrocephalus arundinaceus* (L., 1758), *Anser anser* (L., 1758), *Ciconia ciconia* (L., 1758), *Cygnus olor* (Gmelin, 1803), *Haliaeetus albicilla* (L., 1758), *Larus ridibundus* L., 1766, *Merops apiaster* L., 1758, *Nycticorax nycticorax* (L., 1758), *Parus major* L., 1758, *Parus montanus* Conrad von Baldenstein, 1827, *Passer montanus* (L., 1758), *Remiz pendulinus* (L., 1758), *Vanellus vanellus* (L., 1758) (Mašán 2003; Gwiazdowicz et al. 2005, 2006; Błoszyk et al. 2006; Ba-

jerlein et al. 2006). It is distributed from lowlands up to the mountain zone (1,330 m a.s.l.), and occurs in Europe, Asia, North America and Australia. It is carried phoretically by insects including several genera of scarabaeid beetles, and by flies (Bregetova & Koroleva 1960; Bregetova 1977b; Mařán 2003).

*Proctolaelaps pygmaeus* (Müller, 1860) (Ascidae) is a cosmopolitan species that occurs in soil, moss, decomposing organic matter and in the nests of small mammals, and most probably feeds on other microscopic invertebrates (Bregetova 1977a). It has been found in the nests of the white-tailed eagle and the white stork (Błoszyk et al. 2005; Gwiazdowicz et al. 2005, 2006; Bajerlein et al. 2006).

*Androlaelaps casalis* (Berlese, 1887) (Laelapidae) is found in forest litter, humus, and soil, but most commonly in the nests of small mammals and birds (Karg 1993). It occurs in the entire Palaearctic region and North America, and is probably cosmopolitan. In Poland it has been found in the nests of the white-tailed eagle and the white stork (Błoszyk et al. 2005; Gwiazdowicz et al. 2005, 2006; Bajerlein et al. 2006). It feeds on other mites and the eggs and larvae of insects (Hughes 1976).

*Apionoseius infirmus* (Berlese, 1887) (Trachytiidae). This species is found in decomposing plant remains, tree trunks and tree holes in Europe and Asia. It is commonly found in birds' nests, and is phoretic on beetles (Wiśniewski & Hirschmann 1993; Mařán 2001; Błoszyk et al. 2005, 2006; Bajerlein et al. 2006; Gwiazdowicz et al. 2006).

## Discussion

The most frequent species of Gamasina in the nests was *Macrocheles ancyleus* and the most abundant species was *Dendrolaelaps strenzkei*. Among the Uropodina the most frequent and abundant species was *Apionoseius infirmus*. In total, ten species of Gamasina and only two of Uropodina had occurrence coefficients of greater than 10%. The remaining species occur in the nests of the black stork only incidentally. The two most abundant species, *D. strenzkei* and *A. infirmus*, constituted almost 50% of the mites found, a phenomenon that is often observed in the mite communities of temporary microhabitats (Błoszyk et al. 2005). A further group included species with dominance coefficients between 5 and 10% and occurrence coefficients between 19 and 52%. These were *Macrocheles merdarius*, *Macrocheles ancyleus*, *Dendrolaelaps longiusculus*, *Macrocheles glaber* and *Proctolaelaps pygmaeus*. Those five species constituted 33.9% of the mites found. Among the Uropodina only one species had a high occurrence coefficient, *Oodinychus ovalis* (C.L. Koch, 1839), which was recorded in 23% of the nests sampled.

Most of the specimens recorded during the present study were adults, and females predominated in the majority of species. A few species, especially of Parasitidae, showed a high frequency of deutonymphs, as we also found in the nests of the white stork (Błoszyk

et al. 2005). This observation is somewhat puzzling – one should expect the r-strategy to predominate in species inhabiting unstable habitats, and juvenile specimens should dominate the whole fauna. The explanation is quite straightforward – most of the species recorded in the nests sampled are typically found in rotting wood and decomposing plant material, and multi-annual nests of the stork consist of rotting twigs and sticks. The K-type reproductive strategy is typical for most of those species when recorded outside the nests (Błoszyk 1999; Błoszyk et al. 2003). Since the nests constitute only a marginal habitat for their mite inhabitants, these mites have had no time or reason to evolve a new reproductive strategy.

This explanation also holds true for a second interesting observation. The community of Uropodina found in these nests comprises mainly bisexual species, in contrast with soil communities, where female-only populations are common, but similarly to those found in decomposing plant material (Błoszyk et al. 2003). Of the species observed in this study, only *Uropoda orbicularis* (O.F. Müller, 1776) and *Uroobovella pulchella* (Berlese, 1904) might be described as thelytokous. However, even in these cases occasional males have been found, and these species are facultatively parthenogenetic (Faasch 1967; Błoszyk et al. 2006).

The mite fauna of the nests of the black stork is strikingly different from that of the nests of white stork, which we have described previously (Błoszyk et al. 2005; Bajerlein et al. 2006). The sampling methods used for the two bird species were not always identical, so a rigorous comparison of the results is not possible. However, some general trends can be seen. From a total of 61 species of Mesostigmata, only 17 species were common to both stork species, and the species similarity index (S) amounted to slightly less than 31%. Also the mean abundances of mites in the nests of the two species differed significantly (Mann-Whitney *U* test:  $Z = 3.2$ ;  $P < 0.01$ ; Fig. 1). The most abundant species in the nests of *C. nigra*, *Dendrolaelaps strenzkei*, was not found at all in the nests of *C. ciconia*, and the second most abundant species in the nest of *C. ciconia*, *Macrocheles robustulus* (Berlese, 1904), was absent from the nests of *C. nigra*. At a generic level, *Macrocheles* made up 71% of the mites from *C. ciconia* but only 24% from *C. nigra*, and *Dendrolaelaps* made up 38% of the mites from *C. nigra* but less than 1% from *C. ciconia*. Possible reasons for these differences may be found in the different construction materials used by the two species of birds, differences in the birds' behaviour, and the different habitats in which their nests are built. The interior lining of the nest of *C. nigra* is made of tightly compressed moss, which remains humid and can even support a pool of water, while that of *C. ciconia* is more open in structure, and less likely to retain water. Also, *C. ciconia* nests in exposed synanthropic situations such as buildings and power transmission pylons, while in Poland, the nests of *C. nigra* are built in heavily shaded sites in forest trees (Pugacewicz 1995), where the microclimate is likely to be much more cool and




Fig. 1. Box-plot comparing the number of mite specimens recorded in samples from the nests of both stork species.

humid. We have previously reported that *C. ciconia* brings cow dung and compost into its nests (Bajerlein et al. 2006), and this is likely to introduce coprophilous mite species into the nests, such as *Macrocheles*. These species are likely to be further favoured by the presence of pellets of undigested food in the nests of *C. ciconia* (skin and bones of small mammals and amphibians). *Ciconia nigra* rarely brings dung or compost into its nests, since it feeds mainly in water, unlike *C. ciconia*, which feeds in meadows. Dung or compost was never observed in the nests studied here. Also, *C. nigra* does not leave undigested food in its nests.

The nests that we examined contained large numbers of predatory mites, dominated by the genera *Dendrolaelaps* and *Macrocheles*, which raises the question of what these mites were feeding on. The black stork and other species of *Ciconia* are attacked by a variety of ectoparasites, including lice (Ezealor 1985; Martin Mateo 1988; Zlotorzyczna 1990; Lanzarot et al. 2005; Dik & Uslu 2006), parasitic Astigmata (Fain & Lawrence 1986; Chen & Fan 2003), quill mites (Bochkov & Mironov 1999), ticks (Ezealor 1985; Dik & Uslu 2006) and feather mites (Perez & Atyeo 1992; Janssen et al. 2004). It is possible that predatory mites play a role in nest hygiene by feeding on these parasites. However, parasitic Mesostigmata were not found in the nests we examined. This is consistent with the observations of Lanzarot et al. (2005), who found lice on 11 of the 50 nestlings of *C. nigra* that they examined in Spain, but no other parasites. A near or complete absence of blood-feeding parasitic Mesostigmata has also been reported for nests of the white-tailed eagle (Gwiazdowicz et al. 2005, 2006; Fenda & Lengyel 2007), and the white stork (Bloszyk et al. 2005; Bajerlein et al. 2006). In contrast, large populations of parasitic mites, such as *Dermanyssus hirudinis* (Hermann, 1804), *D. gallinae* (De Geer, 1778), *Ornithonyssus bursa* (Berlese,

1888), *O. sylviarum* (Canestrini et Fanzago, 1877) have commonly been found in the nests of passerine birds (Krištofik et al. 1993, 2001, 2002; Mašan & Krištofik 1993; Gwiazdowicz 2003a). Even more surprising is the fact that the parasite *Dermanyssus gallinae* was the most abundant mite species found in the nests of boreal owl *Aegolius funereus* (L., 1758) in Bohemia by Krištofik et al. (2003), but was absent from nests of the same owl species examined by Philips (1981) in Norway. Several families of saprophytic mites in the suborder Astigmata have been reported from birds' nests (Philips 1981; OConnor 1979), and these are a further possible type of prey for predatory mites. However, in the present study and others, Astigmata were not found. Inconsistencies in the occurrence of Astigmata and parasitic Mesostigmata in the nests of different bird species are difficult to explain, but highlight the fact that the ecology of mites in birds' nests is not yet fully understood.

Apart from being carried with nest material, phoresy on coprophilous insects could be an important mode of migration of mites to and from nests (Faasch 1967). Some of the Mesostigmata species recorded in this study, including *Macrocheles merdarius*, *M. glaber*, *Alliphis halleri* (G. et R. Canestrini, 1881) and *Proctolaelaps pygmaeus*, are phoretic on many species of dung beetles. Some informal observations (J. Bloszyk, unpubl. data) indicate that at least 20 other species reported in the present paper can adopt the same means of dispersal. That would be consistent with previous observations conducted on mite fauna of multi-annual nests of other bird species (Bloszyk et al. 2005, 2006; Gwiazdowicz et al. 2005, 2006; Bajerlein et al. 2006).

We can identify a further possible route by which mites may enter the nests of *C. nigra*, through the activities of other species of birds. Nests of *C. nigra* from the previous breeding season are sometimes occupied by birds of prey in the next breeding season, for example the common buzzard *Buteo buteo* (L., 1758), the Northern goshawk *Accipiter gentilis*, and the lesser spotted eagle *Aquila pomarina* C.L. Brehm, 1831 (Bednorz 1974; Pugacewicz 1995; Snow et al. 1998; P.T. Dolata unpublished data). The effects of these birds on the mite fauna are likely to be variable and unpredictable, and introduce a further unknown level of heterogeneity into the data presented here.

#### Acknowledgements

We are very grateful to Voivodship Nature Conservator in Poznań and the Ministry of the Environment for permission to study black stork nests. The field work was made possible by T. Mizera, the team of South Wielkopolska Group of the Polish Society for the Protection of Birds (PwG OTOP), especially T. Ekiert, J. Pietrowiak and J. Pruchniewicz and foresters from Antonin, Jarocin, Krotoszyn and Taczanów Forest Inspectorates. The research was partially supported by the Voivodeship Fund for Environmental Protection and Water Management in Poznań and grant No. 512 00 066.

## References

- Bajerlein D., Błoszyk J., Gwiazdowicz D.J., Ptaszyk J. & Halliday B. 2006. Community structure and dispersal of mites (Acari, Mesostigmata) in nests of the white stork (*Ciconia ciconia*). *Biologia* **61**: 525–530. DOI 10.2478/s11756-006-0086-9
- Bednorz J. 1974. Bocian czarny, *Ciconia nigra* (L.) w Polsce [Black stork, *Ciconia nigra* (L.) in Poland]. *Ochrona Przyrody* **39**: 201–243.
- Błoszyk J. 1999. Geograficzne i ekologiczne zróżnicowanie zgrupowań roztoczy z kohorty Uropodina (Acari: Mesostigmata) w Polsce. I. Uropodina lasów grądowych (*Carpinion betuli*) [Geographical and ecological variability of mites of the cohort Uropodina (Acari: Mesostigmata) in Poland. I. Uropodine mites of oak-hornbeam forest (*Carpinion betuli*)]. Wydawnictwo Kontekst Poznań, 245 pp.
- Błoszyk J., Bajaczyk R., Markowicz M. & Gulvik M. 2003. Geographical and ecological variability of mites of the suborder Uropodina (Acari: Mesostigmata) in Europe. *Biol. Lett.* **40**: 15–35.
- Błoszyk J., Bajerlein D., Gwiazdowicz D.J., Halliday R.B. & Dylewska M. 2006. Uropodine mite communities (Acari: Mesostigmata) in birds' nests in Poland. *Belg. J. Zool.* **136**: 145–153.
- Błoszyk J., Gwiazdowicz D.J., Bajerlein D. & Halliday R.B. 2005. Nests of the white stork *Ciconia ciconia* (L.) as a habitat for mesostigmatic mites (Acari, Mesostigmata). *Acta Parasitol.* **50**: 171–175.
- Błoszyk J. & Olszanowski Z. 1985. Materiały do znajomości roztoczy gniazd i budek legowych ptaków. I. Uropodina i Nothoidea (Acari: Mesostigmata et Oribatida) [Materials to the knowledge of the mites of birds nests. I. Uropodina and Nothoidea (Acari: Mesostigmata and Oribatida)]. *Przegląd Zoologiczny* **29**: 69–74.
- Błoszyk J. & Olszanowski Z. 1986. Materiały do znajomości fauny roztoczy gniazd i budek legowych ptaków. II. Różnice w liczebności i składzie gatunkowym populacji Uropodina (Acari: Anactodotrichida) budek legowych na Mierzei Wiślanej na podstawie dwuletnich obserwacji [Materials to the knowledge of the mites of birds nests. II. Numerical and species compositional differences of Uropodina (Acari, Anactinotrichida) of birds nests in the sandbar "Mierzeja Wiśłana" on the base of biennial observation]. *Przegląd Zoologiczny* **30**: 63–66.
- Bochkov A.V. & Mironov S.V. 1999. New quill mite species of the family Syringophilidae (Acari: Cheyletoidea) from the European part of Russia. *Acarina* **7**: 35–45.
- Bregetova N.G. 1977a. Family Aceosejidae Baker et Wharton, 1952 (sensu Evans, 1958), pp. 169–226. In: Ghilyarov M.S. & Bregetova N.G. (eds), Key to the Soil Inhabiting Mites. Mesostigmata, Nauka, Leningrad.
- Bregetova N.G. 1977b. Macrochelidae, pp. 346–411. In: Ghilyarov M.S. & Bregetova N.G. (eds), Key to the Soil Inhabiting Mites. Mesostigmata, Nauka, Leningrad.
- Bregetova N.G. & Koroleva E.V. 1960. Mites of the family Macrochelidae Vitzthum, 1930 in the fauna of the SSSR. *Parazit. Sb.* **19**: 32–154.
- Chen F. & Fan H. 2003. The diagnosis and treatment of *Laminosioptes cysticola* disease in *Ciconia nigra*. *Chin. J. Vet. Sci. Technol.* **33**: 66–67.
- Cicolani B. 1983. Una nuova specie di *Macrocheles* (Acarina: Macrochelidae) raccolta nella grotta della Zinzulusa (Castro Marina-Puglie). *Boll. Mus. Civ. St. Nat. Verona* **9**: 151–159.
- Cieślak M. 1988. Gniazda bociana czarnego (*Ciconia nigra*) w Lasach Janowskich, woj. tarnobrzeskie [Nest sites of Black Stork (*Ciconia nigra*) in Lasy Janowskie, prov. Tarnobrzeg]. *Notatki Ornitologiczne* **29**: 227–231.
- Dik B. & Uslu U. 2006. Mallophaga (Insecta) species occurring on storks (*Ciconia ciconia* Linnaeus, 1758). *Türkiye Parazitoloji Dergisi* **30**: 220–225.
- Ezealor A.U. 1985. Parasites and diseases of Abdim's Stork *Ciconia abdimii*. *Malimbus* **7**: 120.
- Faasch H. 1967. Beitrag zur Biologie der einheimischen uropodiden *Uroobovella marginata* (C.L. Koch, 1839) und *Uropoda orbicularis* (O.F. Müller, 1776) und experimentelle Analyse ihres Phoresieverhaltens. *Zoologische Jahrbücher Abteilung für Systematik, Ökologie und Geographie der Tiere* **94**: 521–608.
- Fain A. & Laurence B.R. 1986. Two species of *Neottialges* Fain (Acari, Hypoderatidae) under the skin of birds, with a key to the hypopi of this genus. *J. Nat. Hist.* **20**: 849–856. DOI 10.1080/00222938600770621
- Fenda P. & Lengyel J. 2007. Roztočce (Acarina, Mesostigmata) v hniezdach orliaka morského (*Haliaeetus albicilla*) na Slovensku [Mites (Acarina, Mesostigmata) in nests of the white-tailed sea eagle *Haliaeetus albicilla* (L.) in Slovakia]. *Entomofauna Carpathica* **19**: 48–50.
- Gotzman J. & Jabłoński B. 1972. Gniazda naszych ptaków [Nests of our birds]. PZWS, Warszawa, 282 pp.
- Gwiazdowicz D.J. 2003a. Mites (Acari, Mesostigmata) appearing in Poland, in the bird's nests of Passeriformes, Falconiformes and Strigiformes orders, pp. 562–572. In: Miler, A.T. (ed.), Kształtowanie i ochrona środowiska leśnego, Wyd. Akad. Rol., Poznań.
- Gwiazdowicz D.J. 2003b. Mites (Acari: Mesostigmata) occurring in the nests of birds of prey (Falconiformes) and owls (Strigiformes). *Acarina* **11**: 235–239.
- Gwiazdowicz D.J., Błoszyk J., Bajerlein D., Halliday R.B. & Mizera T. 2006. Mites (Acari: Mesostigmata) inhabiting nests of the white-tailed sea eagle *Haliaeetus albicilla* (L.) in Poland. *Entomol. Fenn.* **8**: 366–372.
- Gwiazdowicz D.J., Błoszyk J., Mizera T. & Tryjanowski P. 2005. Mesostigmatic mites (Acari: Mesostigmata) in white-tailed sea eagle nests (*Haliaeetus albicilla*). *J. Raptor Res.* **39**: 60–65.
- Gwiazdowicz D.J., Mizera T. & Skorupski M. 1999. Mites in greater spotted eagle nests. *J. Raptor Res.* **33**: 257–260.
- Gwiazdowicz D.J., Mizera T. & Skorupski M. 2000. Mites (Acari, Gamasida) from the nests of birds of prey in Poland. *Buteo* **11**: 97–100.
- Hامل R., Beran V. & Dolata P.T. 2007. Potrava mláďat čápa černého (*Ciconia nigra*) v České republice a v Polsku [Nestling diet of the black stork *Ciconia nigra* in the Czech Republic and Poland]. *Sylvia* **43**: 165–172.
- Hirschmann W. & Wiśniewski J. 1982. Weltweite Revision der Gattungen *Dendrolaelaps* Halbert 1915 und *Longoseius* Chant 1961 (Parasitiformes). *Acarologie. Schriftenreihe für Vergleichende Milbenkunde* **29**: 1–190.
- Hughes A.M. 1976. The Mites of Stored Food and Houses. Ministry of Agriculture, Fisheries and Food, Technical Bulletin no. 9 (Second edition), Her Majesty's Stationery Office, London, 400 pp.
- Janssen G., Hormann M. & Rohde C. 2004. Der Schwarzstorch. Die Neue Brehm-Bücherei Bd. 468. Westarp Wissenschaften, Hohenwarsleben, 414 pp.
- Karg W. 1993. Acari (Acarina), Milben Parasitiformes (Anactinochaeta), Cohort Gamasina Leach. *Raubmilben. Die Tierwelt Deutschlands* **59**, pp. 1–523.
- Keller M. & Profus P. 1992. Present situation, reproduction and food of the Black Stork in Poland, pp. 227–236. In: Meriaux J.-L., Schierer A., Tombal C. & Tombal J.-C. (eds), Les Cigognes d'Europe, Actes du Colloque International. Institut Européen d'Ecologie, Metz.
- Kołodzki Z., Wężyk M., Kociniak M. & Zieliński P. 2003. Występowanie bociana czarnego *Ciconia nigra* na obszarze byłego województwa piotrkowskiego w latach 1994–2001 [Occurrence of the Black Stork *Ciconia nigra* in the former Piotrków Voivodship in the years 1994–2001]. *Chrońmy Przyrodę Ojczystą* **59** (4): 5–19.
- Krauss W. 1970. Die europäischen Arten der Gattungen *Macrocheles* Latreille 1829 und *Geholaspis* Berlese 1918. *Acarologie. Schriftenreihe für Vergleichende Milbenkunde* **14**, pp. 1–60, Plates 1–20.
- Křištofik J., Mašán P. & Šustek Z. 2001. Mites (Acari), beetles (Coleoptera) and fleas (Siphonaptera) in the nests of great reed warbler (*Acrocephalus arundinaceus*) and reed warbler (*A. scirpaceus*). *Biologia* **56**: 525–536.

- Krištofik J., Mašán P., Šustek Z. & Gajdoš P. 1993. Arthropods in the nests of penduline tit (*Remiz pendulinus*). *Biologia* **48**: 493–505.
- Krištofik J., Mašán P., Šustek Z. & Kloubec B. 2003. Arthropods (Pseudoscorpionida, Acari, Coleoptera, Siphonaptera) in nests of the tengmalm's owl, *Aegolius funereus*. *Biologia* **58**: 231–240.
- Krištofik J., Šustek Z. & Mašán P. 2002. Arthropods (Pseudoscorpionida, Acari, Coleoptera, Siphonaptera) in the nests of red-backed shrike (*Lanius collurio*) and lesser grey shrike (*Lanius minor*). *Biologia* **57**: 603–613.
- Kurowski M., Konofalski M., Czuchnowski R. & Profus P. 1995. Stan populacji bociana czarnego *Ciconia nigra* na Ziemi Radomskiej w latach 1981–1994 [Population of the black stork *Ciconia nigra* in the district of Radom in 1981–1994]. *Chrońmy Przyrodę Ojczystą* **51** (2): 15–28.
- Kuźniak S., Lorek G. & Lewandowski M. 1999. Stan populacji bociana czarnego *Ciconia nigra* i jego ochrona na Ziemi Leszczyńskiej [The status and protection of the black stork *Ciconia nigra* in the district of Leszno]. *Chrońmy Przyrodę Ojczystą* **55** (3): 5–18.
- Lanzarot M.P., Martín-Mateo M.P., Merino S. & Fernández-García M. 2005. A redescription of *Neophiloaterus tricolor* (Burmeister, 1838) (Insecta: Phthiraptera: Ischnocera: Philopteridae) from the black stork *Ciconia nigra* (L.) (Aves) with notes on its prevalence. *J. Nat. Hist.* **39**: 3471–3480. DOI 10.1080/00222930500392766
- Makatsch W. 1976. Die Eier der Vögel Europas. Eine Darstellung der Brutbiologie aller in Europa brütenden Vogelarten. Bd. 2. Neumann-Neudamm Verlag, Melsungen, 322 pp.
- Martin Mateo M.P. 1988. Malófagos (Insecta) parásitos de cigüeña común, *Ciconia ciconia* (L.) (Aves). *Eos* **64**: 147–158.
- Mašán P. 2001. Mites of the Cohort Uropodina (Acarina, Mesostigmata) in Slovakia. *Annotationes Zoologicae et Botanicae* **223**: 1–320.
- Mašán P. 2003. Macrochelid mites of Slovakia (Acari, Mesostigmata, Macrochelidae). Institute of Zoology, Slovak Academy of Sciences, Bratislava, 149 pp.
- Mašán P. & Krištofik J. 1993. Mites and ticks (Acarina: Mesostigmata et Ixodida) from the nests of *Riparia riparia* L. in South Slovakia. *Biologia* **48**: 155–162.
- Mašán P. & Krištofik J. 1995. Mesostigmatid mites (Acarina: Mesostigmata) in the nests of penduline tit (*Remiz pendulinus*). *Biologia* **50**: 481–485.
- Mašán P. & Zubáčová Z. 2001. First records of macrochelid mites (Acari, Macrochelidae) in Slovakia. *Biologia* **56**: 577–578.
- O'Connor B.M. 1979. Evolutionary origins of Astigmatid mites inhabiting stored products, pp. 273–278. In: Rodriguez J.G. (ed.), *Recent Advances in Acarology*, Vol. 1, Academic Press, New York.
- Perez T.M. & Atyeo W.T. 1992. A review of the Xoloptoidinae (Acari, Pterolichidae) and the description of a new genus. *Entomologische Mitteilungen aus dem Zoologischen Museum Hamburg* **10**: 209–219.
- Philips J.R. 1981. Mites (Acarina) from nest of Norwegian birds of prey. *Fauna Norvegica Series B* **28**: 44–47.
- Philips J.R. 2000. A review and checklist of the parasitic mites (Acarina) of the Falconiformes and Strigiformes. *J. Raptor Res.* **34**: 210–231.
- Philips J.R., Root M. & DeSimone P. 1983. Arthropods from a saw-whet owl (*Aegolius acadicus*) nest in Connecticut. *Entomol. News* **94**: 60–64.
- Profus P. & Wójciak J. 2007. Bocian czarny *Ciconia nigra* [Black Stork *Ciconia nigra*], pp. 126–127. In: Sikora A., Rohde Z., Gromadzki M., Neubauer G. & Chylarecki P. (eds), *Atlas rozmieszczenia ptaków lęgowych Polski 1985–2004* [The atlas of breeding birds in Poland 1985–2004], Bogucki Wyd. Nauk, Poznań.
- Pugacewicz E. 1994. Stan populacji bociana czarnego (*Ciconia nigra*) na Nizinie Północnopodlaskiej w latach 1985–1994 [Population of the Black Stork (*Ciconia nigra*) in the Nizina Północnopodlaska Lowlands in 1985–1994]. *Notatki Ornitologiczne* **35**: 297–308.
- Pugacewicz E. 1995. Populacja bociana czarnego (*Ciconia nigra*) w polskiej części Puszczy Białowieskiej [Population of the Black Stork (*Ciconia nigra*) in the Polish Part of Białowieża Forest]. *Ptaki Północnego Podlasia* **1**: 1–25.
- Snow D.W., Perrins C.M., Doherty P. & Cramp S. 1998. *The Complete Birds of the Western Palearctic on CD-ROM*. Oxford University Press, Oxford.
- Stój M. 1995. Ekologia rozrodu bociana czarnego *Ciconia nigra* w Beskidzie Niskim i okolicach Jasła [Reproduction ecology of the Black Stork in Bekid Niski and in the environs of Jasło]. *Chrońmy Przyrodę Ojczystą* **51** (2): 29–39.
- Strazds M. 2003. Longevity of Black Stork (*Ciconia nigra*) nests and nest site protection in Latvia. *Aves* **40** (1–4): 69–71.
- Tamás E.A., Kalocsa B. & Strazds M. 2006. Foreword. *Biota* **7**: 5–6.
- Tryjanowski P., Baraniak E., Bajaczyk R., Gwiazdowicz D.J., Konwerski S., Olszanowski Z. & Szymkowiak P. 2001. Arthropods in nests of the red-backed shrike (*Lanius collurio*) in Poland. *Belg. J. Zool.* **131**: 69–74.
- Wiśniewski J. & Hirschmann W. 1993. Gangsystematik der Parasitiformes. Teil 548. Katalog der Ganggattungen, Untergattungen, Gruppen und Arten der Uropodiden der Erde (Taxonomie, Literatur, Grösse, Verbreitung, Vorkommen). *Acarologie. Schriftenreihe für Vergleichende Milbenkunde* **40**, pp. 1–220.
- Zawadzka D., Olech B. & Zawadzki J. 1990. Zagęszczenie, rozród i pokarm bociana czarnego (*Ciconia nigra*) w Kampinoskim Parku Narodowym w latach 1979–1987 [Population density, reproduction and food of the black stork in the Kampinoski National Park in years 1979–1987]. *Notatki Ornitologiczne* **31**: 5–20.
- Złotorzycka J. 1990. Pasożytnicze stawonogi Arthropoda parasitica. Katalog Fauny Pasożytniczej Polski. *Catalogus faunae parasiticae poloniae* [Catalogue of Polish parasite fauna. Bird parasites. Parasitic arthropods]. *Katalog Fauny Pasożytniczej Polski* **4** (3), pp. 1–366.

Received October 24, 2008

Accepted January 15, 2009