

Short Conceptual Overview

Long telomeres: too much of a good thing

Michael Chang

European Research Institute for the Biology of Ageing,
University Medical Centre Groningen, University of
Groningen, A. Deusinglaan 1, NL-9713 AV Groningen,
The Netherlands

e-mail: m.chang@umcg.nl

Abstract

Telomeres, the physical ends of linear eukaryotic chromosomes, protect chromosome ends from end fusions and degradation. Telomere length is tightly regulated to ensure that telomeres are neither too short nor too long. Short telomeres are preferentially elongated by the enzyme telomerase. In the absence of telomerase, telomeres progressively shorten with each round of cell division. Critically shortened telomeres lose their ability to protect chromosome ends, inducing cell cycle arrest and senescence. While the consequences and cellular response to short telomeres are frequently explored, long telomeres also pose problems and cells have evolved mechanisms to shorten over-elongated telomeres. These aspects of long telomeres are discussed in this short conceptual overview.

Keywords: telomerase; telomere length regulation; telomere rapid deletion; telomeres.

Introduction

Eukaryotic DNA is organized into linear chromosomes. Maintaining the genetic information encoded within the DNA is an essential biological process. The DNA in our cells is constantly being challenged, both by DNA-damaging agents and by normal DNA metabolism, and any damage to the DNA must be repaired to safeguard the integrity of the genome. Perhaps the most hazardous DNA lesion is a double-stranded DNA break (DSB). DSBs, created by mechanical stress or DNA-damaging agents, need to be recognized and accurately repaired (1). In contrast, natural chromosome ends must be shielded from repair activities. Failure to do so could lead to cell cycle arrest, end-to-end fusion events, and loss of genome integrity. To combat this problem, cells have evolved specialized proteins that bind to short, repetitive, G-rich sequences at chromosome ends, forming protective nucleoprotein complexes called telomeres (2). However, the canonical DNA replication machinery is unable to fully

replicate chromosome ends, resulting in telomere erosion with each round of cell division (3). In the vast majority of eukaryotes, telomere shortening is counteracted by a specialized reverse transcriptase called telomerase, whose core consists of a protein catalytic subunit and an RNA moiety, hTERT and hTR, respectively, in humans (4–6). Telomerase extends a telomere by repeated reverse transcription of a short sequence to the 3' end of the telomere, using the RNA subunit as a template (7–9). The DNA replication machinery that is responsible for lagging strand synthesis presumably fills in the complementary 5' strand. To ensure that telomeres are never in danger of becoming too short, telomerase preferentially extends short telomeres, an evolutionarily conserved feature of telomerase that has been observed in the budding yeast *Saccharomyces cerevisiae* (10), mice (11), and human fibroblasts expressing telomerase (12). Individuals born with reduced telomerase activity have short telomeres, which leads to telomere dysfunction in highly proliferative cells, and several human diseases are associated with shortened telomeres (13). Furthermore, critically shortened, dysfunctional telomeres are unstable and lead to chromosome end-to-end fusion events and genome instability, which can promote tumor progression (14, 15).

A review on telomeres typically includes a discussion on the consequences of harboring short telomeres, and a description of how cells recognize and extend short telomeres. However, telomere length is tightly regulated, not just to ensure that telomeres do not become too short, but also to prevent them from becoming over-elongated. If short telomeres can have such negative consequences, why are longer telomeres not evolutionarily selected? Can longer-than-normal telomeres have detrimental effects as well? This overview aims to highlight some of the important aspects of long telomeres.

Evolutionary considerations of long telomeres

Most human somatic cells do not express telomerase, resulting in progressive telomere shortening with each round of cell division (16). Extensively eroded telomeres trigger a DNA damage checkpoint response, which arrests cell cycle progression and causes cells to either die by apoptosis or enter a state known as replicative senescence (9, 17, 18). A recent report indicates that the presence of approximately five dysfunctional telomeres causes p53-dependent senescence in human cells (19). Senescence limits replicative potential and therefore has been proposed to be a cause of human aging, but it is also thought that replicative senescence is an important

barrier to tumorigenesis as cancer cells need to maintain their telomeres to continue proliferating. Thus, inheriting long telomeres may increase replicative potential, and perhaps life span, but it could result in increased cancer rates (20). However, this model seems unlikely as it has recently been noted that longer blood and epithelial cell telomere length is rarely associated with increased rates of cancer (21). On the other hand, short dysfunctional telomeres promote genome instability, which is a hallmark of cancer cells, and short telomeres are often linked to increased cancer risks (14, 15, 21). Furthermore, most cancers occur late in life when the force of selection pressure is reduced (21). Therefore, it is unlikely that individuals with long telomeres are selected against because of increased cancer rates.

An alternative explanation for keeping telomere lengths in check is the ‘thrifty telomere’ hypothesis, which suggests that long telomeres require more energy to maintain (21). In this model, telomeres should be kept as short as possible provided they are still fully functional. However, the evolutionary reasons for limiting telomere length are still far from being understood. Evolutionary models must be able to account for telomere length variation within a species as well as variation between species, and such models are often difficult to prove. Thus, instead of examining the strengths and weaknesses of these speculative models, this review will focus on the cellular and molecular consequences of long telomeres.

DNA replication stress at telomeres

Telomeric DNA sequences are highly repetitive and GC rich – features that are typically troublesome for the DNA replication machinery during DNA synthesis. Moreover, human telomeres are hypersensitive to UV-induced DNA damage (22), and sites of damage also cause problems for DNA polymerases. In *S. cerevisiae*, replication forks pause while traversing the telomeric repeats, and the strength of this pausing is proportional to telomere length (23, 24). Several proteins have been identified that promote telomeric DNA replication, suggesting that there may be multiple reasons for replication fork pausing at telomeres. The Rrm3 helicase promotes replication through the telomere, likely by facilitating replication past non-histone protein-DNA complexes (23, 25). In the fission yeast *Schizosaccharomyces pombe*, the telomere-binding protein Taz1 promotes DNA replication at telomeres (26). Similarly, mammalian telomeres also pose a challenge to the DNA replication machinery and require the Taz1-homolog TRF1 for efficient replication (27).

The telomeric DNA from most eukaryotic organisms can form G-quadruplex (G4) structures *in vitro* (28, 29). G4 DNA was first observed *in vivo* through studies using anti-G4 DNA-specific antibodies to detect such structures at ciliate telomeres (30, 31). In theory, the formation of G4 DNA structures should be problematic for a passing replication fork. In *S. cerevisiae*, the Pif1 helicase is needed to resolve G4 DNA, and in cells lacking Pif1, DNA replication is impeded and there is an increase in replication fork collapse (32). Other helicases, such as mammalian BLM, WRN, and RTEL, have

also been implicated in promoting telomeric replication through the removal of G4 DNA structures (27, 33, 34). Taz1 and TRF1 have both been suggested to promote telomeric replication by recruiting one or more of these helicases (26, 27).

Increasing the length of a telomere would obviously increase the number of potential barriers to efficient telomeric DNA synthesis, making long telomeres more difficult to fully replicate. Furthermore, although there is some evidence that DNA replication can initiate within the telomeric tracts (27, 35), the majority of telomere replication is accomplished by replication forks originating from subtelomeric regions (24, 27). Thus, increasing the length of a telomere would also increase the distance that the replication fork must travel to fully replicate the telomere, which may result in a prolonged S phase and disruption of cell cycle progression. Taken together, cells may not favor the presence of long telomeres due to problems associated with the telomere replication.

Effect of long telomeres on telomere-binding proteins

Telomeric repeats are bound by telomere-binding proteins, so long telomeres would recruit more of these proteins. If cells maintain longer-than-normal telomeres, sufficient quantities of these telomere-binding proteins must be present to ensure proper capping of the telomeres. Remarkably, both *S. cerevisiae* and human cells can be manipulated to have extremely elongated telomeres with no dramatic effect on cell viability, indicating that telomere-binding proteins are not easily titrated below a threshold needed to maintain essential telomere capping (36, 37).

However, many telomere-binding proteins have both telomeric and non-telomeric functions. For example, *S. cerevisiae* Rap1 binds telomeric repeat DNA and is important to establish telomere length homeostasis (38, 39), but it was first discovered as a protein that can modulate gene expression (40). Furthermore, the Rap1-associated proteins Sir2, Sir3, and Sir4 are important for repressing transcription near telomeres as well as at the silent mating type loci (41). Sir2 is also important for regulating replicative life span through its role at the ribosomal DNA (rDNA) repeats (42, 43). Increasing telomere length may sequester telomere-binding proteins at the telomere and away from non-telomeric sites. Indeed, it has been reported that long telomeres reduce life span by reducing the amount of Sir2 available at the rDNA repeats (44). Similarly, long telomeres increase telomeric silencing, but reduce silencing at the *HMR* mating type locus by sequestering the Sir proteins to telomeres (45, 46). Whether the sequestration of telomere-binding proteins at over-elongated telomeres significantly affects other aspects of cell biology is still largely unknown.

Shortening over-elongated telomeres

The previous sections described potential downsides of harboring long telomeres. So how do cells shorten over-

elongated telomeres? The most obvious mechanism is to limit telomerase-mediated telomere extension (Figure 1A). Short telomeres are preferentially elongated by telomerase in *S. cerevisiae* (10), mice (11), and human fibroblasts expressing telomerase (12). In other words, long telomeres are less likely to be extended by telomerase and will progressively shorten at a rate similar to when telomerase is absent. Indeed, in *S. cerevisiae*, an artificially over-elongated telomere shortens at a rate of ~3–4 base pairs (bp) per generation, and this rate is independent of the presence or absence of telomerase (47). The shortening rate is ~50–150 bp per generation in mouse cells lacking telomerase (48), and in a variety of human cell types with no detectable telomerase activity (18, 49–51).

The reason for this shortening is primarily due to a combination of incomplete DNA replication of chromosome ends and nucleolytic degradation. Incomplete DNA replication occurs because (i) DNA polymerases can only synthesize DNA in a 5' to 3' direction, and (ii) DNA polymerases cannot synthesize DNA *de novo* and require a short primer of ~10 nucleotides (nt) of RNA. On the leading strand, DNA polymerase can theoretically synthesize DNA until it reaches the end of the chromosome, producing a blunt end. However, lagging strand synthesis occurs discontinuously, with each fragment (or Okazaki fragment) beginning with a short RNA primer. The RNA primers must be removed and replaced with DNA, which is synthesized by DNA polymerase from an upstream Okazaki fragment. Removal of the RNA primer of the terminal Okazaki fragment leaves a gap of ~10 nt that cannot be replaced. Without a mechanism to compensate for this loss, chromosome ends will progressively shorten. This

phenomenon was termed the 'end-replication problem' (52, 53). In addition, to the end-replication problem, chromosome ends terminate with 3' overhangs, the size of which varies from species to species. Thus, nucleolytic degradation must occur, at least on the blunt-ended telomere synthesized by the leading strand (3). The combined effect of the end-replication problem and nucleolytic degradation is that telomeres shorten with each round of cell division.

DNA damage can also contribute to the shortening of telomeres (Figure 1B). For example, the accumulation of single-stranded breaks in telomeric DNA, induced by oxidative stress, is a major cause of telomere shortening in human fibroblasts (54, 55). Furthermore, as mentioned above, DNA replication forks have difficulty traversing telomeric DNA (23, 24, 26, 27). If a replication fork collapses before reaching the end of the telomere, there is no replication origin distal to the site of fork collapse to generate a fork to finish the replication of the telomere, resulting in a truncated telomere (Figure 1C). Evidence for such truncated telomeres has been observed in *S. cerevisiae* (56). Normally, if the truncated telomere is significantly shortened, the telomerase will preferentially elongate it (10). However, if the telomere was initially over-elongated, the truncation may still leave the telomere longer than wild-type length. Telomerase would not preferentially elongate such a telomere, and this might be used as a mechanism to shorten over-elongated telomeres.

While incomplete replication, nucleolytic degradation, DNA damage, and replication fork collapse can all shorten over-elongated telomeres, these mechanisms act at telomeres of all lengths. In contrast, over-elongated telomeres are also




Figure 1 Shortening over-elongated telomeres.

(A) In the absence of telomerase, or when telomerase activity is inhibited, telomeres shorten due to a combination of incomplete DNA replication and nucleolytic degradation. (B) DNA damage within the telomeric tracts can lead to telomere truncation events. (C) Replication forks pause while traversing telomeric repeats. The fork pausing can be induced by proteins bound to the telomere or, as depicted here, by the presence of G4 DNA. Collapse of a stalled replication fork can lead to a truncated telomere. (D) Over-elongated telomeres can undergo TRD events, which return the telomere to approximately wild-type length by excising an extrachromosomal DNA circle containing telomeric repeats.




Figure 2 Regulating telomere length via a protein-counting mechanism in *S. cerevisiae*.

(A) Telomerase activity is regulated by telomere length, which is determined by the amount of Rap1, Rif1, and Rif2 bound at a telomere. A short telomere recruits few Rap1/Rif complexes, which favors telomerase-mediated extension of the telomere. A long telomere associates with many Rap1/Rif complexes, which inhibits telomerase activity. (B) While telomere-bound Rap1/Rif complexes regulate telomerase activity, it is unclear whether Rap1/Rif complexes can also mark a telomere for shortening by TRD. It is possible that TRD is regulated by a Rap1/Rif-independent mechanism.

specifically targeted for shortening by a mechanism called ‘telomere rapid deletion’ (TRD; Figure 1D) (57), which has also been referred to as ‘telomere trimming’ to avoid implying that the telomeres are completely deleted (58). TRD was first identified in *S. cerevisiae*, where it was shown that over-elongated telomeres could be shortened to approximately wild-type telomere length via a single intrachromosomal recombination event between telomere repeats (59, 60). A TRD event involves the excision of a telomere loop formed by the invasion of the telomeric 3' overhang into telomeric sequence further upstream in the telomere. TRD has also been observed in *Kluyveromyces lactis* (61), *Arabidopsis thaliana* (62), and human cells (58, 63), indicating that it is a general mechanism for rapidly shortening over-elongated telomeres.

Telomere length determination

Having discussed how cells shorten over-elongated telomeres, an important question still remains: how do cells determine if a telomere is over-elongated? Telomere length determination is best understood in *S. cerevisiae*. Rap1 binds to double-stranded telomeric repeats about once every 18 bp (64). Rap1 recruits two additional proteins, Rif1 and Rif2, which

act synergistically to negatively regulate telomerase (65, 66). Thus, the longer the telomere, the more Rap1/Rif1 and Rap1/Rif2 complexes will be at the telomere and the stronger the inhibition on telomerase (Figure 2A). Tethering Rap1, Rif1, or Rif2 to a telomere shortens the telomere in a manner that is proportional to the number of tethered molecules (38, 39). In human cells, a similar ‘protein-counting’ mechanism was also observed by targeting the telomeric proteins TRF1 and TRF2 to specific telomeres (67).

Although the precise details still need to be worked out, much is already known about the mechanisms by which short telomeres activate telomerase. Both physical and genetic evidence indicates that the Rif proteins inhibit Tel1, the yeast ortholog of human ataxia telangiectasia mutated (ATM), which is a positive regulator of telomerase (68, 69). When a telomere is short, there are fewer Rap1/Rif complexes at the telomere, allowing Tel1 to act. Indeed, recruitment of Tel1 to telomeres is about 10-fold higher at short telomeres than at wild-type length telomeres (70). Tel1 is a kinase and its kinase activity is important for its role in telomere length maintenance (71), but there is currently no consensus on what its critical telomeric phosphorylation targets are.

Considerably less is known about the targeting of over-elongated telomeres for shortening. As mentioned above, it

is known that telomerase activity is inhibited at long telomeres, but it is unclear whether they are specifically targeted for shortening by TRD, and if they are, whether a similar Rap1/Rif protein-counting mechanism is employed (Figure 2B). It is also unclear whether the frequency of TRD is directly proportional to telomere length. However, TRD is a recombination-mediated event, and recombination efficiency is directly proportional to the length of the substrate DNA in prokaryotes, yeast, and mammalian cells (72–77). Consistent with this notion, it has been observed, under certain circumstances, that long telomeres in *S. cerevisiae* preferentially undergo recombination (78). It will be interesting to determine whether TRD is subject to active regulation, and if so, how this is accomplished.

Outlook

Although the consequences of long telomeres have less obvious impact than those of short telomeres, it is clear that over-elongated telomeres must be shortened. More work is needed to characterize the cellular response to over-elongated telomeres, and to determine whether long telomeres are subject to evolutionary selection pressure. Such work is particularly important given the connections between telomere length homeostasis and human health.

Acknowledgments

I thank Brian Luke and Peter Lansdorp for critically reading this manuscript. I apologize to those scientists whose work I could not mention due to the space limitations of the short conceptual overview format.

Conflict of interest statement

The author declares that no conflict of interest exists.

References

- Kanaar R, Wyman C, Rothstein R. Quality control of DNA break metabolism: in the 'end', it's a good thing. *EMBO J* 2008; 27: 581–8.
- de Lange T. How telomeres solve the end-protection problem. *Science* 2009; 326: 948–52.
- Hug N, Lingner J. Telomere length homeostasis. *Chromosoma* 2006; 115: 413–25.
- Greider CW, Blackburn EH. Identification of a specific telomere terminal transferase activity in Tetrahymena extracts. *Cell* 1985; 43: 405–13.
- Nakamura TM, Morin GB, Chapman KB, Weinrich SL, Andrews WH, Lingner J, Harley CB, Cech TR. Telomerase catalytic subunit homologs from fission yeast and human. *Science* 1997; 277: 955–9.
- Feng J, Funk WD, Wang SS, Weinrich SL, Avilion AA, Chiu CP, Adams RR, Chang E, Allsopp RC, Yu J, Le S, West MD, Harley CB, Andrews WH, Greider CW, Villeponteau B. The RNA component of human telomerase. *Science* 1995; 269: 1236–41.
- Greider CW, Blackburn EH. A telomeric sequence in the RNA of Tetrahymena telomerase required for telomere repeat synthesis. *Nature* 1989; 337: 331–7.
- Singer MS, Gottschling DE. TLC1: template RNA component of *Saccharomyces cerevisiae* telomerase. *Science* 1994; 266: 404–9.
- Yu GL, Bradley JD, Attardi LD, Blackburn EH. In vivo alteration of telomere sequences and senescence caused by mutated Tetrahymena telomerase RNAs. *Nature* 1990; 344: 126–32.
- Teixeira MT, Americ M, Sperisen P, Lingner J. Telomere length homeostasis is achieved via a switch between telomerase-extendible and -nonextendible states. *Cell* 2004; 117: 323–35.
- Hemann MT, Strong MA, Hao LY, Greider CW. The shortest telomere, not average telomere length, is critical for cell viability and chromosome stability. *Cell* 2001; 107: 67–77.
- Britt-Compton B, Capper R, Rowson J, Baird DM. Short telomeres are preferentially elongated by telomerase in human cells. *FEBS Lett* 2009; 583: 3076–80.
- Armanios M. Syndromes of telomere shortening. *Annu Rev Genomics Hum Genet* 2009; 10: 45–61.
- Serrano M, Blasco MA. Cancer and ageing: convergent and divergent mechanisms. *Nat Rev Mol Cell Biol* 2007; 8: 715–22.
- Artandi SE, DePinho RA. Telomeres and telomerase in cancer. *Carcinogenesis* 2010; 31: 9–18.
- Verdun RE, Karlseder J. Replication and protection of telomeres. *Nature* 2007; 447: 924–31.
- Lundblad V, Szostak JW. A mutant with a defect in telomere elongation leads to senescence in yeast. *Cell* 1989; 57: 633–43.
- Harley CB, Futcher AB, Greider CW. Telomeres shorten during ageing of human fibroblasts. *Nature* 1990; 345: 458–60.
- Kaul Z, Cesare AJ, Huschtscha LI, Neumann AA, Reddel RR. Five dysfunctional telomeres predict onset of senescence in human cells. *EMBO Rep* 2011; 13: 52–9.
- Weinstein BS, Ciszek D. The reserve-capacity hypothesis: evolutionary origins and modern implications of the trade-off between tumor-suppression and tissue-repair. *Exp Gerontol* 2002; 37: 615–27.
- Eisenberg DT. An evolutionary review of human telomere biology: the thrifty telomere hypothesis and notes on potential adaptive paternal effects. *Am J Hum Biol* 2011; 23: 149–67.
- Rochette PJ, Brash DE. Human telomeres are hypersensitive to UV-induced DNA Damage and refractory to repair. *PLoS Genet* 2010; 6: e1000926.
- Ivessa AS, Zhou JQ, Schulz VP, Monson EK, Zakian VA. *Saccharomyces Rrm3p*, a 5' to 3' DNA helicase that promotes replication fork progression through telomeric and subtelomeric DNA. *Genes Dev* 2002; 16: 1383–96.
- Makovets S, Herskowitz I, Blackburn EH. Anatomy and dynamics of DNA replication fork movement in yeast telomeric regions. *Mol Cell Biol* 2004; 24: 4019–31.
- Ivessa AS, Lenzmeier BA, Bessler JB, Goudsouzian LK, Schnakenberg SL, Zakian VA. The *Saccharomyces cerevisiae* helicase Rrm3p facilitates replication past nonhistone protein-DNA complexes. *Mol Cell* 2003; 12: 1525–36.
- Miller KM, Rog O, Cooper JP. Semi-conservative DNA replication through telomeres requires Taz1. *Nature* 2006; 440: 824–8.
- Sfeir A, Kosiyatrakul ST, Hockemeyer D, MacRae SL, Karlseder J, Schildkraut CL, de Lange T. Mammalian telomeres resemble fragile sites and require TRF1 for efficient replication. *Cell* 2009; 138: 90–103.
- Sen D, Gilbert W. Formation of parallel four-stranded complexes by guanine-rich motifs in DNA and its implications for meiosis. *Nature* 1988; 334: 364–6.

29. Sundquist WI, Klug A. Telomeric DNA dimerizes by formation of guanine tetrads between hairpin loops. *Nature* 1989; 342: 825–9.
30. Paeschke K, Simonsson T, Postberg J, Rhodes D, Lipps HJ. Telomere end-binding proteins control the formation of G-quadruplex DNA structures in vivo. *Nat Struct Mol Biol* 2005; 12: 847–54.
31. Paeschke K, Juranek S, Simonsson T, Hempel A, Rhodes D, Lipps HJ. Telomerase recruitment by the telomere end binding protein- facilitates G-quadruplex DNA unfolding in ciliates. *Nat Struct Mol Biol* 2008; 15: 598–604.
32. Paeschke K, Capra JA, Zakian VA. DNA replication through G-quadruplex motifs is promoted by the *Saccharomyces cerevisiae* Pif1 DNA helicase. *Cell* 2011; 145: 678–91.
33. Crabbe L, Verdun RE, Haggblom CI, Karlseder J. Defective telomere lagging strand synthesis in cells lacking WRN helicase activity. *Science* 2004; 306: 1951–3.
34. Ding H, Schertzer M, Wu X, Gertsenstein M, Selig S, Kammori M, Pourvali R, Poon S, Vulto I, Chavez E, Tam PP, Nagy A, Lansdorff PM. Regulation of murine telomere length by Rtel: an essential gene encoding a helicase-like protein. *Cell* 2004; 117: 873–86.
35. Kurth I, Gautier J. Origin-dependent initiation of DNA replication within telomeric sequences. *Nucleic Acids Res* 2010; 38: 467–76.
36. Puglisi A, Bianchi A, Lemmens L, Damay P, Shore D. Distinct roles for yeast Stn1 in telomere capping and telomerase inhibition. *EMBO J* 2008; 27: 2328–39.
37. Cristofari G, Lingner J. Telomere length homeostasis requires that telomerase levels are limiting. *EMBO J* 2006; 25: 565–74.
38. Marcand S, Gilson E, Shore D. A protein-counting mechanism for telomere length regulation in yeast. *Science* 1997; 275: 986–90.
39. Levy DL, Blackburn EH. Counting of Rif1p and Rif2p on *Saccharomyces cerevisiae* telomeres regulates telomere length. *Mol Cell Biol* 2004; 24: 10857–67.
40. Shore D, Nasmyth K. Purification and cloning of a DNA binding protein from yeast that binds to both silencer and activator elements. *Cell* 1987; 51: 721–32.
41. Aparicio OM, Billington BL, Gottschling DE. Modifiers of position effect are shared between telomeric and silent mating-type loci in *S. cerevisiae*. *Cell* 1991; 66: 1279–87.
42. Sinclair DA, Guarente L. Extrachromosomal rDNA circles – a cause of aging in yeast. *Cell* 1997; 91: 1033–42.
43. Guarente L. Sir2 links chromatin silencing, metabolism, and aging. *Genes Dev* 2000; 14: 1021–6.
44. Austriaco NR Jr, Guarente LP. Changes of telomere length cause reciprocal changes in the lifespan of mother cells in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci USA* 1997; 94: 9768–72.
45. Kyrion G, Liu K, Liu C, Lustig AJ. RAP1 and telomere structure regulate telomere position effects in *Saccharomyces cerevisiae*. *Genes Dev* 1993; 7: 1146–59.
46. Buck SW, Shore D. Action of a RAP1 carboxy-terminal silencing domain reveals an underlying competition between HMR and telomeres in yeast. *Genes Dev* 1995; 9: 370–84.
47. Marcand S, Brevet V, Gilson E. Progressive cis-inhibition of telomerase upon telomere elongation. *EMBO J* 1999; 18: 3509–19.
48. Niida H, Matsumoto T, Satoh H, Shiwa M, Tokutake Y, Furuichi Y, Shinkai Y. Severe growth defect in mouse cells lacking the telomerase RNA component. *Nat Genet* 1998; 19: 203–6.
49. Counter CM, Avilion AA, LeFeuvre CE, Stewart NG, Greider CW, Harley CB, Bacchetti S. Telomere shortening associated with chromosome instability is arrested in immortal cells which express telomerase activity. *EMBO J* 1992; 11: 1921–9.
50. Bodnar AG, Ouellette M, Frolkis M, Holt SE, Chiu CP, Morin GB, Harley CB, Shay JW, Lichtsteiner S, Wright WE. Extension of life-span by introduction of telomerase into normal human cells. *Science* 1998; 279: 349–52.
51. Yang J, Chang E, Cherry AM, Bangs CD, Oei Y, Bodnar A, Bronstein A, Chiu CP, Herron GS. Human endothelial cell life extension by telomerase expression. *J Biol Chem* 1999; 274: 26141–8.
52. Olovnikov AM. [Principle of marginotomy in template synthesis of polynucleotides]. *Dokl Akad Nauk SSSR* 1971; 201: 1496–9.
53. Watson JD. Origin of concatemeric T7 DNA. *Nat New Biol* 1972; 239: 197–201.
54. von Zglinicki T, Saretzki G, Döcke W, Lotze C. Mild hyperoxia shortens telomeres and inhibits proliferation of fibroblasts: a model for senescence? *Exp Cell Res* 1995; 220: 186–93.
55. von Zglinicki T, Pilger R, Sittte N. Accumulation of single-strand breaks is the major cause of telomere shortening in human fibroblasts. *Free Radic Biol Med* 2000; 28: 64–74.
56. Chang M, Arneric M, Lingner J. Telomerase repeat addition processivity is increased at critically short telomeres in a Tel1-dependent manner in *Saccharomyces cerevisiae*. *Genes Dev* 2007; 21: 2485–94.
57. Lustig AJ. Clues to catastrophic telomere loss in mammals from yeast telomere rapid deletion. *Nat Rev Genet* 2003; 4: 916–23.
58. Pickett HA, Cesare AJ, Johnston RL, Neumann AA, Reddel RR. Control of telomere length by a trimming mechanism that involves generation of t-circles. *EMBO J* 2009; 28: 799–809.
59. Li B, Lustig AJ. A novel mechanism for telomere size control in *Saccharomyces cerevisiae*. *Genes Dev* 1996; 10: 1310–26.
60. Bucholc M, Park Y, Lustig AJ. Intrachromatid excision of telomeric DNA as a mechanism for telomere size control in *Saccharomyces cerevisiae*. *Mol Cell Biol* 2001; 21: 6559–73.
61. Bechard LH, Jamieson N, McEachern MJ. Recombination can cause telomere elongations as well as truncations deep within telomeres in wild-type *Kluyveromyces lactis* cells. *Eukaryot Cell* 2011; 10: 226–36.
62. Watson JM, Shippen DE. Telomere rapid deletion regulates telomere length in *Arabidopsis thaliana*. *Mol Cell Biol* 2007; 27: 1706–15.
63. Pickett HA, Henson JD, Au AY, Neumann AA, Reddel RR. Normal mammalian cells negatively regulate telomere length by telomere trimming. *Hum Mol Genet* 2011; 20: 4684–92.
64. Gilson E, Roberge M, Giraldo R, Rhodes D, Gasser SM. Distortion of the DNA double helix by RAP1 at silencers and multiple telomeric binding sites. *J Mol Biol* 1993; 231: 293–310.
65. Hardy CF, Sussel L, Shore D. A RAP1-interacting protein involved in transcriptional silencing and telomere length regulation. *Genes Dev* 1992; 6: 801–14.
66. Wotton D, Shore D. A novel Rap1p-interacting factor, Rif2p, cooperates with Rif1p to regulate telomere length in *Saccharomyces cerevisiae*. *Genes Dev* 1997; 11: 748–60.
67. Ancelin K, Brunori M, Bauwens S, Koering CE, Brun C, Ricoul M, Pommier JP, Sabatier L, Gilson E. Targeting assay to study the cis functions of human telomeric proteins: evidence for inhibition of telomerase by TRF1 and for activation of telomere degradation by TRF2. *Mol Cell Biol* 2002; 22: 3474–87.
68. Hirano Y, Fukunaga K, Sugimoto K. Rif1 and rif2 inhibit localization of tel1 to DNA ends. *Mol Cell* 2009; 33: 312–22.
69. Craven RJ, Petes TD. Dependence of the regulation of telomere length on the type of subtelomeric repeat in the yeast *Saccharomyces cerevisiae*. *Genetics* 1999; 152: 1531–41.

70. Sabourin M, Tuzon CT, Zakian VA. Telomerase and Tel1p preferentially associate with short telomeres in *S. cerevisiae*. *Mol Cell* 2007; 27: 550–61.
71. Mallory JC, Petes TD. Protein kinase activity of Tel1p and Mec1p, two *Saccharomyces cerevisiae* proteins related to the human ATM protein kinase. *Proc Natl Acad Sci USA* 2000; 97: 13749–54.
72. Singer BS, Gold L, Gauss P, Doherty DH. Determination of the amount of homology required for recombination in bacteriophage T4. *Cell* 1982; 31: 25–33.
73. Watt VM, Ingles CJ, Urdea MS, Rutter WJ. Homology requirements for recombination in *Escherichia coli*. *Proc Natl Acad Sci USA* 1985; 82: 4768–72.
74. Shen P, Huang HV. Homologous recombination in *Escherichia coli*: dependence on substrate length and homology. *Genetics* 1986; 112: 441–57.
75. Jinks-Robertson S, Michelitch M, Ramcharan S. Substrate length requirements for efficient mitotic recombination in *Saccharomyces cerevisiae*. *Mol Cell Biol* 1993; 13: 3937–50.
76. Rubnitz J, Subramani S. The minimum amount of homology required for homologous recombination in mammalian cells. *Mol Cell Biol* 1984; 4: 2253–8.
77. Liskay RM, Letsou A, Stachelek JL. Homology requirement for efficient gene conversion between duplicated chromosomal sequences in mammalian cells. *Genetics* 1987; 115: 161–7.
78. Chang M, Dittmar JC, Rothstein R. Long telomeres are preferentially extended during recombination-mediated telomere maintenance. *Nat Struct Mol Biol* 2011; 18: 451–6.

Received March 20, 2012; accepted April 10, 2012



Michael Chang received his PhD degree at the University of Toronto in 2005 under the supervision of Dr. Grant W. Brown, studying DNA damage response pathways using high-throughput functional genomics. After his PhD, he took a position at the Swiss Institute for Experimental Cancer Research in Lausanne as a postdoctoral fellow in the lab of Dr. Joachim

Lingner, whose research is focused on telomerase and chromosome end replication. In 2008, Michael moved to the lab of Dr. Rodney Rothstein at the Columbia University Medical Center in New York, where he continued to study factors that regulate telomerase as well as telomerase-independent mechanisms of telomere maintenance. In 2011, he joined the European Research Institute for the Biology of Ageing in the Netherlands as an Assistant Professor. Work in his lab focuses on telomere maintenance and genome integrity as it relates to cancer and aging.