

# An approximation theorem for solutions of degenerate semilinear elliptic equations

*Albo Carlos Cavalheiro*

**Abstract.** The main result establishes that a weak solution of degenerate semilinear elliptic equations can be approximated by a sequence of solutions for non-degenerate semilinear elliptic equations.

## 1 Introduction

Let  $L$  be a degenerate elliptic operator in divergence form

$$Lu = - \sum_{i,j=1}^n D_j(a_{ij}(x)D_i u(x)), \quad D_j = \frac{\partial}{\partial x_j}, \quad (1)$$

where the coefficients  $a_{ij}$  are measurable, real-valued functions whose coefficient matrix  $\mathcal{A} = (a_{ij})$  is symmetric and satisfies the degenerate ellipticity condition

$$\lambda|\xi|^2\omega(x) \leq \sum_{i,j=1}^n a_{ij}(x)\xi_i\xi_j \leq \Lambda|\xi|^2\omega(x), \quad (2)$$

for all  $\xi \in \mathbb{R}^n$  and almost everywhere  $x \in \Omega \subset \mathbb{R}^n$  a bounded open set,  $\omega$  is a weight function,  $\lambda$  and  $\Lambda$  are positive constants.

The main purpose of this paper (see Theorem 1) is to establish that a weak solution  $u \in W_0^{1,2}(\Omega, \omega)$  for the semilinear Dirichlet problem

$$(P) \begin{cases} Lu(x) - \gamma u(x)g_1(x) + h(u(x))g_2(x) = f(x) & \text{in } \Omega, \\ u(x) = 0 & \text{on } \partial\Omega, \end{cases}$$

where  $\gamma \in \mathbb{R}$ , can be approximated by a sequence of solutions of non-degenerate semilinear elliptic equations.

---

2010 MSC: 35J61, 35J70, 35D30.

Key words: Degenerate semilinear elliptic equations, weighted Sobolev Spaces.

DOI: 10.1515/cm-2017-0004

By a *weight*, we shall mean a locally integrable function  $\omega$  on  $\mathbb{R}^n$  such that  $\omega(x) > 0$  for a.e.  $x \in \mathbb{R}^n$ . Every weight  $\omega$  gives rise to a measure on the measurable subsets on  $\mathbb{R}^n$  through integration. This measure will be denoted by  $\mu$ . Thus,  $\mu(E) = \int_E \omega(x) dx$  for measurable sets  $E \subset \mathbb{R}^n$ .

In general, the Sobolev spaces  $W^{k,p}(\Omega)$  without weights occur as spaces of solutions for elliptic and parabolic partial differential equations. For degenerate partial differential equations, i.e., equations with various types of singularities in the coefficients, it is natural to look for solutions in weighted Sobolev spaces (see [1], [2], [4], [3], [5], [6], [7], [9] and [13]).

A class of weights, which is particularly well understood, is the class of  $A_p$ -weights (or Muckenhoupt class) that was introduced by B. Muckenhoupt (see [11]). These classes have found many useful applications in harmonic analysis (see [12]). Another reason for studying  $A_p$ -weights is the fact that powers of the distance to submanifolds of  $\mathbb{R}^n$  often belong to  $A_p$  (see [10]). There are, in fact, many interesting examples of weights (see [9] for  $p$ -admissible weights).

The following lemma can be proved in exactly the same way as Lemma 2.1 in [7] (see also, Lemma 3.1 and Lemma 4.13 in [1]). Our lemma provides a general approximation theorem for  $A_p$ -weights ( $1 \leq p < \infty$ ) by means of weights which are bounded away from 0 and infinity and whose  $A_p$ -constants depend only on the  $A_p$ -constant of  $\omega$ . Lemma 1 is the key point for Theorem 1, and the crucial point consists of showing that a weak limit of a sequence of solutions of approximate problems is in fact a solution of the original problem.

**Lemma 1.** *Let  $\alpha, \beta > 1$  be given and let  $\omega \in A_p$  ( $1 \leq p < \infty$ ), with  $A_p$ -constant  $C(\omega, p)$  and let  $a_{ij} = a_{ji}$  be measurable, real-valued functions satisfying*

$$\lambda\omega(x)|\xi|^2 \leq \sum_{i,j=1}^n a_{ij}(x)\xi_i\xi_j \leq \Lambda\omega(x)|\xi|^2, \quad (3)$$

for all  $\xi \in \mathbb{R}^n$  and a.e.  $x \in \Omega$ . Then there exist weights  $\omega_{\alpha\beta} \geq 0$  a.e. and measurable real-valued functions  $a_{ij}^{\alpha\beta}$  such that the following conditions are met.

- (i)  $c_1(1/\beta) \leq \omega_{\alpha\beta} \leq c_2\alpha$  in  $\Omega$ , where  $c_1$  and  $c_2$  depend only on  $\omega$  and  $\Omega$ .
- (ii) There exist weights  $\tilde{\omega}_1$  and  $\tilde{\omega}_2$  such that  $\tilde{\omega}_1 \leq \omega_{\alpha\beta} \leq \tilde{\omega}_2$ , where  $\tilde{\omega}_i \in A_p$  and  $C(\tilde{\omega}_i, p)$  depends only on  $C(\omega, p)$  ( $i = 1, 2$ ).
- (iii)  $\omega_{\alpha\beta} \in A_p$ , with constant  $C(\omega_{\alpha\beta}, p)$  depending only on  $C(\omega, p)$  uniformly on  $\alpha$  and  $\beta$ .
- (iv) There exists a closed set  $F_{\alpha\beta}$  such that  $\omega_{\alpha\beta} \equiv \omega$  in  $F_{\alpha\beta}$  and  $\omega_{\alpha\beta} \sim \tilde{\omega}_1 \sim \tilde{\omega}_2$  in  $F_{\alpha\beta}$  with equivalence constants depending on  $\alpha$  and  $\beta$  (i.e., there are positive constants  $c_{\alpha\beta}$  and  $C_{\alpha\beta}$  such that  $c_{\alpha\beta}\tilde{\omega}_i \leq \omega_{\alpha\beta} \leq C_{\alpha\beta}\tilde{\omega}_i$ ,  $i = 1, 2$ ). Moreover,  $F_{\alpha\beta} \subset F_{\alpha'\beta'}$  if  $\alpha \leq \alpha'$ ,  $\beta \leq \beta'$ , and the complement of  $\bigcup_{\alpha, \beta \geq 1} F_{\alpha\beta}$  has zero measure.
- (v)  $\omega_{\alpha\beta} \rightarrow \omega$  a.e. in  $\mathbb{R}^n$  as  $\alpha, \beta \rightarrow \infty$ .

$$(vi) \quad \lambda \omega_{\alpha\beta}(x)|\xi|^2 \leq \sum_{i,j=1}^n a_{ij}^{\alpha\beta}(x)\xi_i\xi_j \leq \Lambda \omega_{\alpha\beta}(x)|\xi|^2, \text{ for every } \xi \in \mathbb{R} \text{ and a.e. } x \in \Omega.$$

*Proof.* See [1], Lemma 3.1 or Lemma 4.13. □

The following theorem will be proved in Section 3.

**Theorem 1.** *Suppose that*

(H1) *The function  $h: \mathbb{R} \rightarrow \mathbb{R}$  is Lipschitz continuous (i.e., there exists a constant  $C_h > 0$  such that  $|h(t_1) - h(t_2)| \leq C_h|t_1 - t_2|$  for all  $t_1, t_2 \in \mathbb{R}$ ) and  $h(0) = 0$ ;*

(H2)  $\omega \in A_2$ ;

(H3)  $g_1/\omega \in L^\infty(\Omega)$ ,  $g_2/\omega \in L^\infty(\Omega)$  and  $f/\omega \in L^2(\Omega, \omega)$ ;

(H4)  $\gamma > 0$  is not an eigenvalue of the linearized problem

$$(LP) \quad \begin{cases} Lu(x) - \gamma u(x)\omega(x) = 0 & \text{in } \Omega, \\ u(x) = 0 & \text{on } \partial\Omega; \end{cases}$$

(H5) *The constant  $M = \lambda - \gamma C_\Omega^2 \|g_1/\omega\|_{L^\infty(\Omega)} - C_h C_\Omega^2 \|g_2/\omega\|_{L^\infty(\Omega)} > 0$  (with  $C_\Omega$  as in Theorem 2).*

*Then the problem (P) has a unique solution  $u \in W_0^{1,2}(\Omega, \omega)$  and there exists a constant  $C > 0$  such that*

$$\|u\|_{W_0^{1,2}(\Omega, \omega)} \leq C \left\| \frac{f}{\omega} \right\|_{L^2(\Omega, \omega)}. \tag{4}$$

*Moreover,  $u$  is the weak limit in  $W_0^{1,2}(\Omega, \tilde{\omega}_1)$  of a sequence of solutions  $u_m \in W_0^{1,2}(\Omega, \omega_m)$  of the problems*

$$(P_m) \quad \begin{cases} L_m u_m(x) - \gamma u_m(x)g_{1m}(x) + h(u_m(x))g_{2m}(x) = f_m(x) & \text{in } \Omega, \\ u_m(x) = 0 & \text{on } \partial\Omega, \end{cases}$$

*with  $L_m u_m = - \sum_{i,j=1}^n D_j(a_{ij}^{mm}(x)D_i u_m(x))$ ,  $g_{1m} = g_1\omega_m/\omega$ ,  $g_{2m} = g_2\omega_m/\omega$ ,*

*$f_m = f(\omega/\omega_m)^{-1/2}$  and  $\omega_m = \omega_{mm}$  (where  $\omega_{mm}$ ,  $a_{ij}^{mm}$  and  $\tilde{\omega}_1$  are as Lemma 1).*

## 2 Definitions and basic results

Let  $\omega$  be a locally integrable nonnegative function in  $\mathbb{R}^n$  and assume that  $0 < \omega(x) < \infty$  almost everywhere. We say that  $\omega$  belongs to the Muckenhoupt class  $A_p$ ,  $1 < p < \infty$ , or that  $\omega$  is an  $A_p$ -weight, if there is a constant  $C = C(p, \omega)$  such that

$$\left( \frac{1}{|B|} \int_B \omega(x) dx \right) \left( \frac{1}{|B|} \int_B \omega^{1/(1-p)}(x) dx \right)^{p-1} \leq C,$$

for all balls  $B \subset \mathbb{R}^n$ , where  $|\cdot|$  denotes the  $n$ -dimensional Lebesgue measure in  $\mathbb{R}^n$ . If  $1 < q \leq p$ , then  $A_q \subset A_p$  (see [8], [9] or [13] for more information about  $A_p$ -weights). The weight  $\omega$  satisfies the doubling condition if there exists a positive constant  $C$  such that  $\mu(B(x; 2r)) \leq C\mu(B(x; r))$  for every ball  $B = B(x; r) \subset \mathbb{R}^n$ , where  $\mu(B) = \int_B \omega(x) dx$ . If  $\omega \in A_p$ , then  $\mu$  is doubling (see Corollary 15.7 in [9]).

As an example of  $A_p$ -weight, the function  $\omega(x) = |x|^\alpha$ ,  $x \in \mathbb{R}^n$ , is in  $A_p$  if and only if  $-n < \alpha < n(p-1)$  (see Corollary 4.4, Chapter IX in [12]).

If  $\omega \in A_p$ , then  $\left(\frac{|E|}{|B|}\right)^p \leq C \frac{\mu(E)}{\mu(B)}$  whenever  $B$  is a ball in  $\mathbb{R}^n$  and  $E$  is a measurable subset of  $B$  (see 15.5 *strong doubling property* in [9]). Therefore,  $\mu(E) = 0$  if and only if  $|E| = 0$ ; so there is no need to specify the measure when using the ubiquitous expression almost everywhere and almost every, both abbreviated a.e.

**Definition 1.** Let  $\omega$  be a weight, and let  $\Omega \subset \mathbb{R}^n$  be open. For  $0 < p < \infty$  we define  $L^p(\Omega, \omega)$  as the set of measurable functions  $f$  on  $\Omega$  such that

$$\|f\|_{L^p(\Omega, \omega)} = \left( \int_{\Omega} |f(x)|^p \omega(x) dx \right)^{1/p} < \infty.$$

If  $\omega \in A_p$ ,  $1 < p < \infty$ , then  $\omega^{-1/(p-1)}$  is locally integrable and we have  $L^p(\Omega, \omega) \subset L^1_{\text{loc}}(\Omega)$  for every open set  $\Omega$  (see Remark 1.2.4 in [13]). It thus makes sense to talk about weak derivatives of functions in  $L^p(\Omega, \omega)$ .

**Definition 2.** Let  $\Omega \subset \mathbb{R}^n$  be open,  $k$  be a nonnegative integer and  $\omega \in A_p$  ( $1 < p < \infty$ ). We define the weighted Sobolev space  $W^{k,p}(\Omega, \omega)$  as the set of functions  $u \in L^p(\Omega, \omega)$  with weak derivatives  $D^\alpha u \in L^p(\Omega, \omega)$  for  $1 \leq |\alpha| \leq k$ . The norm of  $u$  in  $W^{k,p}(\Omega, \omega)$  is defined by

$$\|u\|_{W^{k,p}(\Omega, \omega)} = \left( \int_{\Omega} |u(x)|^p \omega(x) dx + \sum_{1 \leq |\alpha| \leq k} \int_{\Omega} |D^\alpha u(x)|^p \omega(x) dx \right)^{1/p}. \quad (5)$$

We also define  $W_0^{k,p}(\Omega, \omega)$  as the closure of  $C_0^\infty(\Omega)$  with respect to the norm (5).

If  $\omega \in A_p$ , then  $W^{k,p}(\Omega, \omega)$  is the closure of  $C^\infty(\Omega)$  with respect to the norm (5) (see Theorem 2.1.4 in [13]). The spaces  $W^{k,p}(\Omega, \omega)$  and  $W_0^{k,p}(\Omega, \omega)$  are Banach spaces.

It is evident that the weight function  $\omega$  which satisfies  $0 < c_1 \leq \omega(x) \leq c_2$  for  $x \in \Omega$  ( $c_1$  and  $c_2$  positive constants), gives nothing new (the space  $W_0^{k,p}(\Omega, \omega)$  is then identical with the classical Sobolev space  $W_0^{k,p}(\Omega)$ ). Consequently, we shall be interested above in all such weight functions  $\omega$  which either vanish in somewhere  $\Omega \cup \partial\Omega$  or increase to infinity (or both).

The dual space of  $W_0^{1,p}(\Omega, \omega)$  is the space

$$\begin{aligned} [W_0^{1,p}(\Omega, \omega)]^* &= W^{-1,p'}(\Omega, \omega) \\ &= \left\{ T = f_0 - \operatorname{div} F : F = (f_1, \dots, f_n), \frac{f_j}{\omega} \in L^{p'}(\Omega, \omega) \right\}. \end{aligned}$$

**Definition 3.** We say that an element  $u \in W_0^{1,2}(\Omega, \omega)$  is weak solution of problem (P) if

$$\int_{\Omega} a_{ij}(x)D_i u(x)D_j \varphi(x) \, dx - \int_{\Omega} \gamma u(x)g_1(x)\varphi(x) \, dx + \int_{\Omega} h(u(x))g_2(x)\varphi(x) \, dx = \int_{\Omega} f(x)\varphi(x) \, dx,$$

for every  $\varphi \in W_0^{1,2}(\Omega, \omega)$ .

**Theorem 2.** (The weighted Sobolev inequality) Let  $\Omega$  be an open bounded set in  $\mathbb{R}^n$  and  $\omega \in A_p$  ( $1 < p < \infty$ ). There exist positive constants  $C_{\Omega}$  and  $\delta$  such that for all  $u \in W_0^{1,p}(\Omega, \omega)$  and all  $\theta$  satisfying  $1 \leq \theta \leq n/(n-1) + \delta$ ,

$$\|u\|_{L^{\theta p}(\Omega, \omega)} \leq C_{\Omega} \|\nabla u\|_{L^p(\Omega, \omega)}. \tag{6}$$

*Proof.* Its suffices to prove the inequality for functions  $u \in C_0^{\infty}(\Omega)$  (see Theorem 1.3 in [6]). To extend the estimates (6) to arbitrary  $u \in W_0^{1,p}(\Omega, \omega)$ , we let  $\{u_m\}$  be a sequence of  $C_0^{\infty}(\Omega)$  functions tending to  $u$  in  $W_0^{1,p}(\Omega, \omega)$ . Applying the estimates (6) to differences  $u_{m_1} - u_{m_2}$ , we see that  $\{u_m\}$  will be a Cauchy sequence in  $L^{kp}(\Omega, \omega)$ . Consequently the limit function  $u$  will lie in the desired spaces and satisfy (6).  $\square$

### 3 Proof of Theorem 1

**Step 1.** The existence of solution  $u \in W_0^{1,2}(\Omega, \omega)$  for the problem (P) has been demonstrated in [2], Theorem 1. In particular, for  $\varphi = u$  in Definition 3, we have

$$\int_{\Omega} a_{ij}(x)D_i u(x)D_j u(x) \, dx - \int_{\Omega} \gamma u^2(x)g_1(x) \, dx + \int_{\Omega} h(u(x))g_2(x)u(x) \, dx = \int_{\Omega} f(x)u(x) \, dx. \tag{7}$$

(i) By (2) we have

$$\int_{\Omega} a_{ij}(x)D_i u(x)D_j u(x) \, dx \geq \lambda \int_{\Omega} |\nabla u|^2 \omega \, dx.$$

(ii) By (H3) and Theorem 2 (with  $p = 2$  and  $\theta = 1$ ) we obtain

$$\begin{aligned} \left| \int_{\Omega} \gamma u^2 g_1 \, dx \right| &\leq \gamma \int_{\Omega} u^2 \frac{|g_1|}{\omega} \omega \, dx \\ &\leq \gamma \|g_1/\omega\|_{L^{\infty}(\Omega)} \int_{\Omega} u^2 \omega \, dx \\ &\leq \gamma C_{\Omega}^2 \|g_1/\omega\|_{L^{\infty}(\Omega)} \int_{\Omega} |\nabla u|^2 \omega \, dx, \end{aligned}$$

and

$$\begin{aligned} \left| \int_{\Omega} f u \, dx \right| &\leq \int_{\Omega} \frac{|f|}{\omega} |u| \omega \, dx \\ &\leq \|f/\omega\|_{L^2(\Omega, \omega)} \|u\|_{L^2(\Omega, \omega)} \\ &\leq C_{\Omega} \|f/\omega\|_{L^2(\Omega, \omega)} \|\nabla u\|_{L^2(\Omega, \omega)}. \end{aligned}$$

(iii) By (H1), since  $h(0) = 0$ , then  $|h(t)| \leq C_h |t|$  for all  $t \in \mathbb{R}$ . By (H3) and Theorem 2, we obtain

$$\begin{aligned} \left| \int_{\Omega} h(u) u g_2 \, dx \right| &\leq \int_{\Omega} |h(u)| |u| \frac{|g_2|}{\omega} \omega \, dx \\ &\leq C_h \|g_2/\omega\|_{L^{\infty}(\Omega)} \int_{\Omega} |u|^2 \omega \, dx \\ &\leq C_h C_{\Omega}^2 \|g_2/\omega\|_{L^{\infty}(\Omega)} \int_{\Omega} |\nabla u|^2 \omega \, dx. \end{aligned}$$

Hence, in (7), we obtain

$$\begin{aligned} \lambda \int_{\Omega} |\nabla u|^2 \omega \, dx - \gamma C_{\Omega}^2 \|g_1/\omega\|_{L^{\infty}(\Omega)} \int_{\Omega} |\nabla u|^2 \omega \, dx \\ - C_h C_{\Omega}^2 \|g_2/\omega\|_{L^{\infty}(\Omega)} \int_{\Omega} |\nabla u|^2 \omega \, dx \leq C_{\Omega} \|f/\omega\| \|\nabla u\|_{L^2(\Omega, \omega)}. \end{aligned}$$

Therefore,

$$\|\nabla u\|_{L^2(\Omega, \omega)} \leq \frac{C_{\Omega}}{M} \|f/\omega\|_{L^2(\Omega, \omega)}$$

where

$$M = \lambda - \gamma C_{\Omega}^2 \|g_1/\omega\|_{L^{\infty}(\Omega)} - C_h C_{\Omega}^2 \|g_2/\omega\|_{L^{\infty}(\Omega)} > 0.$$

Consequently, we obtain

$$\begin{aligned} \|u\|_{W_0^{1,2}(\Omega, \omega)}^2 &= \int_{\Omega} u^2 \omega \, dx + \int_{\Omega} |\nabla u|^2 \omega \, dx \\ &\leq (C_{\Omega}^2 + 1) \int_{\Omega} |\nabla u|^2 \omega \, dx \\ &\leq (C_{\Omega}^2 + 1) \frac{C_{\Omega}^2}{M^2} \|f/\omega\|_{L^2(\Omega, \omega)}^2. \end{aligned}$$

Therefore,

$$\begin{aligned} \|u\|_{W_0^{1,2}(\Omega, \omega)} &\leq (C_{\Omega}^2 + 1)^{1/2} \frac{C_{\Omega}}{M} \|f/\omega\|_{L^2(\Omega, \omega)} \\ &= C \|f/\omega\|_{L^2(\Omega, \omega)}. \end{aligned} \tag{8}$$

**Step 2.** Uniqueness.

If  $u_1, u_2 \in W_0^{1,2}(\Omega, \omega)$  are solutions of the problem (P), then

$$\int_{\Omega} a_{ij} D_i u_k D_j \varphi \, dx - \gamma \int_{\Omega} u_k g_1 \varphi \, dx + \int_{\Omega} h(u_k) g_2 \varphi \, dx = \int_{\Omega} f \varphi \, dx, \quad (k = 1, 2).$$

Hence,

$$\int_{\Omega} a_{ij} (D_i u_1 - D_i u_2) D_j \varphi \, dx - \gamma \int_{\Omega} (u_1 - u_2) g_1 \varphi \, dx + \int_{\Omega} (h(u_1) - h(u_2)) g_2 \varphi \, dx = 0, \tag{9}$$

for all  $\varphi \in W_0^{1,2}(\Omega, \omega)$ . In particular, for  $\varphi = u_1 - u_2$  in (9) we obtain:

(i) By (2),

$$\int_{\Omega} a_{ij} D_i (u_1 - u_2) D_j (u_1 - u_2) \, dx \geq \lambda \int_{\Omega} |\nabla(u_1 - u_2)|^2 \omega \, dx.$$

(ii) By (H3) and Theorem 2 (with  $p = 2$  and  $\theta = 1$ ),

$$\begin{aligned} \left| \int_{\Omega} (u_1 - u_2)^2 g_1 \, dx \right| &\leq \int_{\Omega} |u_1 - u_2|^2 \frac{|g_1|}{\omega} \omega \, dx \\ &\leq \|g_1/\omega\|_{L^\infty(\Omega)} \int_{\Omega} |u_1 - u_2|^2 \omega \, dx \\ &\leq C_\Omega^2 \|g_1/\omega\|_{L^\infty(\Omega)} \int_{\Omega} |\nabla(u_1 - u_2)|^2 \omega \, dx. \end{aligned}$$

(iii) By (H1), (H3) and Theorem 2,

$$\begin{aligned} \left| \int_{\Omega} (h(u_1) - h(u_2))(u_1 - u_2) g_2 \, dx \right| &\leq \int_{\Omega} |h(u_1) - h(u_2)| |u_1 - u_2| \frac{|g_2|}{\omega} \omega \, dx \\ &\leq C_h \int_{\Omega} |u_1 - u_2|^2 \frac{|g_2|}{\omega} \omega \, dx \\ &\leq C_h C_\Omega^2 \|g_2/\omega\|_{L^\infty(\Omega)} \int_{\Omega} |\nabla(u_1 - u_2)|^2 \omega \, dx. \end{aligned}$$

Hence,

$$\left( \lambda - \gamma C_\Omega^2 \|g_1/\omega\|_{L^\infty(\Omega)} - C_h C_\Omega^2 \|g_2/\omega\|_{L^\infty(\Omega)} \right) \int_{\Omega} |\nabla(u_1 - u_2)|^2 \omega \, dx \leq 0,$$

and since  $M = \lambda - \gamma C_\Omega^2 \|g_1/\omega\|_{L^\infty(\Omega)} - C_h C_\Omega^2 \|g_2/\omega\|_{L^\infty(\Omega)} > 0$ , then

$$\int_{\Omega} |\nabla(u_1 - u_2)|^2 \omega \, dx = 0,$$

and  $\int_{\Omega} |u_1 - u_2|^2 \omega \, dx = 0$ . Therefore,  $u_1 = u_2$  a.e.

**Step 3.** First, if  $f_m = f(\omega/\omega_m)^{-1/2}$ ,  $g_{1m} = g_1\omega_m/\omega$  and  $g_{2m} = g_2\omega_m/\omega$ , we note that

$$\begin{aligned} \left\| \frac{f_m}{\omega_m} \right\|_{L^2(\Omega, \omega_m)} &= \left\| \frac{f}{\omega} \right\|_{L^2(\Omega, \omega)}, & \left\| \frac{g_{1m}}{\omega_m} \right\|_{L^\infty(\Omega)} &= \left\| \frac{g_1}{\omega} \right\|_{L^\infty(\Omega)}, \\ \text{and } \left\| \frac{g_{2m}}{\omega_m} \right\|_{L^\infty(\Omega)} &= \left\| \frac{g_2}{\omega} \right\|_{L^\infty(\Omega)}. \end{aligned}$$

Then, we have

$$\begin{aligned} M_m &= \lambda - \gamma C_\Omega^2 \|g_{1m}/\omega_m\|_{L^\infty(\Omega)} - C_h C_\Omega^2 \|g_{2m}\omega_m\|_{L^\infty(\Omega)} \\ &= \lambda - \gamma C_\Omega^2 \|g_1/\omega\|_{L^\infty(\Omega)} - C_h C_\Omega^2 \|g_2/\omega\|_{L^\infty(\Omega)} = M > 0. \end{aligned}$$

If  $u_m \in W_0^{1,2}(\Omega, \omega_m)$  is a solution of problem  $(P_m)$  we have (by (8))

$$\begin{aligned} \|u_m\|_{W_0^{1,2}(\Omega, \omega_m)} &\leq \frac{(C_\Omega^2 + 1)^{1/2} C_\Omega}{M_m} \|f_m/\omega_m\|_{L^2(\Omega, \omega_m)} \\ &= \frac{(C_\Omega^2 + 1)^{1/2} C_\Omega}{M} \|f/\omega\|_{L^2(\Omega, \omega)} = C_1. \end{aligned}$$

Using Lemma 1,  $\tilde{\omega}_1 \leq \omega_m$ , we obtain

$$\|u_m\|_{W_0^{1,2}(\Omega, \tilde{\omega}_1)} \leq \|u_m\|_{W_0^{1,2}(\Omega, \omega_m)} \leq C_1. \quad (10)$$

Consequently,  $\{u_m\}$  is a bounded sequence in  $W_0^{1,2}(\Omega, \tilde{\omega}_1)$ . Therefore, there is a subsequence, again denoted by  $\{u_m\}$ , and  $\tilde{u} \in W_0^{1,2}(\Omega, \tilde{\omega}_1)$  such that

$$u_m \rightharpoonup \tilde{u} \quad \text{in } L^2(\Omega, \tilde{\omega}_1), \quad (11)$$

$$\nabla u_m \rightharpoonup \nabla \tilde{u} \quad \text{in } L^2(\Omega, \tilde{\omega}_1), \quad (12)$$

$$u_m \rightarrow \tilde{u} \quad \text{a.e. in } \Omega, \quad (13)$$

where the symbol “ $\rightharpoonup$ ” denotes weak convergence (see Theorem 1.31 in [9]).

**Step 4.** We have that  $\tilde{u} \in W_0^{1,2}(\Omega, \omega)$ . In fact, for  $F_k$  fixed, we have by (11) and (12), for all  $\varphi \in W_0^{1,2}(\Omega, \tilde{\omega}_1)$ ,

$$\begin{aligned} \int_\Omega u_m \varphi \tilde{\omega}_1 \, dx &\rightarrow \int_\Omega \tilde{u} \varphi \tilde{\omega}_1 \, dx, \\ \int_\Omega D_i u_m D_i \varphi \tilde{\omega}_1 \, dx &\rightarrow \int_\Omega D_i \tilde{u} D_i \varphi \tilde{\omega}_1 \, dx. \end{aligned}$$

If  $\psi \in W_0^{1,2}(\Omega, \omega)$ , then  $\varphi = \psi \chi_{F_k} \in W_0^{1,2}(\Omega, \tilde{\omega}_1)$  (since  $\omega \sim \tilde{\omega}_1$  in  $F_k$ , i.e., there is a constant  $c > 0$  such that  $\tilde{\omega}_1 \leq c\omega$  in  $F_k$ , and  $\chi_E$  denotes the characteristic function of a measurable set  $E \subset \mathbb{R}^n$ ) and

$$\begin{aligned} \int_\Omega \varphi^2 \tilde{\omega}_1 \, dx &= \int_{F_k} \psi^2 \tilde{\omega}_1 \, dx \leq c \int_{F_k} \psi^2 \omega \, dx \leq c \int_\Omega \psi^2 \omega \, dx < \infty, \\ \int_\Omega (D_i \varphi)^2 \tilde{\omega}_1 \, dx &= \int_{F_k} (D_i \psi)^2 \tilde{\omega}_1 \, dx \leq c \int_{F_k} (D_i \psi)^2 \omega \, dx \leq c \int_\Omega (D_i \psi)^2 \omega \, dx < \infty. \end{aligned}$$



Consequently,

$$\begin{aligned} \int_{\Omega} u_m \psi \chi_{F_k} \tilde{\omega}_1 \, dx &\rightarrow \int_{\Omega} \tilde{u} \psi \chi_{F_k} \tilde{\omega}_1 \, dx, \\ \int_{\Omega} D_i u_m D_i \psi \chi_{F_k} \tilde{\omega}_1 \, dx &\rightarrow \int_{\Omega} D_i \tilde{u} D_i \psi \chi_{F_k} \tilde{\omega}_1 \, dx, \end{aligned}$$

for all  $\psi \in W_0^{1,2}(\Omega, \omega)$ , that is, the sequence  $\{u_m \chi_{F_k}\}$  is weakly convergent in  $W_0^{1,2}(\Omega, \omega)$ . Therefore, we have

$$\|\nabla \tilde{u}\|_{L^2(F_k, \omega)}^2 = \int_{F_k} |\nabla \tilde{u}|^2 \omega \, dx \leq \limsup_{m \rightarrow \infty} \int_{F_k} |\nabla u_m|^2 \omega \, dx,$$

and for  $m \geq k$  we have  $\omega = \omega_m$  in  $F_k$ . Hence, by (10), we obtain

$$\begin{aligned} \|\nabla \tilde{u}\|_{L^2(F_k, \omega)}^2 &\leq \limsup_{m \rightarrow \infty} \int_{F_k} |\nabla u_m|^2 \omega \, dx \\ &= \limsup_{m \rightarrow \infty} \int_{F_k} |\nabla u_m|^2 \omega_m \, dx \\ &\leq \limsup_{m \rightarrow \infty} \int_{\Omega} |\nabla u_m|^2 \omega_m \, dx \leq C_1^2. \end{aligned}$$

By the Monotone Convergence Theorem we obtain  $\|\nabla \tilde{u}\|_{L^2(\Omega, \omega)} \leq C_1$ . Therefore, we have  $\tilde{u} \in W_0^{1,2}(\Omega, \omega)$ .

**Step 5.** We need to show that  $\tilde{u}$  is a solution of problem (P), i.e., for every  $\varphi \in W_0^{1,2}(\Omega, \omega)$  we have

$$\begin{aligned} \int_{\Omega} a_{ij}(x) D_i \tilde{u}(x) D_j \varphi(x) \, dx - \int_{\Omega} \gamma \tilde{u}(x) g_1(x) \varphi(x) \, dx + \int_{\Omega} h(\tilde{u}(x)) g_2(x) \varphi(x) \, dx \\ = \int_{\Omega} f(x) \varphi(x) \, dx. \end{aligned}$$

Using the fact that  $u_m$  is a solution of  $(P_m)$ , we have

$$\begin{aligned} \int_{\Omega} a_{ij}^{mm}(x) D_i u_m(x) D_j \varphi(x) \, dx - \int_{\Omega} \gamma u_m(x) g_{1m}(x) \varphi(x) \, dx \\ + \int_{\Omega} h(u_m(x)) g_{2m}(x) \varphi(x) \, dx = \int_{\Omega} f_m(x) \varphi(x) \, dx, \end{aligned}$$

for every  $\varphi \in W_0^{1,2}(\Omega, \omega_m)$ . Moreover, over  $F_k$  (for  $m \geq k$ ) we have the following properties:

- (i)  $\omega = \omega_m$ ;
- (ii)  $g_{1m} = g_1$ ;

(iii)  $g_{2m} = g_2$ ;

(iv)  $f_m = f$ ;

(v)  $a_{ij}^{mm}(x) = a_{ij}(x)$ .

For  $\varphi \in W_0^{1,2}(\Omega, \omega)$  and  $k > 0$  (fixed), we define  $G_1, G_2: W_0^{1,2}(\Omega, \tilde{\omega}_1) \rightarrow \mathbb{R}$  by

$$G_1(u) = \int_{\Omega} a_{ij}(x) D_i u(x) D_j \varphi(x) \chi_{F_k} dx - \gamma \int_{\Omega} u(x) g_1(x) \varphi(x) \chi_{F_k}(x) dx,$$

$$G_2(u) = \int_{\Omega} h(u(x)) g_2(x) \varphi(x) \chi_{F_k}(x) dx.$$

(a) We have that  $G_1$  is linear and continuous functional. In fact, since the matrix  $\mathcal{A} = (a_{ij})$  is symmetric, we have

$$|\langle \mathcal{A} \nabla u, \nabla \varphi \rangle| \leq \langle \mathcal{A} \nabla u, \nabla u \rangle^{1/2} \langle \mathcal{A} \nabla \varphi, \nabla \varphi \rangle^{1/2},$$

where  $\langle \cdot, \cdot \rangle$  denotes here the Euclidian scalar product in  $\mathbb{R}^n$ . We also have  $\omega \sim \tilde{\omega}_1$  in  $F_k$  ( $\omega \leq c\tilde{\omega}_1$ ). By (2) and (H3) we obtain

$$\begin{aligned} |G_1(u)| &\leq \int_{F_k} |\langle \mathcal{A} \nabla u, \nabla \varphi \rangle| dx + \gamma \int_{F_k} |u| |g_1| |\varphi| dx \\ &\leq \int_{F_k} \langle \mathcal{A} \nabla u, \nabla u \rangle^{1/2} \langle \mathcal{A} \nabla \varphi, \nabla \varphi \rangle^{1/2} dx + \gamma \int_{F_k} |u| \frac{|g_1|}{\omega} |\varphi| \omega dx \\ &\leq \left( \int_{F_k} \langle \mathcal{A} \nabla u, \nabla u \rangle dx \right)^{1/2} \left( \int_{F_k} \langle \mathcal{A} \nabla \varphi, \nabla \varphi \rangle dx \right)^{1/2} \\ &\quad + \gamma \|g_1/\omega\|_{L^\infty(\Omega)} \left( \int_{F_k} |u|^2 \omega dx \right)^{1/2} \left( \int_{F_k} |\varphi|^2 \omega \right)^{1/2} \\ &\leq \Lambda \left( \int_{F_k} |\nabla u|^2 \omega dx \right)^{1/2} \left( \int_{\Omega} |\nabla \varphi|^2 \omega dx \right)^{1/2} \\ &\quad + \gamma \left( \int_{F_k} |u|^2 \omega dx \right)^{1/2} \left( \int_{\Omega} |\varphi|^2 \omega dx \right)^{1/2} \\ &\leq \Lambda \left( \int_{F_k} c |\nabla u|^2 \tilde{\omega}_1 dx \right)^{1/2} \left( \int_{\Omega} |\nabla \varphi|^2 \omega dx \right)^{1/2} \\ &\quad + \gamma \left( \int_{F_k} c |u|^2 \tilde{\omega}_1 dx \right)^{1/2} \left( \int_{\Omega} |\varphi|^2 \omega dx \right)^{1/2} \\ &\leq (\Lambda c^{1/2} + \gamma c^{1/2}) \|\varphi\|_{W_0^{1,2}(\Omega, \omega)} \|u\|_{W_0^{1,2}(\Omega, \tilde{\omega}_1)}. \end{aligned}$$

(b) We have that  $G_2$  is continuous functional. In fact, if  $u_1, u_2 \in W_0^{1,2}(\Omega, \tilde{\omega}_1)$ , we obtain by (H1) and (H3)

$$\begin{aligned} |G_2(u_2) - G_2(u_1)| &\leq \int_{F_k} |h(u_2) - h(u_1)| |g_2| |\varphi| dx \\ &\leq \int_{F_k} C_h |u_1 - u_2| \frac{|g_1|}{\omega} |\varphi| \omega dx \\ &\leq C_h \|g_2/\omega\|_{L^\infty(\Omega)} \left( \int_{\Omega} |\varphi|^2 \omega dx \right)^{1/2} \left( \int_{F_k} c |u_1 - u_2|^2 \tilde{\omega}_1 dx \right)^{1/2} \\ &\leq c^{1/2} C_h \|g_2/\omega\|_{L^\infty(\Omega)} \|\varphi\|_{W_0^{1,2}(\Omega, \omega)} \|u_1 - u_2\|_{W_0^{1,2}(\Omega, \tilde{\omega}_1)}. \end{aligned}$$

Using (a), (b), properties (i), (ii), (iii), (iv) and (v), and that  $u_m$  is solution of  $(P_m)$ , we obtain

$$\begin{aligned} &\int_{F_k} a_{ij} D_i \tilde{u} D_j \varphi dx - \gamma \int_{F_k} \tilde{u} g_1 \varphi dx + \int_{F_k} h(\tilde{u}) g_2 \varphi dx = \lim_{m \rightarrow \infty} [G_1(u_m) + G_2(u_m)] \\ &= \lim_{m \rightarrow \infty} \left( \int_{F_k} a_{ij}^{mm} D_i u_m D_j \varphi dx - \gamma \int_{F_k} u_m g_1 \varphi dx + \int_{F_k} h(u_m) g_2 \varphi dx \right) \\ &= \lim_{m \rightarrow \infty} \left( \int_{\Omega} a_{ij}^{mm} D_i u_m D_j \varphi dx - \gamma \int_{\Omega} u_m g_1 \varphi dx + \int_{\Omega} h(u_m) g_2 \varphi dx \right. \\ &\quad \left. - \int_{\Omega \cap F_k^c} a_{ij}^{mm} D_i u_m D_j \varphi dx + \gamma \int_{\Omega \cap F_k^c} u_m g_1 \varphi dx - \int_{\Omega \cap F_k^c} h(u_m) g_2 \varphi dx \right) \\ &= \lim_{m \rightarrow \infty} \left( \int_{\Omega} f_m \varphi dx - \int_{\Omega \cap F_k^c} a_{ij}^{mm} D_i u_m D_j \varphi dx + \gamma \int_{\Omega \cap F_k^c} u_m g_1 \varphi dx \right. \\ &\quad \left. - \int_{\Omega \cap F_k^c} h(u_m) g_2 \varphi dx \right), \end{aligned} \tag{14}$$

where  $E^c$  denotes the complement of a set  $E \subset \mathbb{R}^n$ .

(I) By the Lebesgue Dominated Convergence Theorem we obtain

$$\int_{\Omega} f_m \varphi dx \rightarrow \int_{\Omega} f \varphi dx. \tag{15}$$

(II) Since the matrix  $\mathcal{A}^m = (a_{ij}^{mm})$  is symmetric, we have

$$|\langle \mathcal{A}^m \nabla u_m, \nabla \varphi \rangle| \leq \langle \mathcal{A}^m \nabla u_m, \nabla u_m \rangle^{1/2} \langle \mathcal{A}^m \nabla \varphi, \nabla \varphi \rangle^{1/2}.$$

Then, by (2) and (10), we obtain

$$\begin{aligned}
\left| \int_{\Omega \cap F_k^c} a_{ij}^{mm} D_i u_m D_j \varphi \, dx \right| &\leq \int_{\Omega \cap F_k^c} |\langle \mathcal{A}^m \nabla u_m, \nabla \varphi \rangle| \, dx \\
&\leq \Lambda \left( \int_{\Omega \cap F_k^c} |\nabla u_m|^2 \omega_m \, dx \right)^{1/2} \left( \int_{\Omega \cap F_k^c} |\nabla \varphi|^2 \omega_m \, dx \right)^{1/2} \\
&\leq \Lambda \|u_m\|_{W_0^{1,2}(\Omega, \omega_m)} \left( \int_{\Omega \cap F_k^c} |\nabla \varphi|^2 \omega_m \, dx \right)^{1/2} \\
&\leq \Lambda C_1 \left( \int_{\Omega \cap F_k^c} |\nabla \varphi|^2 \omega_m \, dx \right)^{1/2}. \tag{16}
\end{aligned}$$

(III) By (H3) and (10), we obtain

$$\begin{aligned}
\left| \int_{\Omega \cap F_k^c} u_m g_{1m} \varphi \, dx \right| &\leq \int_{\Omega \cap F_k^c} |u_m| \frac{|g_{1m}|}{\omega_m} |\varphi| \omega_m \, dx \\
&\leq \|g_{1m}/\omega_m\|_{L^\infty(\Omega)} \left( \int_{\Omega} |u_m|^2 \omega_m \, dx \right)^{1/2} \left( \int_{\Omega \cap F_k^c} |\varphi|^2 \omega_m \, dx \right)^{1/2} \\
&\leq \|g_1/\omega\|_{L^\infty(\Omega)} \|u_m\|_{W_0^{1,2}(\Omega, \omega_m)} \left( \int_{\Omega \cap F_k^c} |\varphi|^2 \omega_m \, dx \right)^{1/2} \\
&\leq C_1 \|g_1/\omega\|_{L^\infty(\Omega)} \left( \int_{\Omega \cap F_k^c} |\varphi|^2 \omega_m \, dx \right)^{1/2}. \tag{17}
\end{aligned}$$

(IV) By (H1), (H3) and  $|h(t)| \leq C_h |t|$  (for all  $t \in \mathbb{R}$ ) we have

$$\begin{aligned}
\left| \int_{\Omega \cap F_k^c} h(u_m) g_{2m} \varphi \, dx \right| &\leq \int_{\Omega \cap F_k^c} |h(u_m)| \frac{|g_{2m}|}{\omega_m} |\varphi| \omega_m \, dx \\
&\leq C_h \|g_{2m}/\omega_m\|_{L^\infty(\Omega)} \int_{\Omega \cap F_k^c} |u_m| |\varphi| \omega_m \, dx \\
&\leq C_h \|g_2/\omega\|_{L^\infty(\Omega)} \left( \int_{\Omega \cap F_k^c} |u_m|^2 \omega_m \, dx \right)^{1/2} \left( \int_{\Omega \cap F_k^c} |\varphi|^2 \omega_m \, dx \right)^{1/2} \\
&\leq C_h \|g_2/\omega\|_{L^\infty(\Omega)} \|u_m\|_{W_0^{1,2}(\Omega, \omega_m)} \left( \int_{\Omega \cap F_k^c} |\varphi|^2 \omega_m \, dx \right)^{1/2} \\
&\leq C_h C_1 \|g_2/\omega\|_{L^\infty(\Omega)} \left( \int_{\Omega \cap F_k^c} |\varphi|^2 \omega_m \, dx \right)^{1/2} \tag{18}
\end{aligned}$$

Using Lemma 1, we know that  $|\Omega \cap F_k^c| \rightarrow 0$  when  $k \rightarrow \infty$ . Then

$$\lim_{k \rightarrow \infty} \left( \int_{\Omega \cap F_k^c} |\varphi|^2 \omega_m \, dx \right)^{1/2} = \lim_{k \rightarrow \infty} \left( \int_{\Omega \cap F_k^c} |\nabla \varphi|^2 \omega_m \, dx \right)^{1/2} = 0,$$

and we obtain in (16), (17) and (18)

$$\lim_{k \rightarrow \infty} \int_{\Omega \cap F_k^c} a_{ij}^{mm}(x) D_i u(x) D_j \varphi(x) \, dx = 0, \tag{19}$$

$$\lim_{k \rightarrow \infty} \int_{\Omega \cap F_k^c} u_m g_{1m} \varphi \, dx = 0, \tag{20}$$

$$\lim_{k \rightarrow \infty} \int_{\Omega \cap F_k^c} h(u_m(x)) g_{2m}(x) \varphi(x) \, dx = 0. \tag{21}$$

Therefore, by (14), (19), (20) and (21) we conclude, when  $k \rightarrow \infty$  (and  $m \geq k$ ),

$$\int_{\Omega} a_{ij} D_i \tilde{u} D_j \varphi(x) \, dx - \gamma \int_{\Omega} g_1 \tilde{u} \varphi \, dx + \int_{\Omega} h(\tilde{u}) g_2 \varphi \, dx = \int_{\Omega} f \varphi \, dx,$$

for all  $\varphi \in W_0^{1,2}(\Omega, \omega)$ , that is,  $\tilde{u}$  is a solution of problem (P). Therefore,  $u = \tilde{u}$  (by the uniqueness).

**Example 1.** Let  $\Omega = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 < 1\}$  and  $0 < a < b$ . By Theorem 1, with  $h(t) = \sin(t)$ ,  $f(x, y) = x|y|$ ,  $\omega(x, y) = (x^2 + y^2)^{-1/2}$ ,

$$g_1(x, y) = (x^2 + y^2)^{-1/2} \cos(xy),$$

$$g_2(x, y) = (x^2 + y^2)^{-1/4} \sin(xy)$$

and

$$A(x, y) = \begin{pmatrix} a(x^2 + y^2)^{-1/2} & 0 \\ 0 & b(x^2 + y^2)^{-1/2} \end{pmatrix},$$

the problem

$$\begin{cases} Lu(x) - \gamma u(x) g_1(x) + h(u(x)) g_2(x) = f(x) & \text{in } \Omega, \\ u(x) = 0 & \text{on } \partial\Omega, \end{cases}$$

where

$$Lu(x) = -\frac{\partial}{\partial x} \left( a(x^2 + y^2)^{-1/2} \frac{\partial u}{\partial x} \right) - \frac{\partial y}{\partial x} \left( b(x^2 + y^2)^{-1/2} \frac{\partial u}{\partial y} \right),$$

has a unique solution  $u \in W_0^{1,2}(\Omega, \omega)$  if  $\gamma > 0$  is not an eigenvalue of linearized problem (LP), and  $u$  can be approximated by a sequence of solutions for non-degenerate semilinear elliptic equations.

## References

- [1] A. C. Cavalheiro: An approximation theorem for solutions of degenerate elliptic equations. *Proc. Edinb. Math. Soc.* 45 (2002) 363–389. doi: 10.1017/S0013091500000079
- [2] A. C. Cavalheiro: Existence of solutions in weighted Sobolev spaces for some degenerate semilinear elliptic equations. *Appl. Math. Lett.* 17 (2004) 387–391. doi:10.1016/S0893-9659(04)00043-6

- [3] A. C. Cavalheiro: Existence results for the Dirichlet problem of some degenerate nonlinear elliptic equations. *J. Appl. Anal.* 20 (2) (2014) 145–154.  
doi:10.1515/jaa-2014-0016
- [4] A. C. Cavalheiro: Uniqueness of solutions for some degenerate nonlinear elliptic equations. *Appl. Math. (Warsaw)* 41 (1) (2014) 93–106.
- [5] A. C. Cavalheiro: Existence and uniqueness of solutions for the Navier problems with degenerate nonlinear elliptic equations. *Note Mat.* 25 (2) (2015) 1–16.
- [6] E. Fabes, C. Kenig, R. Serapioni: The local regularity of solutions of degenerate elliptic equations. *Comm. Partial Differential Equations* 7 (1982) 77–116.  
doi:10.1080/03605308208820218
- [7] J. C. Fernandes, B. Franchi: Existence and properties of the Green function for a class of degenerate parabolic equations. *Rev. Mat. Iberoam.* 12 (1996) 491–525.
- [8] J. Garcia-Cuerva, J. L. Rubio de Francia: Weighted Norm Inequalities and Related Topics. *North-Holland Mathematics Studies* 116 (1985).
- [9] J. Heinonen, T. Kilpeläinen, O. Martio: *Nonlinear Potential Theory of Degenerate Elliptic Equations*. Oxford Math. Monographs, Clarendon Press (1993).
- [10] A. Kufner: *Weighted Sobolev Spaces*. John Wiley & Sons, New York (1985).
- [11] B. Muckenhoupt: Weighted norm inequalities for the Hardy maximal function. *Trans. Amer. Math. Soc.* 165 (1972) 207–226.
- [12] A. Torchinsky: *Real-Variable Methods in Harmonic Analysis*. Academic Press, San Diego (1986).
- [13] B. O. Turesson: *Nonlinear Potential Theory and Weighted Sobolev Spaces*. Springer-Verlag (2000). Lecture Notes in Math.

*Author's address:*

DEPARTMENT OF MATHEMATICS, STATE UNIVERSITY OF LONDRINA, LONDRINA – PR –  
BRAZIL, 86057-970

*E-mail:* [accava@gmail.com](mailto:accava@gmail.com)

*Received:* 16 November, 2016

*Accepted for publication:* 16 March, 2017

*Communicated by:* Geoff Prince