

Several Differentiation Formulas of Special Functions. Part III

Bo Li
Qingdao University of Science
and Technology
China

Yan Zhang
Qingdao University of Science
and Technology
China

Xiquan Liang
Qingdao University of Science
and Technology
China

Summary. In this article, we give several differentiation formulas of special and composite functions including trigonometric function, inverse trigonometric function, polynomial function and logarithmic function.

MML identifier: FDIFF_7, version: 7.7.01 4.60.940

The articles [13], [15], [16], [1], [4], [10], [11], [17], [5], [14], [12], [2], [6], [9], [7], [8], and [3] provide the terminology and notation for this paper.

For simplicity, we follow the rules: x, r, a, b denote real numbers, n denotes a natural number, Z denotes an open subset of \mathbb{R} , and f, f_1, f_2, f_3 denote partial functions from \mathbb{R} to \mathbb{R} .

One can prove the following propositions:

- (1) $x_{\mathbb{Z}}^2 = x^2$.
- (2) If $x > 0$, then $x_{\mathbb{R}}^{\frac{1}{2}} = \sqrt{x}$.
- (3) If $x > 0$, then $x_{\mathbb{R}}^{-\frac{1}{2}} = \frac{1}{\sqrt{x}}$.
- (4) Suppose $Z \subseteq]-1, 1[$ and $Z \subseteq \text{dom}(r$ (the function arcsin)). Then
 - (i) r (the function arcsin) is differentiable on Z , and
 - (ii) for every x such that $x \in Z$ holds $(r$ (the function arcsin))' $\big|_Z(x) = \frac{r}{\sqrt{1-x^2}}$.

- (5) Suppose $Z \subseteq]-1, 1[$ and $Z \subseteq \text{dom}(r$ (the function arccos)). Then
- (i) r (the function arccos) is differentiable on Z , and
 - (ii) for every x such that $x \in Z$ holds $(r$ (the function arccos))' $_{|Z}(x) = -\frac{x}{\sqrt{1-x^2}}$.
- (6) Suppose f is differentiable in x and $f(x) > -1$ and $f(x) < 1$. Then (the function arcsin) $\cdot f$ is differentiable in x and ((the function arcsin) $\cdot f$)'(x) = $\frac{f'(x)}{\sqrt{1-f(x)^2}}$.
- (7) Suppose f is differentiable in x and $f(x) > -1$ and $f(x) < 1$. Then (the function arccos) $\cdot f$ is differentiable in x and ((the function arccos) $\cdot f$)'(x) = $-\frac{f'(x)}{\sqrt{1-f(x)^2}}$.
- (8) Suppose $Z \subseteq \text{dom}(\log_-(e) \cdot$ (the function arcsin)) and $Z \subseteq]-1, 1[$ and for every x such that $x \in Z$ holds (the function arcsin)(x) > 0. Then
- (i) $\log_-(e) \cdot$ (the function arcsin) is differentiable on Z , and
 - (ii) for every x such that $x \in Z$ holds $(\log_-(e) \cdot$ (the function arcsin))' $_{|Z}(x) = \frac{1}{\sqrt{1-x^2} \cdot (\text{the function arcsin})(x)}$.
- (9) Suppose $Z \subseteq \text{dom}(\log_-(e) \cdot$ (the function arccos)) and $Z \subseteq]-1, 1[$ and for every x such that $x \in Z$ holds (the function arccos)(x) > 0. Then
- (i) $\log_-(e) \cdot$ (the function arccos) is differentiable on Z , and
 - (ii) for every x such that $x \in Z$ holds $(\log_-(e) \cdot$ (the function arccos))' $_{|Z}(x) = -\frac{1}{\sqrt{1-x^2} \cdot (\text{the function arccos})(x)}$.
- (10) Suppose $Z \subseteq \text{dom}(\binom{n}{Z} \cdot$ (the function arcsin)) and $Z \subseteq]-1, 1[$. Then
- (i) $\binom{n}{Z} \cdot$ (the function arcsin) is differentiable on Z , and
 - (ii) for every x such that $x \in Z$ holds $(\binom{n}{Z} \cdot$ (the function arcsin))' $_{|Z}(x) = \frac{n \cdot (\text{the function arcsin})(x)_{|Z}^{n-1}}{\sqrt{1-x^2}}$.
- (11) Suppose $Z \subseteq \text{dom}(\binom{n}{Z} \cdot$ (the function arccos)) and $Z \subseteq]-1, 1[$. Then
- (i) $\binom{n}{Z} \cdot$ (the function arccos) is differentiable on Z , and
 - (ii) for every x such that $x \in Z$ holds $(\binom{n}{Z} \cdot$ (the function arccos))' $_{|Z}(x) = -\frac{n \cdot (\text{the function arccos})(x)_{|Z}^{n-1}}{\sqrt{1-x^2}}$.
- (12) Suppose $Z \subseteq \text{dom}(\frac{1}{2} (\binom{2}{Z} \cdot$ (the function arcsin))) and $Z \subseteq]-1, 1[$. Then
- (i) $\frac{1}{2} (\binom{2}{Z} \cdot$ (the function arcsin)) is differentiable on Z , and
 - (ii) for every x such that $x \in Z$ holds $(\frac{1}{2} (\binom{2}{Z} \cdot$ (the function arcsin)))' $_{|Z}(x) = \frac{(\text{the function arcsin})(x)}{\sqrt{1-x^2}}$.
- (13) Suppose $Z \subseteq \text{dom}(\frac{1}{2} (\binom{2}{Z} \cdot$ (the function arccos))) and $Z \subseteq]-1, 1[$. Then
- (i) $\frac{1}{2} (\binom{2}{Z} \cdot$ (the function arccos)) is differentiable on Z , and
 - (ii) for every x such that $x \in Z$ holds $(\frac{1}{2} (\binom{2}{Z} \cdot$ (the function arccos)))' $_{|Z}(x) = -\frac{(\text{the function arccos})(x)}{\sqrt{1-x^2}}$.

- (14) Suppose $Z \subseteq \text{dom}(\text{(the function arcsin)} \cdot f)$ and for every x such that $x \in Z$ holds $f(x) = a \cdot x + b$ and $f(x) > -1$ and $f(x) < 1$. Then
- (the function arcsin) $\cdot f$ is differentiable on Z , and
 - for every x such that $x \in Z$ holds $((\text{the function arcsin}) \cdot f)'_{|Z}(x) = \frac{a}{\sqrt{1-(a \cdot x + b)^2}}$.
- (15) Suppose $Z \subseteq \text{dom}(\text{(the function arccos)} \cdot f)$ and for every x such that $x \in Z$ holds $f(x) = a \cdot x + b$ and $f(x) > -1$ and $f(x) < 1$. Then
- (the function arccos) $\cdot f$ is differentiable on Z , and
 - for every x such that $x \in Z$ holds $((\text{the function arccos}) \cdot f)'_{|Z}(x) = -\frac{a}{\sqrt{1-(a \cdot x + b)^2}}$.
- (16) Suppose $Z \subseteq \text{dom}(\text{id}_Z (\text{the function arcsin}))$ and $Z \subseteq]-1, 1[$. Then
- $\text{id}_Z (\text{the function arcsin})$ is differentiable on Z , and
 - for every x such that $x \in Z$ holds $(\text{id}_Z (\text{the function arcsin}))'_{|Z}(x) = (\text{the function arcsin})(x) + \frac{x}{\sqrt{1-x^2}}$.
- (17) Suppose $Z \subseteq \text{dom}(\text{id}_Z (\text{the function arccos}))$ and $Z \subseteq]-1, 1[$. Then
- $\text{id}_Z (\text{the function arccos})$ is differentiable on Z , and
 - for every x such that $x \in Z$ holds $(\text{id}_Z (\text{the function arccos}))'_{|Z}(x) = (\text{the function arccos})(x) - \frac{x}{\sqrt{1-x^2}}$.
- (18) Suppose $Z \subseteq \text{dom}(f (\text{the function arcsin}))$ and $Z \subseteq]-1, 1[$ and for every x such that $x \in Z$ holds $f(x) = a \cdot x + b$. Then
- $f (\text{the function arcsin})$ is differentiable on Z , and
 - for every x such that $x \in Z$ holds $(f (\text{the function arcsin}))'_{|Z}(x) = a \cdot (\text{the function arcsin})(x) + \frac{a \cdot x + b}{\sqrt{1-x^2}}$.
- (19) Suppose $Z \subseteq \text{dom}(f (\text{the function arccos}))$ and $Z \subseteq]-1, 1[$ and for every x such that $x \in Z$ holds $f(x) = a \cdot x + b$. Then
- $f (\text{the function arccos})$ is differentiable on Z , and
 - for every x such that $x \in Z$ holds $(f (\text{the function arccos}))'_{|Z}(x) = a \cdot (\text{the function arccos})(x) - \frac{a \cdot x + b}{\sqrt{1-x^2}}$.
- (20) Suppose $Z \subseteq \text{dom}(\frac{1}{2} ((\text{the function arcsin}) \cdot f))$ and for every x such that $x \in Z$ holds $f(x) = 2 \cdot x$ and $f(x) > -1$ and $f(x) < 1$. Then
- $\frac{1}{2} ((\text{the function arcsin}) \cdot f)$ is differentiable on Z , and
 - for every x such that $x \in Z$ holds $(\frac{1}{2} ((\text{the function arcsin}) \cdot f))'_{|Z}(x) = \frac{1}{\sqrt{1-(2 \cdot x)^2}}$.
- (21) Suppose $Z \subseteq \text{dom}(\frac{1}{2} ((\text{the function arccos}) \cdot f))$ and for every x such that $x \in Z$ holds $f(x) = 2 \cdot x$ and $f(x) > -1$ and $f(x) < 1$. Then
- $\frac{1}{2} ((\text{the function arccos}) \cdot f)$ is differentiable on Z , and
 - for every x such that $x \in Z$ holds $(\frac{1}{2} ((\text{the function arccos}) \cdot f))'_{|Z}(x) = -\frac{1}{\sqrt{1-(2 \cdot x)^2}}$.

(22) Suppose $Z \subseteq \text{dom}((\frac{1}{\mathbb{R}}) \cdot f)$ and $f = f_1 - f_2$ and $f_2 = \frac{2}{Z}$ and for every x such that $x \in Z$ holds $f_1(x) = 1$ and $f(x) > 0$. Then $(\frac{1}{\mathbb{R}}) \cdot f$ is differentiable on Z and for every x such that $x \in Z$ holds $((\frac{1}{\mathbb{R}}) \cdot f)'_{|Z}(x) = -x \cdot (1 - x^2_{\mathbb{Z}})_{\mathbb{R}}^{-\frac{1}{2}}$.

(23) Suppose that

- (i) $Z \subseteq \text{dom}(\text{id}_Z(\text{the function arcsin}) + (\frac{1}{\mathbb{R}}) \cdot f)$,
- (ii) $Z \subseteq]-1, 1[$,
- (iii) $f = f_1 - f_2$,
- (iv) $f_2 = \frac{2}{Z}$, and
- (v) for every x such that $x \in Z$ holds $f_1(x) = 1$ and $f(x) > 0$ and $x \neq 0$.

Then

- (vi) $\text{id}_Z(\text{the function arcsin}) + (\frac{1}{\mathbb{R}}) \cdot f$ is differentiable on Z , and
- (vii) for every x such that $x \in Z$ holds $(\text{id}_Z(\text{the function arcsin}) + (\frac{1}{\mathbb{R}}) \cdot f)'_{|Z}(x) = (\text{the function arcsin})(x)$.

(24) Suppose that

- (i) $Z \subseteq \text{dom}(\text{id}_Z(\text{the function arccos}) - (\frac{1}{\mathbb{R}}) \cdot f)$,
- (ii) $Z \subseteq]-1, 1[$,
- (iii) $f = f_1 - f_2$,
- (iv) $f_2 = \frac{2}{Z}$, and
- (v) for every x such that $x \in Z$ holds $f_1(x) = 1$ and $f(x) > 0$ and $x \neq 0$.

Then

- (vi) $\text{id}_Z(\text{the function arccos}) - (\frac{1}{\mathbb{R}}) \cdot f$ is differentiable on Z , and
- (vii) for every x such that $x \in Z$ holds $(\text{id}_Z(\text{the function arccos}) - (\frac{1}{\mathbb{R}}) \cdot f)'_{|Z}(x) = (\text{the function arccos})(x)$.

(25) Suppose $Z \subseteq \text{dom}(\text{id}_Z((\text{the function arcsin}) \cdot f))$ and for every x such that $x \in Z$ holds $f(x) = \frac{x}{a}$ and $f(x) > -1$ and $f(x) < 1$. Then

- (i) $\text{id}_Z((\text{the function arcsin}) \cdot f)$ is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $(\text{id}_Z((\text{the function arcsin}) \cdot f))'_{|Z}(x) = (\text{the function arcsin})(\frac{x}{a}) + \frac{x}{a \cdot \sqrt{1 - (\frac{x}{a})^2}}$.

(26) Suppose $Z \subseteq \text{dom}(\text{id}_Z((\text{the function arccos}) \cdot f))$ and for every x such that $x \in Z$ holds $f(x) = \frac{x}{a}$ and $f(x) > -1$ and $f(x) < 1$. Then

- (i) $\text{id}_Z((\text{the function arccos}) \cdot f)$ is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $(\text{id}_Z((\text{the function arccos}) \cdot f))'_{|Z}(x) = (\text{the function arccos})(\frac{x}{a}) - \frac{x}{a \cdot \sqrt{1 - (\frac{x}{a})^2}}$.

(27) Suppose $Z \subseteq \text{dom}((\frac{1}{\mathbb{R}}) \cdot f)$ and $f = f_1 - f_2$ and $f_2 = \frac{2}{Z}$ and for every x such that $x \in Z$ holds $f_1(x) = a^2$ and $f(x) > 0$. Then $(\frac{1}{\mathbb{R}}) \cdot f$ is differentiable on Z and for every x such that $x \in Z$ holds $((\frac{1}{\mathbb{R}}) \cdot f)'_{|Z}(x) = -x \cdot (a^2 - x^2_{\mathbb{Z}})_{\mathbb{R}}^{-\frac{1}{2}}$.

(28) Suppose that

- (i) $Z \subseteq \text{dom}(\text{id}_Z ((\text{the function arcsin}) \cdot f_3) + (\frac{1}{\mathbb{R}}) \cdot f)$,
- (ii) $Z \subseteq]-1, 1[$,
- (iii) $f = f_1 - f_2$,
- (iv) $f_2 = \frac{2}{Z}$, and
- (v) for every x such that $x \in Z$ holds $f_1(x) = a^2$ and $f(x) > 0$ and $f_3(x) = \frac{x}{a}$ and $f_3(x) > -1$ and $f_3(x) < 1$ and $x \neq 0$ and $a > 0$.

Then

- (vi) $\text{id}_Z ((\text{the function arcsin}) \cdot f_3) + (\frac{1}{\mathbb{R}}) \cdot f$ is differentiable on Z , and
- (vii) for every x such that $x \in Z$ holds $(\text{id}_Z ((\text{the function arcsin}) \cdot f_3) + (\frac{1}{\mathbb{R}}) \cdot f)'_{|Z}(x) = (\text{the function arcsin})(\frac{x}{a})$.

(29) Suppose that

- (i) $Z \subseteq \text{dom}(\text{id}_Z ((\text{the function arccos}) \cdot f_3) - (\frac{1}{\mathbb{R}}) \cdot f)$,
- (ii) $Z \subseteq]-1, 1[$,
- (iii) $f = f_1 - f_2$,
- (iv) $f_2 = \frac{2}{Z}$, and
- (v) for every x such that $x \in Z$ holds $f_1(x) = a^2$ and $f(x) > 0$ and $f_3(x) = \frac{x}{a}$ and $f_3(x) > -1$ and $f_3(x) < 1$ and $x \neq 0$ and $a > 0$.

Then

- (vi) $\text{id}_Z ((\text{the function arccos}) \cdot f_3) - (\frac{1}{\mathbb{R}}) \cdot f$ is differentiable on Z , and
- (vii) for every x such that $x \in Z$ holds $(\text{id}_Z ((\text{the function arccos}) \cdot f_3) - (\frac{1}{\mathbb{R}}) \cdot f)'_{|Z}(x) = (\text{the function arccos})(\frac{x}{a})$.

(30) Suppose $Z \subseteq \text{dom}((-\frac{1}{n}) ((\frac{n}{Z}) \cdot \frac{1}{\text{the function sin}}))$ and $n > 0$ and for every x such that $x \in Z$ holds $(\text{the function sin})(x) \neq 0$. Then

- (i) $(-\frac{1}{n}) ((\frac{n}{Z}) \cdot \frac{1}{\text{the function sin}})$ is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $((-\frac{1}{n}) ((\frac{n}{Z}) \cdot \frac{1}{\text{the function sin}}))'_{|Z}(x) = \frac{(\text{the function cos})(x)}{(\text{the function sin})(x)_Z^{n+1}}$.

(31) Suppose $Z \subseteq \text{dom}(\frac{1}{n} ((\frac{n}{Z}) \cdot \frac{1}{\text{the function cos}}))$ and $n > 0$ and for every x such that $x \in Z$ holds $(\text{the function cos})(x) \neq 0$. Then

- (i) $\frac{1}{n} ((\frac{n}{Z}) \cdot \frac{1}{\text{the function cos}})$ is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $(\frac{1}{n} ((\frac{n}{Z}) \cdot \frac{1}{\text{the function cos}}))'_{|Z}(x) = \frac{(\text{the function sin})(x)}{(\text{the function cos})(x)_Z^{n+1}}$.

(32) Suppose $Z \subseteq \text{dom}((\text{the function sin}) \cdot \log_-(e))$ and for every x such that $x \in Z$ holds $x > 0$. Then

- (i) $(\text{the function sin}) \cdot \log_-(e)$ is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $((\text{the function sin}) \cdot \log_-(e))'_{|Z}(x) = \frac{(\text{the function cos})((\log_-(e))(x))}{x}$.

(33) Suppose $Z \subseteq \text{dom}((\text{the function cos}) \cdot \log_-(e))$ and for every x such that $x \in Z$ holds $x > 0$. Then

- (i) (the function \cos) $\cdot \log_-(e)$ is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $((\text{the function } \cos) \cdot \log_-(e))'_{|Z}(x) = -\frac{(\text{the function } \sin)((\log_-(e))(x))}{x}$.
- (34) Suppose $Z \subseteq \text{dom}((\text{the function } \sin) \cdot (\text{the function } \exp))$. Then
- (i) (the function \sin) \cdot (the function \exp) is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $((\text{the function } \sin) \cdot (\text{the function } \exp))'_{|Z}(x) = (\text{the function } \exp)(x) \cdot (\text{the function } \cos)((\text{the function } \exp)(x))$.
- (35) Suppose $Z \subseteq \text{dom}((\text{the function } \cos) \cdot (\text{the function } \exp))$. Then
- (i) (the function \cos) \cdot (the function \exp) is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $((\text{the function } \cos) \cdot (\text{the function } \exp))'_{|Z}(x) = -(\text{the function } \exp)(x) \cdot (\text{the function } \sin)((\text{the function } \exp)(x))$.
- (36) Suppose $Z \subseteq \text{dom}((\text{the function } \exp) \cdot (\text{the function } \cos))$. Then
- (i) (the function \exp) \cdot (the function \cos) is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $((\text{the function } \exp) \cdot (\text{the function } \cos))'_{|Z}(x) = -(\text{the function } \exp)((\text{the function } \cos)(x)) \cdot (\text{the function } \sin)(x)$.
- (37) Suppose $Z \subseteq \text{dom}((\text{the function } \exp) \cdot (\text{the function } \sin))$. Then
- (i) (the function \exp) \cdot (the function \sin) is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $((\text{the function } \exp) \cdot (\text{the function } \sin))'_{|Z}(x) = (\text{the function } \exp)((\text{the function } \sin)(x)) \cdot (\text{the function } \cos)(x)$.
- (38) Suppose $Z \subseteq \text{dom}((\text{the function } \sin) + (\text{the function } \cos))$. Then
- (i) (the function \sin) + (the function \cos) is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $((\text{the function } \sin) + (\text{the function } \cos))'_{|Z}(x) = (\text{the function } \cos)(x) - (\text{the function } \sin)(x)$.
- (39) Suppose $Z \subseteq \text{dom}((\text{the function } \sin) - (\text{the function } \cos))$. Then
- (i) (the function \sin) - (the function \cos) is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $((\text{the function } \sin) - (\text{the function } \cos))'_{|Z}(x) = (\text{the function } \cos)(x) + (\text{the function } \sin)(x)$.
- (40) Suppose $Z \subseteq \text{dom}((\text{the function } \exp) ((\text{the function } \sin) - (\text{the function } \cos)))$. Then
- (i) (the function \exp) $((\text{the function } \sin) - (\text{the function } \cos))$ is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $((\text{the function } \exp) ((\text{the function } \sin) - (\text{the function } \cos)))'_{|Z}(x) = 2 \cdot (\text{the function } \exp)(x) \cdot (\text{the function } \sin)(x)$.
- (41) Suppose $Z \subseteq \text{dom}((\text{the function } \exp) ((\text{the function } \sin) + (\text{the function } \cos)))$. Then

- (i) (the function exp) ((the function sin)+(the function cos)) is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds ((the function exp) ((the function sin)+(the function cos)))' $\upharpoonright_Z(x) = 2 \cdot$ (the function exp)(x) \cdot (the function cos)(x).
- (42) Suppose $Z \subseteq \text{dom}(\frac{\text{(the function sin)+(the function cos)}}{\text{the function exp}})$. Then
- (i) $\frac{\text{(the function sin)+(the function cos)}}{\text{the function exp}}$ is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $(\frac{\text{(the function sin)+(the function cos)}}{\text{the function exp}})' \upharpoonright_Z(x) = \frac{2 \cdot \text{(the function sin)}(x)}{\text{(the function exp)}(x)}$.
- (43) Suppose $Z \subseteq \text{dom}(\frac{\text{(the function sin)-(the function cos)}}{\text{the function exp}})$. Then
- (i) $\frac{\text{(the function sin)-(the function cos)}}{\text{the function exp}}$ is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $(\frac{\text{(the function sin)-(the function cos)}}{\text{the function exp}})' \upharpoonright_Z(x) = \frac{2 \cdot \text{(the function cos)}(x)}{\text{(the function exp)}(x)}$.
- (44) Suppose $Z \subseteq \text{dom}(\text{(the function exp) (the function sin)})$. Then
- (i) (the function exp) (the function sin) is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds ((the function exp) (the function sin))' $\upharpoonright_Z(x) =$ (the function exp)(x) \cdot ((the function sin)(x) + (the function cos)(x)).
- (45) Suppose $Z \subseteq \text{dom}(\text{(the function exp) (the function cos)})$. Then
- (i) (the function exp) (the function cos) is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds ((the function exp) (the function cos))' $\upharpoonright_Z(x) =$ (the function exp)(x) \cdot ((the function cos)(x) - (the function sin)(x)).
- (46) Suppose (the function cos)(x) $\neq 0$. Then
- (i) $\frac{\text{the function sin}}{\text{the function cos}}$ is differentiable in x , and
- (ii) $(\frac{\text{the function sin}}{\text{the function cos}})'(x) = \frac{1}{\text{(the function cos)}(x)^2}$.
- (47) Suppose (the function sin)(x) $\neq 0$. Then
- (i) $\frac{\text{the function cos}}{\text{the function sin}}$ is differentiable in x , and
- (ii) $(\frac{\text{the function cos}}{\text{the function sin}})'(x) = -\frac{1}{\text{(the function sin)}(x)^2}$.
- (48) Suppose $Z \subseteq \text{dom}(\frac{2}{Z} \cdot \frac{\text{the function sin}}{\text{the function cos}})$ and for every x such that $x \in Z$ holds (the function cos)(x) $\neq 0$. Then
- (i) $\frac{2}{Z} \cdot \frac{\text{the function sin}}{\text{the function cos}}$ is differentiable on Z , and
- (ii) for every x such that $x \in Z$ holds $(\frac{2}{Z} \cdot \frac{\text{the function sin}}{\text{the function cos}})' \upharpoonright_Z(x) = \frac{2 \cdot \text{(the function sin)}(x)}{\text{(the function cos)}(x)^{\frac{3}{Z}}}$.
- (49) Suppose $Z \subseteq \text{dom}(\frac{2}{Z} \cdot \frac{\text{the function cos}}{\text{the function sin}})$ and for every x such that $x \in Z$ holds (the function sin)(x) $\neq 0$. Then
- (i) $\frac{2}{Z} \cdot \frac{\text{the function cos}}{\text{the function sin}}$ is differentiable on Z , and

- (ii) for every x such that $x \in Z$ holds $\left(\frac{2}{Z}\right) \cdot \frac{\text{the function cos}}{\text{the function sin}} \Big|_Z(x) = \frac{2 \cdot (\text{the function cos})(x)}{(\text{the function sin})(x)^3}$.
- (50) Suppose that
- (i) $Z \subseteq \text{dom}\left(\frac{\text{the function sin}}{\text{the function cos}} \cdot f\right)$, and
- (ii) for every x such that $x \in Z$ holds $f(x) = \frac{x}{2}$ and $(\text{the function cos})(f(x)) \neq 0$.
- Then
- (iii) $\frac{\text{the function sin}}{\text{the function cos}} \cdot f$ is differentiable on Z , and
- (iv) for every x such that $x \in Z$ holds $\left(\frac{\text{the function sin}}{\text{the function cos}} \cdot f\right) \Big|_Z(x) = \frac{1}{1 + (\text{the function cos})(x)}$.
- (51) Suppose that
- (i) $Z \subseteq \text{dom}\left(\frac{\text{the function cos}}{\text{the function sin}} \cdot f\right)$, and
- (ii) for every x such that $x \in Z$ holds $f(x) = \frac{x}{2}$ and $(\text{the function sin})(f(x)) \neq 0$.
- Then
- (iii) $\frac{\text{the function cos}}{\text{the function sin}} \cdot f$ is differentiable on Z , and
- (iv) for every x such that $x \in Z$ holds $\left(\frac{\text{the function cos}}{\text{the function sin}} \cdot f\right) \Big|_Z(x) = \frac{1}{1 - (\text{the function cos})(x)}$.

REFERENCES

- [1] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [2] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [3] Artur Korniłowicz and Yasunari Shidama. Inverse trigonometric functions arcsin and arccos. *Formalized Mathematics*, 13(1):73–79, 2005.
- [4] Jarosław Kotowicz. Partial functions from a domain to a domain. *Formalized Mathematics*, 1(4):697–702, 1990.
- [5] Jarosław Kotowicz. Partial functions from a domain to the set of real numbers. *Formalized Mathematics*, 1(4):703–709, 1990.
- [6] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(2):269–272, 1990.
- [7] Konrad Raczkowski. Integer and rational exponents. *Formalized Mathematics*, 2(1):125–130, 1991.
- [8] Konrad Raczkowski and Andrzej Nędzusiak. Real exponents and logarithms. *Formalized Mathematics*, 2(2):213–216, 1991.
- [9] Konrad Raczkowski and Paweł Sadowski. Real function differentiability. *Formalized Mathematics*, 1(4):797–801, 1990.
- [10] Konrad Raczkowski and Paweł Sadowski. Topological properties of subsets in real numbers. *Formalized Mathematics*, 1(4):777–780, 1990.
- [11] Yasunari Shidama. The Taylor expansions. *Formalized Mathematics*, 12(2):195–200, 2004.
- [12] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics*.
- [13] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [14] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers. *Formalized Mathematics*, 1(3):445–449, 1990.
- [15] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [16] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.

- [17] Yuguang Yang and Yasunari Shidama. Trigonometric functions and existence of circle ratio. *Formalized Mathematics*, 7(2):255–263, 1998.

Received March 22, 2006
