

# Recognizing Chordal Graphs: Lex BFS and MCS<sup>1</sup>

Broderick Arneson  
University of Alberta  
Edmonton, Canada

Piotr Rudnicki  
University of Alberta  
Edmonton, Canada

**Summary.** We are formalizing the algorithm for recognizing chordal graphs by lexicographic breadth-first search as presented in [13, Section 3 of Chapter 4, pp. 81–84]. Then we follow with a formalization of another algorithm serving the same end but based on maximum cardinality search as presented by Tarjan and Yannakakis [25].

This work is a part of the MSc work of the first author under supervision of the second author. We would like to thank one of the anonymous reviewers for very useful suggestions.

MML identifier: LEXBFS, version: 7.8.03 4.75.958

The notation and terminology used in this paper are introduced in the following articles: [28], [11], [26], [32], [33], [35], [30], [10], [7], [8], [20], [29], [4], [2], [14], [23], [12], [3], [6], [9], [18], [15], [19], [16], [17], [24], [21], [1], [5], [31], [27], [22], and [34].

## 1. PRELIMINARIES

The following propositions are true:

- (1) Let  $A, B$  be elements of  $\mathbb{N}$ ,  $X$  be a non empty set, and  $F$  be a function from  $\mathbb{N}$  into  $X$ . If  $F$  is one-to-one, then  $\overline{\{F(w); w \text{ ranges over elements of } \mathbb{N}: A \leq w \wedge w \leq A + B\}} = B + 1$ .
- (2) For all natural numbers  $n, m, k$  such that  $m \leq k$  and  $n < m$  holds  $k -' m < k -' n$ .

---

<sup>1</sup>This work has been partially supported by the NSERC grant OGP 9207.

- (3) For all natural numbers  $n, k$  such that  $n < k$  holds  $(k -' (n + 1)) + 1 = k -' n$ .
- (4) For all natural numbers  $n, m, k$  such that  $k \neq 0$  holds  $(n + m \cdot k) \div k = (n \div k) + m$ .

Let  $S$  be a set. We say that  $S$  has finite elements if and only if:

(Def. 1) Every element of  $S$  is finite.

Let us note that there exists a set which is non empty and has finite elements and there exists a subset of  $2^{\mathbb{N}}$  which is non empty and finite and has finite elements.

Let  $S$  be a set with finite elements. One can check that every element of  $S$  is finite.

Let  $f, g$  be functions. The functor  $f[\cup]g$  yielding a function is defined by:

(Def. 2)  $\text{dom}(f[\cup]g) = \text{dom } f \cup \text{dom } g$  and for every set  $x$  such that  $x \in \text{dom } f \cup \text{dom } g$  holds  $(f[\cup]g)(x) = f(x) \cup g(x)$ .

The following three propositions are true:

- (5) For all natural numbers  $m, n, k$  holds  $m \in \text{Seg } k \setminus \text{Seg}(k -' n)$  iff  $k -' n < m$  and  $m \leq k$ .
- (6) For all natural numbers  $n, k, m$  such that  $n \leq m$  holds  $\text{Seg } k \setminus \text{Seg}(k -' n) \subseteq \text{Seg } k \setminus \text{Seg}(k -' m)$ .
- (7) For all natural numbers  $n, k$  such that  $n < k$  holds  $(\text{Seg } k \setminus \text{Seg}(k -' n)) \cup \{k -' n\} = \text{Seg } k \setminus \text{Seg}(k -' (n + 1))$ .

Let  $f$  be a binary relation. We say that  $f$  is natsubset yielding if and only if:

(Def. 3)  $\text{rng } f \subseteq 2^{\mathbb{N}}$ .

Let us mention that there exists a function which is finite-yielding and natsubset yielding.

Let  $f$  be a finite-yielding natsubset yielding function and let  $x$  be a set. Then  $f(x)$  is a finite subset of  $\mathbb{N}$ .

One can prove the following proposition

- (8) For every ordinal number  $X$  and for all finite subsets  $a, b$  of  $X$  such that  $a \neq b$  holds  $(a, 1)\text{-bag} \neq (b, 1)\text{-bag}$ .

Let  $F$  be a natural-yielding function, let  $S$  be a set, and let  $k$  be a natural number. The functor  $F.\text{incSubset}(S, k)$  yielding a natural-yielding function is defined by the conditions (Def. 4).

- (Def. 4)(i)  $\text{dom}(F.\text{incSubset}(S, k)) = \text{dom } F$ , and
- (ii) for every set  $y$  holds if  $y \in S$  and  $y \in \text{dom } F$ , then  $(F.\text{incSubset}(S, k))(y) = F(y) + k$  and if  $y \notin S$ , then  $(F.\text{incSubset}(S, k))(y) = F(y)$ .

Let  $n$  be an ordinal number, let  $T$  be a connected term order of  $n$ , and let  $B$  be a non empty finite subset of Bags  $n$ . The functor  $\max(B, T)$  yields a bag of  $n$  and is defined as follows:

(Def. 5)  $\max(B, T) \in B$  and for every bag  $x$  of  $n$  such that  $x \in B$  holds  $x \leq_T \max(B, T)$ .

Let  $O$  be an ordinal number. Observe that  $\text{InvLexOrder } O$  is connected.

## 2. MISCELLANY ON GRAPHS

Let  $G$  be a graph. Note that there exists a vertex sequence of  $G$  which is non empty and one-to-one.

Let  $G$  be a graph and let  $V$  be a non empty vertex sequence of  $G$ . A walk of  $G$  is called a walk of  $V$  if:

(Def. 6)  $\text{It.vertexSeq}() = V$ .

Let  $G$  be a graph and let  $V$  be a non empty one-to-one vertex sequence of  $G$ . One can check that every walk of  $V$  is path-like.

We now state two propositions:

- (9) For every graph  $G$  and for all walks  $W_1, W_2$  of  $G$  such that  $W_1$  is trivial and  $W_1.\text{last}() = W_2.\text{first}()$  holds  $W_1.\text{append}(W_2) = W_2$ .
- (10) Let  $G, H$  be graphs,  $A, B, C$  be sets,  $G_1$  be a subgraph of  $G$  induced by  $A$ ,  $H_1$  be a subgraph of  $H$  induced by  $B$ ,  $G_2$  be a subgraph of  $G_1$  induced by  $C$ , and  $H_2$  be a subgraph of  $H_1$  induced by  $C$ . Suppose  $G =_G H$  and  $A \subseteq B$  and  $C \subseteq A$  and  $C$  is a non empty subset of the vertices of  $G$ . Then  $G_2 =_G H_2$ .

Let  $G$  be a v-graph. We say that  $G$  is natural v-labeled if and only if:

(Def. 7) The vlabel of  $G$  is natural-yielding.

## 3. GRAPHS WITH TWO VERTEX LABELS

The natural number  $\text{V2-LabelSelector}$  is defined by:

(Def. 8)  $\text{V2-LabelSelector} = 8$ .

Let  $G$  be a graph structure. We say that  $G$  is v2-labeled if and only if:

(Def. 9)  $\text{V2-LabelSelector} \in \text{dom } G$  and there exists a function  $f$  such that  $G(\text{V2-LabelSelector}) = f$  and  $\text{dom } f \subseteq \text{the vertices of } G$ .

Let us note that there exists a graph structure which is graph-like, weighted, elabeled, vlabeled, and v2-labeled.

A v2-graph is a v2-labeled graph. A vv-graph is a vlabeled v2-labeled graph.

Let  $G$  be a v2-graph. The v2-label of  $G$  yields a function and is defined as follows:

(Def. 10) The v2-label of  $G = G(\text{V2-LabelSelector})$ .

Next we state the proposition

(11) For every v2-graph  $G$  holds  $\text{dom}(\text{the v2-label of } G) \subseteq \text{the vertices of } G$ .

Let  $G$  be a graph and let  $X$  be a set. Note that  $G.\text{set}(\text{V2-LabelSelector}, X)$  is graph-like.

We now state the proposition

(12) For every graph  $G$  and for every set  $X$  holds

$$G.\text{set}(\text{V2-LabelSelector}, X) =_G G.$$

Let  $G$  be a finite graph and let  $X$  be a set.

Note that  $G.\text{set}(\text{V2-LabelSelector}, X)$  is finite.

Let  $G$  be a loopless graph and let  $X$  be a set.

Observe that  $G.\text{set}(\text{V2-LabelSelector}, X)$  is loopless.

Let  $G$  be a trivial graph and let  $X$  be a set.

Note that  $G.\text{set}(\text{V2-LabelSelector}, X)$  is trivial.

Let  $G$  be a non trivial graph and let  $X$  be a set. One can check that  $G.\text{set}(\text{V2-LabelSelector}, X)$  is non trivial.

Let  $G$  be a non-multi graph and let  $X$  be a set. One can check that  $G.\text{set}(\text{V2-LabelSelector}, X)$  is non-multi.

Let  $G$  be a non-directed-multi graph and let  $X$  be a set. One can verify that  $G.\text{set}(\text{V2-LabelSelector}, X)$  is non-directed-multi.

Let  $G$  be a connected graph and let  $X$  be a set.

Note that  $G.\text{set}(\text{V2-LabelSelector}, X)$  is connected.

Let  $G$  be an acyclic graph and let  $X$  be a set.

One can verify that  $G.\text{set}(\text{V2-LabelSelector}, X)$  is acyclic.

Let  $G$  be a v-graph and let  $X$  be a set.

One can check that  $G.\text{set}(\text{V2-LabelSelector}, X)$  is v-labeled.

Let  $G$  be a e-graph and let  $X$  be a set. Observe that  $G.\text{set}(\text{V2-LabelSelector}, X)$  is e-labeled.

Let  $G$  be a w-graph and let  $X$  be a set. Observe that  $G.\text{set}(\text{V2-LabelSelector}, X)$  is w-labeled.

Let  $G$  be a v2-graph and let  $X$  be a set.

One can verify that  $G.\text{set}(\text{VLabelSelector}, X)$  is v2-labeled.

Let  $G$  be a graph, let  $Y$  be a set, and let  $X$  be a partial function from the vertices of  $G$  to  $Y$ . Observe that  $G.\text{set}(\text{V2-LabelSelector}, X)$  is v2-labeled.

Let  $G$  be a graph and let  $X$  be a many sorted set indexed by the vertices of  $G$ . Observe that  $G.\text{set}(\text{V2-LabelSelector}, X)$  is v2-labeled.

Let  $G$  be a graph. One can verify that  $G.\text{set}(\text{V2-LabelSelector}, \emptyset)$  is v2-labeled.

Let  $G$  be a v2-graph. We say that  $G$  is natural v2-labeled if and only if:

(Def. 11) The v2-label of  $G$  is natural-yielding.

We say that  $G$  is finite v2-labeled if and only if:

(Def. 12) The  $v_2$ -label of  $G$  is finite-yielding.

We say that  $G$  is natsubset  $v_2$ -labeled if and only if:

(Def. 13) The  $v_2$ -label of  $G$  is natsubset yielding.

One can check that there exists a weighted elabeled vlabeled  $v_2$ -labeled graph which is finite, natural  $v$ -labeled, finite  $v_2$ -labeled, natsubset  $v_2$ -labeled, and chordal and there exists a weighted elabeled vlabeled  $v_2$ -labeled graph which is finite, natural  $v$ -labeled, natural  $v_2$ -labeled, and chordal.

Let  $G$  be a natural  $v$ -labeled  $v$ -graph. Observe that the  $v$ label of  $G$  is natural-yielding.

Let  $G$  be a natural  $v_2$ -labeled  $v_2$ -graph. Observe that the  $v_2$ -label of  $G$  is natural-yielding.

Let  $G$  be a finite  $v_2$ -labeled  $v_2$ -graph. Observe that the  $v_2$ -label of  $G$  is finite-yielding.

Let  $G$  be a natsubset  $v_2$ -labeled  $v_2$ -graph. One can verify that the  $v_2$ -label of  $G$  is natsubset yielding.

Let  $G$  be a  $vv$ -graph and let  $v, x$  be sets. One can check that  $G.\text{labelVertex}(v, x)$  is  $v_2$ -labeled.

Next we state the proposition

(13) For every  $vv$ -graph  $G$  and for all sets  $v, x$  holds the  $v_2$ -label of  $G =$  the  $v_2$ -label of  $G.\text{labelVertex}(v, x)$ .

Let  $G$  be a natural  $v$ -labeled  $vv$ -graph, let  $v$  be a set, and let  $x$  be a natural number. Observe that  $G.\text{labelVertex}(v, x)$  is natural  $v$ -labeled.

Let  $G$  be a natural  $v_2$ -labeled  $vv$ -graph, let  $v$  be a set, and let  $x$  be a natural number. Observe that  $G.\text{labelVertex}(v, x)$  is natural  $v_2$ -labeled.

Let  $G$  be a finite  $v_2$ -labeled  $vv$ -graph, let  $v$  be a set, and let  $x$  be a natural number. Note that  $G.\text{labelVertex}(v, x)$  is finite  $v_2$ -labeled.

Let  $G$  be a natsubset  $v_2$ -labeled  $vv$ -graph, let  $v$  be a set, and let  $x$  be a natural number. One can check that  $G.\text{labelVertex}(v, x)$  is natsubset  $v_2$ -labeled.

Let  $G$  be a graph. Note that there exists a subgraph of  $G$  which is vlabeled and  $v_2$ -labeled.

Let  $G$  be a  $v_2$ -graph and let  $G_2$  be a  $v_2$ -labeled subgraph of  $G$ . We say that  $G_2$  inherits  $v_2$ -label if and only if:

(Def. 14) The  $v_2$ -label of  $G_2 =$  (the  $v_2$ -label of  $G$ )|(the vertices of  $G_2$ ).

Let  $G$  be a  $v_2$ -graph. Note that there exists a  $v_2$ -labeled subgraph of  $G$  which inherits  $v_2$ -label.

Let  $G$  be a  $v_2$ -graph. A  $v_2$ -subgraph of  $G$  is a  $v_2$ -labeled subgraph of  $G$  inheriting  $v_2$ -label.

Let  $G$  be a  $vv$ -graph. Note that there exists a vlabeled  $v_2$ -labeled subgraph of  $G$  which inherits  $v$ label and  $v_2$ -label.

Let  $G$  be a  $vv$ -graph. A  $vv$ -subgraph of  $G$  is a vlabeled  $v_2$ -labeled subgraph of  $G$  inheriting  $v$ label and  $v_2$ -label.

Let  $G$  be a natural  $v$ -labeled  $v$ -graph. Note that every  $v$ -subgraph of  $G$  is natural  $v$ -labeled.

Let  $G$  be a graph and let  $V, E$  be sets. Observe that there exists a subgraph of  $G$  induced by  $V$  and  $E$  which is weighted, elabeled, vlabeled, and  $v2$ -labeled.

Let  $G$  be a  $vv$ -graph and let  $V, E$  be sets. Observe that there exists a vlabeled  $v2$ -labeled subgraph of  $G$  induced by  $V$  and  $E$  which inherits vlabeled and  $v2$ -label.

Let  $G$  be a  $vv$ -graph and let  $V, E$  be sets. A  $(V, E)$ -induced  $vv$ -subgraph of  $G$  is a vlabeled  $v2$ -labeled subgraph of  $G$  induced by  $V$  and  $E$  inheriting vlabeled and  $v2$ -label.

Let  $G$  be a  $vv$ -graph and let  $V$  be a set. A  $V$ -induced  $vv$ -subgraph of  $G$  is a  $(V, G.edgesBetween(V))$ -induced  $vv$ -subgraph of  $G$ .

#### 4. MORE ON GRAPH SEQUENCES

Let  $s$  be a many sorted set indexed by  $\mathbb{N}$ . We say that  $s$  is iterative if and only if:

(Def. 15) For all natural numbers  $k, n$  such that  $s(k) = s(n)$  holds  $s(k + 1) = s(n + 1)$ .

Let  $G_3$  be a many sorted set indexed by  $\mathbb{N}$ . We say that  $G_3$  is eventually constant if and only if:

(Def. 16) There exists a natural number  $n$  such that for every natural number  $m$  such that  $n \leq m$  holds  $G_3(n) = G_3(m)$ .

Let us observe that there exists a many sorted set indexed by  $\mathbb{N}$  which is halting, iterative, and eventually constant.

The following proposition is true

(14) For every many sorted set  $G_4$  indexed by  $\mathbb{N}$  such that  $G_4$  is halting and iterative holds  $G_4$  is eventually constant.

One can check that every many sorted set indexed by  $\mathbb{N}$  which is halting and iterative is also eventually constant.

The following proposition is true

(15) For every many sorted set  $G_4$  indexed by  $\mathbb{N}$  such that  $G_4$  is eventually constant holds  $G_4$  is halting.

Let us mention that every many sorted set indexed by  $\mathbb{N}$  which is eventually constant is also halting.

One can prove the following two propositions:

(16) Let  $G_4$  be an iterative eventually constant many sorted set indexed by  $\mathbb{N}$  and  $n$  be a natural number. If  $G_4.Lifespan() \leq n$ , then  $G_4(G_4.Lifespan()) = G_4(n)$ .

(17) Let  $G_4$  be an iterative eventually constant many sorted set indexed by  $\mathbb{N}$  and  $n, m$  be natural numbers. If  $G_4.\text{Lifespan}() \leq n$  and  $n \leq m$ , then  $G_4(m) = G_4(n)$ .

Let  $G_3$  be a v-graph sequence. We say that  $G_3$  is natural v-labeled if and only if:

(Def. 17) For every natural number  $x$  holds  $G_3(x)$  is natural v-labeled.

Let  $G_3$  be a graph sequence. We say that  $G_3$  is chordal if and only if:

(Def. 18) For every natural number  $x$  holds  $G_3(x)$  is chordal.

We say that  $G_3$  has fixed vertices if and only if:

(Def. 19) For all natural numbers  $n, m$  holds the vertices of  $G_3(n) =$  the vertices of  $G_3(m)$ .

We say that  $G_3$  is v2-labeled if and only if:

(Def. 20) For every natural number  $x$  holds  $G_3(x)$  is v2-labeled.

Let us observe that there exists a graph sequence which is weighted, elabeled, vlabeled, and v2-labeled.

A v2-graph sequence is a v2-labeled graph sequence. A vv-graph sequence is a vlabeled v2-labeled graph sequence.

Let  $G_5$  be a v2-graph sequence and let  $x$  be a natural number. Note that  $G_5(x)$  is v2-labeled.

Let  $G_5$  be a v2-graph sequence. We say that  $G_5$  is natural v2-labeled if and only if:

(Def. 21) For every natural number  $x$  holds  $G_5(x)$  is natural v2-labeled.

We say that  $G_5$  is finite v2-labeled if and only if:

(Def. 22) For every natural number  $x$  holds  $G_5(x)$  is finite v2-labeled.

We say that  $G_5$  is natsubset v2-labeled if and only if:

(Def. 23) For every natural number  $x$  holds  $G_5(x)$  is natsubset v2-labeled.

Let us mention that there exists a weighted elabeled vlabeled v2-labeled graph sequence which is finite, natural v-labeled, finite v2-labeled, natsubset v2-labeled, and chordal and there exists a weighted elabeled vlabeled v2-labeled graph sequence which is finite, natural v-labeled, natural v2-labeled, and chordal.

Let  $G_4$  be a v-graph sequence and let  $x$  be a natural number. Then  $G_4(x)$  is a v-graph.

Let  $G_5$  be a natural v-labeled v-graph sequence and let  $x$  be a natural number. Observe that  $G_5(x)$  is natural v-labeled.

Let  $G_5$  be a natural v2-labeled v2-graph sequence and let  $x$  be a natural number. One can check that  $G_5(x)$  is natural v2-labeled.

Let  $G_5$  be a finite v2-labeled v2-graph sequence and let  $x$  be a natural number. One can verify that  $G_5(x)$  is finite v2-labeled.

Let  $G_5$  be a natsubset v2-labeled v2-graph sequence and let  $x$  be a natural number. Note that  $G_5(x)$  is natsubset v2-labeled.

Let  $G_5$  be a chordal graph sequence and let  $x$  be a natural number. One can check that  $G_5(x)$  is chordal.

Let  $G_4$  be a v-graph sequence and let  $n$  be a natural number. Then  $G_4(n)$  is a v-graph.

Let  $G_4$  be a finite v-graph sequence and let  $n$  be a natural number. One can check that  $G_4(n)$  is finite.

Let  $G_4$  be a vv-graph sequence and let  $n$  be a natural number. Then  $G_4(n)$  is a vv-graph.

Let  $G_4$  be a finite vv-graph sequence and let  $n$  be a natural number. One can verify that  $G_4(n)$  is finite.

Let  $G_4$  be a chordal vv-graph sequence and let  $n$  be a natural number. Note that  $G_4(n)$  is chordal.

Let  $G_4$  be a natural v-labeled vv-graph sequence and let  $n$  be a natural number. One can check that  $G_4(n)$  is natural v-labeled.

Let  $G_4$  be a finite v2-labeled vv-graph sequence and let  $n$  be a natural number. Note that  $G_4(n)$  is finite v2-labeled.

Let  $G_4$  be a natsubset v2-labeled vv-graph sequence and let  $n$  be a natural number. One can check that  $G_4(n)$  is natsubset v2-labeled.

Let  $G_4$  be a natural v2-labeled vv-graph sequence and let  $n$  be a natural number. Observe that  $G_4(n)$  is natural v2-labeled.

## 5. VERTICES NUMBERING SEQUENCES

Let  $G_3$  be a v-graph sequence. We say that  $G_3$  has initially empty v-label if and only if:

(Def. 24) The vlabel of  $G_3(0) = \emptyset$ .

We say that  $G_3$  is adding one at a step if and only if the condition (Def. 25) is satisfied.

(Def. 25) Let  $n$  be a natural number. Suppose  $n < G_3.\text{Lifespan}()$ . Then there exists a set  $w$  such that  $w \notin \text{dom}(\text{the vlabel of } G_3(n))$  and the vlabel of  $G_3(n+1) = (\text{the vlabel of } G_3(n)) + \cdot (w \mapsto (G_3.\text{Lifespan}() -' n))$ .

Let  $G_3$  be a v-graph sequence. We say that  $G_3$  is v-label numbering if and only if the condition (Def. 26) is satisfied.

(Def. 26)  $G_3$  is iterative, eventually constant, finite, natural v-labeled, and adding one at a step and has fixed vertices and initially empty v-label.

One can check that there exists a v-graph sequence which is iterative, eventually constant, finite, natural v-labeled, and adding one at a step and has fixed vertices and initially empty v-label.



Let us observe that there exists a v-graph sequence which is v-label numbering.

One can check the following observations:

- \* every v-graph sequence which is v-label numbering is also iterative,
- \* every v-graph sequence which is v-label numbering is also eventually constant,
- \* every v-graph sequence which is v-label numbering is also finite,
- \* every v-graph sequence which is v-label numbering has also fixed vertices,
- \* every v-graph sequence which is v-label numbering is also natural v-labeled,
- \* every v-graph sequence which is v-label numbering has also initially empty v-label, and
- \* every v-graph sequence which is v-label numbering is also adding one at a step.

A v-label numbering sequence is a v-label numbering v-graph sequence.

Let  $G_3$  be a v-label numbering sequence and let  $n$  be a natural number. The functor  $G_3.PickedAt\ n$  yields a set and is defined by:

- (Def. 27)(i)  $G_3.PickedAt\ n = choose(\text{the vertices of } G_3(0))$  if  $n \geq G_3.Lifespan()$ ,  
 (ii)  $G_3.PickedAt\ n \notin \text{dom}(\text{the vlabel of } G_3(n))$  and the vlabel of  $G_3(n + 1) = (\text{the vlabel of } G_3(n)) + ((G_3.PickedAt\ n) \mapsto (G_3.Lifespan() -' n))$ , otherwise.

The following propositions are true:

- (18) Let  $G_3$  be a v-label numbering sequence and  $n$  be a natural number. If  $n < G_3.Lifespan()$ , then  $G_3.PickedAt\ n \in G_3(n + 1).labeledV()$  and  $G_3(n + 1).labeledV() = G_3(n).labeledV() \cup \{G_3.PickedAt\ n\}$ .
- (19) Let  $G_3$  be a v-label numbering sequence and  $n$  be a natural number. If  $n < G_3.Lifespan()$ , then  $(\text{the vlabel of } G_3(n + 1))(G_3.PickedAt\ n) = G_3.Lifespan() -' n$ .
- (20) For every v-label numbering sequence  $G_3$  and for every natural number  $n$  such that  $n \leq G_3.Lifespan()$  holds  $card(G_3(n).labeledV()) = n$ .
- (21) For every v-label numbering sequence  $G_3$  and for every natural number  $n$  holds  $rng(\text{the vlabel of } G_3(n)) = Seg(G_3.Lifespan()) \setminus Seg(G_3.Lifespan() -' n)$ .
- (22) Let  $G_3$  be a v-label numbering sequence,  $n$  be a natural number, and  $x$  be a set. Then  $(\text{the vlabel of } G_3(n))(x) \leq G_3.Lifespan()$  and if  $x \in G_3(n).labeledV()$ , then  $1 \leq (\text{the vlabel of } G_3(n))(x)$ .
- (23) Let  $G_3$  be a v-label numbering sequence and  $n, m$  be natural numbers. Suppose  $G_3.Lifespan() -' n < m$  and  $m \leq G_3.Lifespan()$ . Then there exists a vertex  $v$  of  $G_3(n)$  such that  $v \in G_3(n).labeledV()$  and  $(\text{the vlabel of } G_3(n))(v) = m$ .

of  $G_3(n)(v) = m$ .

- (24) Let  $G_3$  be a v-label numbering sequence and  $m, n$  be natural numbers. If  $m \leq n$ , then the vlabel of  $G_3(m) \subseteq$  the vlabel of  $G_3(n)$ .
- (25) For every v-label numbering sequence  $G_3$  and for every natural number  $n$  holds the vlabel of  $G_3(n)$  is one-to-one.
- (26) Let  $G_3$  be a v-label numbering sequence,  $m, n$  be natural numbers, and  $v$  be a set. Suppose  $v \in G_3(m).\text{labeledV}()$  and  $v \in G_3(n).\text{labeledV}()$ . Then (the vlabel of  $G_3(m))(v) =$  (the vlabel of  $G_3(n))(v)$ .
- (27) Let  $G_3$  be a v-label numbering sequence,  $v$  be a set, and  $m, n$  be natural numbers. If  $v \in G_3(m).\text{labeledV}()$  and (the vlabel of  $G_3(m))(v) = n$ , then  $G_3.\text{PickedAt}(G_3.\text{Lifespan}() -' n) = v$ .
- (28) Let  $G_3$  be a v-label numbering sequence and  $m, n$  be natural numbers. If  $n < G_3.\text{Lifespan}()$  and  $n < m$ , then  $G_3.\text{PickedAt } n \in G_3(m).\text{labeledV}()$  and (the vlabel of  $G_3(m))(G_3.\text{PickedAt } n) = G_3.\text{Lifespan}() -' n$ .
- (29) Let  $G_3$  be a v-label numbering sequence,  $m$  be a natural number, and  $v$  be a set. Suppose  $v \in G_3(m).\text{labeledV}()$ . Then  $G_3.\text{Lifespan}() -'$  (the vlabel of  $G_3(m))(v) < m$  and  $G_3.\text{Lifespan}() -' m <$  (the vlabel of  $G_3(m))(v)$ .
- (30) Let  $G_3$  be a v-label numbering sequence,  $i$  be a natural number, and  $a, b$  be sets. Suppose  $a \in G_3(i).\text{labeledV}()$  and  $b \in G_3(i).\text{labeledV}()$  and (the vlabel of  $G_3(i))(a) <$  (the vlabel of  $G_3(i))(b)$ . Then  $b \in G_3(G_3.\text{Lifespan}() -' (\text{the vlabel of } G_3(i))(a)).\text{labeledV}()$ .
- (31) Let  $G_3$  be a v-label numbering sequence,  $i$  be a natural number, and  $a, b$  be sets. Suppose  $a \in G_3(i).\text{labeledV}()$  and  $b \in G_3(i).\text{labeledV}()$  and (the vlabel of  $G_3(i))(a) <$  (the vlabel of  $G_3(i))(b)$ . Then  $a \notin G_3(G_3.\text{Lifespan}() -' (\text{the vlabel of } G_3(i))(b)).\text{labeledV}()$ .

## 6. LEXICOGRAPHICAL BREADTH-FIRST SEARCH

Let  $G$  be a graph. The functor  $\text{LexBFS:Init } G$  yields a natural v-labeled finite v2-labeled natsubset v2-labeled vv-graph and is defined as follows:

- (Def. 28)  $\text{LexBFS:Init } G = G.\text{set}(\text{VLabelSelector}, \emptyset).\text{set}(\text{V2-LabelSelector}, (\text{the vertices of } G) \mapsto \emptyset)$ .

Let  $G$  be a finite graph. Then  $\text{LexBFS:Init } G$  is a finite natural v-labeled finite v2-labeled natsubset v2-labeled vv-graph.

Let  $G$  be a finite finite v2-labeled natsubset v2-labeled vv-graph. Let us assume that  $\text{dom}(\text{the v2-label of } G) = \text{the vertices of } G$ . The functor  $\text{LexBFS:PickUnnumbered } G$  yields a vertex of  $G$  and is defined by:

- (Def. 29)(i)  $\text{LexBFS:PickUnnumbered } G = \text{choose}(\text{the vertices of } G)$  if  $\text{dom}(\text{the vlabel of } G) = \text{the vertices of } G$ ,

- (ii) there exists a non empty finite subset  $S$  of  $2^{\mathbb{N}}$  and there exists a non empty finite subset  $B$  of Bags  $\mathbb{N}$  and there exists a function  $F$  such that  $S = \text{rng } F$  and  $F = (\text{the v2-label of } G) \upharpoonright ((\text{the vertices of } G) \setminus \text{dom}(\text{the vlabel of } G))$  and for every finite subset  $x$  of  $\mathbb{N}$  such that  $x \in S$  holds  $(x, 1)\text{-bag} \in B$  and for every set  $x$  such that  $x \in B$  there exists a finite subset  $y$  of  $\mathbb{N}$  such that  $y \in S$  and  $x = (y, 1)\text{-bag}$  and  $\text{LexBFS:PickUnnumbered } G = \text{choose}(F^{-1}(\{\text{support max}(B, \text{InvLexOrder } \mathbb{N})\}))$ , otherwise.

Let  $G$  be a vv-graph, let  $v$  be a set, and let  $k$  be a natural number. The functor  $\text{LexBFS:LabelAdjacent}(G, v, k)$  yielding a vv-graph is defined as follows:

(Def. 30)  $\text{LexBFS:LabelAdjacent}(G, v, k) = G.\text{set}(\text{V2-LabelSelector}, (\text{the v2-label of } G) \upharpoonright ((G.\text{adjacentSet}(\{v\}) \setminus \text{dom}(\text{the vlabel of } G)) \mapsto \{k\}))$ .

Next we state four propositions:

- (32) Let  $G$  be a vv-graph,  $v, x$  be sets, and  $k$  be a natural number. If  $x \notin G.\text{adjacentSet}(\{v\})$ , then  $(\text{the v2-label of } G)(x) = (\text{the v2-label of } \text{LexBFS:LabelAdjacent}(G, v, k))(x)$ .
- (33) Let  $G$  be a vv-graph,  $v, x$  be sets, and  $k$  be a natural number. Suppose  $x \in \text{dom}(\text{the vlabel of } G)$ . Then  $(\text{the v2-label of } G)(x) = (\text{the v2-label of } \text{LexBFS:LabelAdjacent}(G, v, k))(x)$ .
- (34) Let  $G$  be a vv-graph,  $v, x$  be sets, and  $k$  be a natural number. Suppose  $x \in G.\text{adjacentSet}(\{v\})$  and  $x \notin \text{dom}(\text{the vlabel of } G)$ . Then  $(\text{the v2-label of } \text{LexBFS:LabelAdjacent}(G, v, k))(x) = (\text{the v2-label of } G)(x) \cup \{k\}$ .
- (35) Let  $G$  be a vv-graph,  $v$  be a set, and  $k$  be a natural number. Suppose  $\text{dom}(\text{the v2-label of } G) = \text{the vertices of } G$ . Then  $\text{dom}(\text{the v2-label of } \text{LexBFS:LabelAdjacent}(G, v, k)) = \text{the vertices of } G$ .

Let  $G$  be a finite natural v-labeled finite v2-labeled natsubset v2-labeled vv-graph, let  $v$  be a vertex of  $G$ , and let  $k$  be a natural number. Then  $\text{LexBFS:LabelAdjacent}(G, v, k)$  is a finite natural v-labeled finite v2-labeled natsubset v2-labeled vv-graph.

Let  $G$  be a finite natural v-labeled finite v2-labeled natsubset v2-labeled vv-graph, let  $v$  be a vertex of  $G$ , and let  $n$  be a natural number. The functor  $\text{LexBFS:Update}(G, v, n)$  yielding a finite natural v-labeled finite v2-labeled natsubset v2-labeled vv-graph is defined by:

(Def. 31)  $\text{LexBFS:Update}(G, v, n) = \text{LexBFS:LabelAdjacent}(G.\text{labelVertex}(v, G.\text{order}() - 'n), v, G.\text{order}() - 'n)$ .

Let  $G$  be a finite natural v-labeled finite v2-labeled natsubset v2-labeled vv-graph. The functor  $\text{LexBFS:Step } G$  yields a finite natural v-labeled finite v2-labeled natsubset v2-labeled vv-graph and is defined as follows:

(Def. 32)  $\text{LexBFS:Step } G = \begin{cases} G, & \text{if } G.\text{order}() \leq \text{card dom}(\text{the vlabel of } G), \\ \text{LexBFS:Update}(G, \text{LexBFS:PickUnnumbered } G, & \\ \quad \text{card dom}(\text{the vlabel of } G)), & \text{otherwise.} \end{cases}$

Let  $G$  be a finite graph. The functor  $\text{LexBFS:CSeq } G$  yields a finite natural  $v$ -labeled finite  $v2$ -labeled natsubset  $v2$ -labeled  $vv$ -graph sequence and is defined by:

- (Def. 33)  $(\text{LexBFS:CSeq } G)(0) = \text{LexBFS:Init } G$  and for every natural number  $n$  holds  $(\text{LexBFS:CSeq } G)(n+1) = \text{LexBFS:Step}(\text{LexBFS:CSeq } G)(n)$ .

We now state the proposition

- (36) For every finite graph  $G$  holds  $\text{LexBFS:CSeq } G$  is iterative.

Let  $G$  be a finite graph. Observe that  $\text{LexBFS:CSeq } G$  is iterative.

Next we state a number of propositions:

- (37) For every graph  $G$  holds the  $v$ label of  $\text{LexBFS:Init } G = \emptyset$ .

- (38) Let  $G$  be a graph and  $v$  be a set. Then  $\text{dom}(\text{the } v2\text{-label of } \text{LexBFS:Init } G) = \text{the vertices of } G$  and  $(\text{the } v2\text{-label of } \text{LexBFS:Init } G)(v) = \emptyset$ .

- (39) For every graph  $G$  holds  $G =_G \text{LexBFS:Init } G$ .

- (40) Let  $G$  be a finite finite  $v2$ -labeled natsubset  $v2$ -labeled  $vv$ -graph and  $x$  be a set. Suppose that

- (i)  $x \notin \text{dom}(\text{the } v\text{label of } G)$ ,
- (ii)  $\text{dom}(\text{the } v2\text{-label of } G) = \text{the vertices of } G$ , and
- (iii)  $\text{dom}(\text{the } v\text{label of } G) \neq \text{the vertices of } G$ .

Then  $((\text{the } v2\text{-label of } G)(x), 1)\text{-bag} \leq_{\text{InvLexOrder } \mathbb{N}} ((\text{the } v2\text{-label of } G)(\text{LexBFS:PickUnnumbered } G), 1)\text{-bag}$ .

- (41) Let  $G$  be a finite finite  $v2$ -labeled natsubset  $v2$ -labeled  $vv$ -graph. Suppose  $\text{dom}(\text{the } v2\text{-label of } G) = \text{the vertices of } G$  and  $\text{dom}(\text{the } v\text{label of } G) \neq \text{the vertices of } G$ . Then  $\text{LexBFS:PickUnnumbered } G \notin \text{dom}(\text{the } v\text{label of } G)$ .

- (42) For every finite graph  $G$  and for every natural number  $n$  holds  $(\text{LexBFS:CSeq } G)(n) =_G G$ .

- (43) For every finite graph  $G$  and for all natural numbers  $m, n$  holds  $(\text{LexBFS:CSeq } G)(m) =_G (\text{LexBFS:CSeq } G)(n)$ .

- (44) Let  $G$  be a finite graph and  $n$  be a natural number. Suppose  $\text{card } \text{dom}(\text{the } v\text{label of } (\text{LexBFS:CSeq } G)(n)) < G.\text{order}()$ . Then the  $v$ label of  $(\text{LexBFS:CSeq } G)(n+1) = (\text{the } v\text{label of } (\text{LexBFS:CSeq } G)(n) + (\text{LexBFS:PickUnnumbered}(\text{LexBFS:CSeq } G)(n) \mapsto (G.\text{order}() - \text{card } \text{dom}(\text{the } v\text{label of } (\text{LexBFS:CSeq } G)(n))))$ .

- (45) For every finite graph  $G$  and for every natural number  $n$  holds  $\text{dom}(\text{the } v2\text{-label of } (\text{LexBFS:CSeq } G)(n)) = \text{the vertices of } (\text{LexBFS:CSeq } G)(n)$ .

- (46) For every finite graph  $G$  and for every natural number  $n$  such that  $n \leq G.\text{order}()$  holds  $\text{card } \text{dom}(\text{the } v\text{label of } (\text{LexBFS:CSeq } G)(n)) = n$ .

- (47) For every finite graph  $G$  and for every natural number  $n$  such that  $G.\text{order}() \leq n$  holds  $(\text{LexBFS:CSeq } G)(G.\text{order}()) =$

- (LexBFS:CSeq  $G$ )( $n$ ).
- (48) For every finite graph  $G$  and for all natural numbers  $m, n$  such that  $G.order() \leq m$  and  $m \leq n$  holds  $(LexBFS:CSeq G)(m) = (LexBFS:CSeq G)(n)$ .
- (49) For every finite graph  $G$  holds LexBFS:CSeq  $G$  is eventually constant.
- Let  $G$  be a finite graph. Note that LexBFS:CSeq  $G$  is eventually constant. We now state two propositions:
- (50) Let  $G$  be a finite graph and  $n$  be a natural number. Then  $dom$  (the vlabel of  $(LexBFS:CSeq G)(n)$ ) = the vertices of  $(LexBFS:CSeq G)(n)$  and only if  $G.order() \leq n$ .
- (51) For every finite graph  $G$  holds  $(LexBFS:CSeq G).Lifespan() = G.order()$ .

Let  $G$  be a finite chordal graph and let  $i$  be a natural number. One can check that  $(LexBFS:CSeq G)(i)$  is chordal.

Let  $G$  be a finite chordal graph. One can check that LexBFS:CSeq  $G$  is chordal.

One can prove the following proposition

- (52) For every finite graph  $G$  holds LexBFS:CSeq  $G$  is v-label numbering.
- Let  $G$  be a finite graph. Note that LexBFS:CSeq  $G$  is v-label numbering. We now state several propositions:
- (53) For every finite graph  $G$  and for every natural number  $n$  such that  $n < G.order()$  holds  $LexBFS:CSeq G.PickedAt n = LexBFS:PickUnnumbered(LexBFS:CSeq G)(n)$ .
- (54) Let  $G$  be a finite graph and  $n$  be a natural number. Suppose  $n < G.order()$ . Then there exists a vertex  $w$  of  $(LexBFS:CSeq G)(n)$  such that
- (i)  $w = LexBFS:PickUnnumbered(LexBFS:CSeq G)(n)$ , and
  - (ii) for every set  $v$  holds if  $v \in G.adjacentSet(\{w\})$  and  $v \notin dom$  (the vlabel of  $(LexBFS:CSeq G)(n)$ ), then (the v2-label of  $(LexBFS:CSeq G)(n + 1)(v) = (the v2-label of (LexBFS:CSeq G)(n)(v) \cup \{G.order() - 'n\}$  and if  $v \notin G.adjacentSet(\{w\})$  or  $v \in dom$  (the vlabel of  $(LexBFS:CSeq G)(n)$ ), then (the v2-label of  $(LexBFS:CSeq G)(n + 1)(v) = (the v2-label of (LexBFS:CSeq G)(n)(v)$ .

- (55) Let  $G$  be a finite graph,  $i$  be a natural number, and  $v$  be a set. Then (the v2-label of  $(LexBFS:CSeq G)(i)(v) \subseteq Seg(G.order()) \setminus Seg(G.order() - 'i)$ .
- (56) Let  $G$  be a finite graph,  $x$  be a set, and  $i, j$  be natural numbers. If  $i \leq j$ , then (the v2-label of  $(LexBFS:CSeq G)(i)(x) \subseteq (the v2-label of (LexBFS:CSeq G)(j)(x)$ .
- (57) Let  $G$  be a finite graph,  $m, n$  be natural numbers, and  $x, y$  be sets. Suppose  $n < G.order()$  and  $n < m$  and  $y = LexBFS:PickUnnumbered(LexBFS:CSeq G)(n)$  and  $x \notin dom$  (the vlabel of

$(\text{LexBFS:CSeq } G)(n))$  and  $x \in G.\text{adjacentSet}(\{y\})$ . Then  $G.\text{order}() -' n \in$  (the v2-label of  $(\text{LexBFS:CSeq } G)(m))(x)$ .

- (58) Let  $G$  be a finite graph and  $m, n$  be natural numbers. Suppose  $m < n$ . Let  $x$  be a set. Suppose  $G.\text{order}() -' m \notin$  (the v2-label of  $(\text{LexBFS:CSeq } G)(m+1))(x)$ . Then  $G.\text{order}() -' m \notin$  (the v2-label of  $(\text{LexBFS:CSeq } G)(n))(x)$ .
- (59) Let  $G$  be a finite graph and  $m, n, k$  be natural numbers. Suppose  $k < n$  and  $n \leq m$ . Let  $x$  be a set. Suppose  $G.\text{order}() -' k \notin$  (the v2-label of  $(\text{LexBFS:CSeq } G)(n))(x)$ . Then  $G.\text{order}() -' k \notin$  (the v2-label of  $(\text{LexBFS:CSeq } G)(m))(x)$ .
- (60) Let  $G$  be a finite graph,  $m, n$  be natural numbers, and  $x$  be a vertex of  $(\text{LexBFS:CSeq } G)(m)$ . Suppose  $n \in$  (the v2-label of  $(\text{LexBFS:CSeq } G)(m))(x)$ . Then there exists a vertex  $y$  of  $(\text{LexBFS:CSeq } G)(m)$  such that  $\text{LexBFS:PickUnnumbered}(\text{LexBFS:CSeq } G)(G.\text{order}() -' n) = y$  and  $y \notin \text{dom}$  (the vlabel of  $(\text{LexBFS:CSeq } G)(G.\text{order}() -' n)$ ) and  $x \in G.\text{adjacentSet}(\{y\})$ .

Let  $G_4$  be a finite natural v-labeled vv-graph sequence. Then  $G_4.\text{Result}()$  is a finite natural v-labeled vv-graph.

The following four propositions are true:

- (61) For every finite graph  $G$  holds  $(\text{LexBFS:CSeq } G).\text{Result}().\text{labeledV}() =$  the vertices of  $G$ .
- (62) For every finite graph  $G$  holds (the vlabel of  $(\text{LexBFS:CSeq } G).\text{Result}()^{-1}$ ) is a vertex scheme of  $G$ .
- (63) Let  $G$  be a finite graph,  $i, j$  be natural numbers, and  $a, b$  be vertices of  $(\text{LexBFS:CSeq } G)(i)$ . Suppose that
- (i)  $a \in \text{dom}$  (the vlabel of  $(\text{LexBFS:CSeq } G)(i)$ ),
  - (ii)  $b \in \text{dom}$  (the vlabel of  $(\text{LexBFS:CSeq } G)(i)$ ),
  - (iii) (the vlabel of  $(\text{LexBFS:CSeq } G)(i))(a) <$  (the vlabel of  $(\text{LexBFS:CSeq } G)(i))(b)$ , and
  - (iv)  $j = G.\text{order}() -'$  (the vlabel of  $(\text{LexBFS:CSeq } G)(i))(b)$ .
- Then  $((\text{the v2-label of } (\text{LexBFS:CSeq } G)(j))(a), 1)\text{-bag} \leq_{\text{InvLexOrder } \mathbb{N}}$   $((\text{the v2-label of } (\text{LexBFS:CSeq } G)(j))(b), 1)\text{-bag}$ .
- (64) Let  $G$  be a finite graph,  $i, j$  be natural numbers, and  $v$  be a vertex of  $(\text{LexBFS:CSeq } G)(i)$ . Suppose  $j \in$  (the v2-label of  $(\text{LexBFS:CSeq } G)(i))(v)$ . Then there exists a vertex  $w$  of  $(\text{LexBFS:CSeq } G)(i)$  such that  $w \in \text{dom}$  (the vlabel of  $(\text{LexBFS:CSeq } G)(i)$ ) and (the vlabel of  $(\text{LexBFS:CSeq } G)(i))(w) = j$  and  $v \in G.\text{adjacentSet}(\{w\})$ .

Let  $G$  be a natural v-labeled v-graph. We say that  $G$  has property  $L3$  if and only if the condition (Def. 34) is satisfied.

(Def. 34) Let  $a, b, c$  be vertices of  $G$ . Suppose that  $a \in \text{dom}(\text{the vlabel of } G)$  and  $b \in \text{dom}(\text{the vlabel of } G)$  and  $c \in \text{dom}(\text{the vlabel of } G)$  and  $(\text{the vlabel of } G)(a) < (\text{the vlabel of } G)(b)$  and  $(\text{the vlabel of } G)(b) < (\text{the vlabel of } G)(c)$  and  $a$  and  $c$  are adjacent and  $b$  and  $c$  are not adjacent. Then there exists a vertex  $d$  of  $G$  such that

- (i)  $d \in \text{dom}(\text{the vlabel of } G)$ ,
- (ii)  $(\text{the vlabel of } G)(c) < (\text{the vlabel of } G)(d)$ ,
- (iii)  $b$  and  $d$  are adjacent,
- (iv)  $a$  and  $d$  are not adjacent, and
- (v) for every vertex  $e$  of  $G$  such that  $e \neq d$  and  $e$  and  $b$  are adjacent and  $e$  and  $a$  are not adjacent holds  $(\text{the vlabel of } G)(e) < (\text{the vlabel of } G)(d)$ .

One can prove the following three propositions:

- (65) For every finite graph  $G$  and for every natural number  $n$  holds  $(\text{LexBFS:CSeq } G)(n)$  has property  $L3$ .
- (66) Let  $G$  be a finite chordal natural v-labeled v-graph. Suppose  $G$  has property  $L3$  and  $\text{dom}(\text{the vlabel of } G) = \text{the vertices of } G$ . Let  $V$  be a vertex scheme of  $G$ . If  $V^{-1} = \text{the vlabel of } G$ , then  $V$  is perfect.
- (67) For every finite chordal vv-graph  $G$  holds  $(\text{the vlabel of } (\text{LexBFS:CSeq } G).\text{Result}())^{-1}$  is a perfect vertex scheme of  $G$ .

## 7. THE MAXIMUM CARDINALITY SEARCH ALGORITHM

Let  $G$  be a finite graph. The functor  $\text{MCS:Init } G$  yields a finite natural v-labeled natural v2-labeled vv-graph and is defined by:

(Def. 35)  $\text{MCS:Init } G = G.\text{set}(\text{VLabelSelector}, \emptyset).\text{set}(\text{V2-LabelSelector}, (\text{the vertices of } G) \mapsto 0)$ .

Let  $G$  be a finite natural v2-labeled vv-graph. Let us assume that  $\text{dom}(\text{the v2-label of } G) = \text{the vertices of } G$ . The functor  $\text{MCS:PickUnnumbered } G$  yields a vertex of  $G$  and is defined by:

- (Def. 36)(i)  $\text{MCS:PickUnnumbered } G = \text{choose}(\text{the vertices of } G)$  if  $\text{dom}(\text{the vlabel of } G) = \text{the vertices of } G$ ,
- (ii) there exists a finite non empty natural-membered set  $S$  and there exists a function  $F$  such that  $S = \text{rng } F$  and  $F = (\text{the v2-label of } G) \upharpoonright ((\text{the vertices of } G) \setminus \text{dom}(\text{the vlabel of } G))$  and  $\text{MCS:PickUnnumbered } G = \text{choose}(F^{-1}(\{\max S\}))$ , otherwise.

Let  $G$  be a finite natural v2-labeled vv-graph and let  $v$  be a set. The functor  $\text{MCS:LabelAdjacent}(G, v)$  yields a finite natural v2-labeled vv-graph and is defined by:

(Def. 37)  $\text{MCS:LabelAdjacent}(G, v) = G.\text{set}(\text{V2-LabelSelector}, (\text{the v2-label of } G).\text{incSubset}((G.\text{adjacentSet}(\{v\}) \setminus \text{dom}(\text{the vlabel of } G), 1))$ .

Let  $G$  be a finite natural  $v$ -labeled natural  $v2$ -labeled  $vv$ -graph and let  $v$  be a vertex of  $G$ . Then  $\text{MCS:LabelAdjacent}(G, v)$  is a finite natural  $v$ -labeled natural  $v2$ -labeled  $vv$ -graph.

Let  $G$  be a finite natural  $v$ -labeled natural  $v2$ -labeled  $vv$ -graph, let  $v$  be a vertex of  $G$ , and let  $n$  be a natural number. The functor  $\text{MCS:Update}(G, v, n)$  yielding a finite natural  $v$ -labeled natural  $v2$ -labeled  $vv$ -graph is defined as follows:

(Def. 38)  $\text{MCS:Update}(G, v, n) = \text{MCS:LabelAdjacent}(G.\text{labelVertex}(v, G.\text{order}() - n), v)$ .

Let  $G$  be a finite natural  $v$ -labeled natural  $v2$ -labeled  $vv$ -graph. The functor  $\text{MCS:Step } G$  yielding a finite natural  $v$ -labeled natural  $v2$ -labeled  $vv$ -graph is defined by:

(Def. 39)  $\text{MCS:Step } G = \begin{cases} G, & \text{if } G.\text{order}() \leq \text{card dom}(\text{the vlabel of } G), \\ \text{MCS:Update}(G, \text{MCS:PickUnnumbered } G, \text{card dom}(\text{the vlabel of } G)), & \text{otherwise.} \end{cases}$

Let  $G$  be a finite graph. The functor  $\text{MCS:CSeq } G$  yields a finite natural  $v$ -labeled natural  $v2$ -labeled  $vv$ -graph sequence and is defined by:

(Def. 40)  $(\text{MCS:CSeq } G)(0) = \text{MCS:Init } G$  and for every natural number  $n$  holds  $(\text{MCS:CSeq } G)(n + 1) = \text{MCS:Step}(\text{MCS:CSeq } G)(n)$ .

The following proposition is true

(68) For every finite graph  $G$  holds  $\text{MCS:CSeq } G$  is iterative.

Let  $G$  be a finite graph. Observe that  $\text{MCS:CSeq } G$  is iterative.

We now state a number of propositions:

(69) For every finite graph  $G$  holds the  $v$ label of  $\text{MCS:Init } G = \emptyset$ .

(70) Let  $G$  be a finite graph and  $v$  be a set. Then  $\text{dom}(\text{the } v2\text{-label of } \text{MCS:Init } G) = \text{the vertices of } G$  and  $(\text{the } v2\text{-label of } \text{MCS:Init } G)(v) = 0$ .

(71) For every finite graph  $G$  holds  $G =_G \text{MCS:Init } G$ .

(72) Let  $G$  be a finite natural  $v2$ -labeled  $vv$ -graph and  $x$  be a set. Suppose that

- (i)  $x \notin \text{dom}(\text{the vlabel of } G)$ ,
- (ii)  $\text{dom}(\text{the } v2\text{-label of } G) = \text{the vertices of } G$ , and
- (iii)  $\text{dom}(\text{the vlabel of } G) \neq \text{the vertices of } G$ .

Then  $(\text{the } v2\text{-label of } G)(x) \leq (\text{the } v2\text{-label of } G)(\text{MCS:PickUnnumbered } G)$ .

(73) Let  $G$  be a finite natural  $v2$ -labeled  $vv$ -graph. Suppose  $\text{dom}(\text{the } v2\text{-label of } G) = \text{the vertices of } G$  and  $\text{dom}(\text{the vlabel of } G) \neq \text{the vertices of } G$ . Then  $\text{MCS:PickUnnumbered } G \notin \text{dom}(\text{the vlabel of } G)$ .

(74) Let  $G$  be a finite natural  $v2$ -labeled  $vv$ -graph and  $v, x$  be sets. If  $x \notin G.\text{adjacentSet}(\{v\})$ , then  $(\text{the } v2\text{-label of } G)(x) = (\text{the } v2\text{-label of } G)(\text{MCS:PickUnnumbered } G)$ .



$\text{MCS:LabelAdjacent}(G, v)(x)$ .

- (75) Let  $G$  be a finite natural v2-labeled vv-graph and  $v, x$  be sets. Suppose  $x \in \text{dom}(\text{the vlabel of } G)$ . Then  $(\text{the v2-label of } G)(x) = (\text{the v2-label of } \text{MCS:LabelAdjacent}(G, v))(x)$ .
- (76) Let  $G$  be a finite natural v2-labeled vv-graph and  $v, x$  be sets. Suppose  $x \in \text{dom}(\text{the v2-label of } G)$  and  $x \in G.\text{adjacentSet}(\{v\})$  and  $x \notin \text{dom}(\text{the vlabel of } G)$ . Then  $(\text{the v2-label of } \text{MCS:LabelAdjacent}(G, v))(x) = (\text{the v2-label of } G)(x) + 1$ .
- (77) Let  $G$  be a finite natural v2-labeled vv-graph and  $v$  be a set. Suppose  $\text{dom}(\text{the v2-label of } G) = \text{the vertices of } G$ . Then  $\text{dom}(\text{the v2-label of } \text{MCS:LabelAdjacent}(G, v)) = \text{the vertices of } G$ .
- (78) For every finite graph  $G$  and for every natural number  $n$  holds  $(\text{MCS:CSeq } G)(n) =_G G$ .
- (79) For every finite graph  $G$  and for all natural numbers  $m, n$  holds  $(\text{MCS:CSeq } G)(m) =_G (\text{MCS:CSeq } G)(n)$ .

Let  $G$  be a finite chordal graph and let  $n$  be a natural number. Observe that  $(\text{MCS:CSeq } G)(n)$  is chordal.

Let  $G$  be a finite chordal graph. Observe that  $\text{MCS:CSeq } G$  is chordal.

One can prove the following propositions:

- (80) For every finite graph  $G$  and for every natural number  $n$  holds  $\text{dom}(\text{the v2-label of } (\text{MCS:CSeq } G)(n)) = \text{the vertices of } (\text{MCS:CSeq } G)(n)$ .
- (81) Let  $G$  be a finite graph and  $n$  be a natural number. Suppose  $\text{card } \text{dom}(\text{the vlabel of } (\text{MCS:CSeq } G)(n)) < G.\text{order}()$ . Then the vlabel of  $(\text{MCS:CSeq } G)(n + 1) = (\text{the vlabel of } (\text{MCS:CSeq } G)(n) + (\text{MCS:PickUnnumbered}(\text{MCS:CSeq } G)(n) \mapsto (G.\text{order}() - \text{card } \text{dom}(\text{the vlabel of } (\text{MCS:CSeq } G)(n))))$ .
- (82) For every finite graph  $G$  and for every natural number  $n$  such that  $n \leq G.\text{order}()$  holds  $\text{card } \text{dom}(\text{the vlabel of } (\text{MCS:CSeq } G)(n)) = n$ .
- (83) For every finite graph  $G$  and for every natural number  $n$  such that  $G.\text{order}() \leq n$  holds  $(\text{MCS:CSeq } G)(G.\text{order}()) = (\text{MCS:CSeq } G)(n)$ .
- (84) For every finite graph  $G$  and for all natural numbers  $m, n$  such that  $G.\text{order}() \leq m$  and  $m \leq n$  holds  $(\text{MCS:CSeq } G)(m) = (\text{MCS:CSeq } G)(n)$ .
- (85) For every finite graph  $G$  holds  $\text{MCS:CSeq } G$  is eventually constant.

Let  $G$  be a finite graph. Observe that  $\text{MCS:CSeq } G$  is eventually constant.

The following propositions are true:

- (86) Let  $G$  be a finite graph and  $n$  be a natural number. Then  $\text{dom}(\text{the vlabel of } (\text{MCS:CSeq } G)(n)) = \text{the vertices of } (\text{MCS:CSeq } G)(n)$  if and only if  $G.\text{order}() \leq n$ .
- (87) For every finite graph  $G$  holds  $(\text{MCS:CSeq } G).\text{Lifespan}() = G.\text{order}()$ .

- (88) For every finite graph  $G$  holds  $\text{MCS:CSeq } G$  is v-label numbering.  
 Let  $G$  be a finite graph. Note that  $\text{MCS:CSeq } G$  is v-label numbering.  
 Next we state three propositions:
- (89) For every finite graph  $G$  and for every natural number  $n$  such that  $n < G.\text{order}()$  holds  $\text{MCS:CSeq } G.\text{PickedAt } n = \text{MCS:PickUnnumbered}(\text{MCS:CSeq } G)(n)$ .
- (90) Let  $G$  be a finite graph and  $n$  be a natural number. Suppose  $n < G.\text{order}()$ . Then there exists a vertex  $w$  of  $(\text{MCS:CSeq } G)(n)$  such that
- (i)  $w = \text{MCS:PickUnnumbered}(\text{MCS:CSeq } G)(n)$ , and
  - (ii) for every set  $v$  holds if  $v \in G.\text{adjacentSet}(\{w\})$  and  $v \notin \text{dom}(\text{the vlabel of } (\text{MCS:CSeq } G)(n))$ , then  $(\text{the v2-label of } (\text{MCS:CSeq } G)(n+1))(v) = (\text{the v2-label of } (\text{MCS:CSeq } G)(n))(v) + 1$  and if  $v \notin G.\text{adjacentSet}(\{w\})$  or  $v \in \text{dom}(\text{the vlabel of } (\text{MCS:CSeq } G)(n))$ , then  $(\text{the v2-label of } (\text{MCS:CSeq } G)(n+1))(v) = (\text{the v2-label of } (\text{MCS:CSeq } G)(n))(v)$ .
- (91) Let  $G$  be a finite graph,  $n$  be a natural number, and  $x$  be a set. Suppose  $x \notin \text{dom}(\text{the vlabel of } (\text{MCS:CSeq } G)(n))$ . Then  $(\text{the v2-label of } (\text{MCS:CSeq } G)(n))(x) = \text{card}((G.\text{adjacentSet}(\{x\})) \cap \text{dom}(\text{the vlabel of } (\text{MCS:CSeq } G)(n)))$ .

Let  $G$  be a natural v-labeled v-graph. We say that  $G$  has property  $T$  if and only if the condition (Def. 41) is satisfied.

- (Def. 41) Let  $a, b, c$  be vertices of  $G$ . Suppose that  $a \in \text{dom}(\text{the vlabel of } G)$  and  $b \in \text{dom}(\text{the vlabel of } G)$  and  $c \in \text{dom}(\text{the vlabel of } G)$  and  $(\text{the vlabel of } G)(a) < (\text{the vlabel of } G)(b)$  and  $(\text{the vlabel of } G)(b) < (\text{the vlabel of } G)(c)$  and  $a$  and  $c$  are adjacent and  $b$  and  $c$  are not adjacent. Then there exists a vertex  $d$  of  $G$  such that
- (i)  $d \in \text{dom}(\text{the vlabel of } G)$ ,
  - (ii)  $(\text{the vlabel of } G)(b) < (\text{the vlabel of } G)(d)$ ,
  - (iii)  $b$  and  $d$  are adjacent, and
  - (iv)  $a$  and  $d$  are not adjacent.

We now state three propositions:

- (92) For every finite graph  $G$  and for every natural number  $n$  holds  $(\text{MCS:CSeq } G)(n)$  has property  $T$ .
- (93) For every finite graph  $G$  holds  $(\text{LexBFS:CSeq } G).\text{Result}()$  has property  $T$ .
- (94) Let  $G$  be a finite chordal natural v-labeled v-graph. Suppose  $G$  has property  $T$  and  $\text{dom}(\text{the vlabel of } G) = \text{the vertices of } G$ . Let  $V$  be a vertex scheme of  $G$ . If  $V^{-1} = \text{the vlabel of } G$ , then  $V$  is perfect.

## REFERENCES

- [1] Broderick Arneson and Piotr Rudnicki. Chordal graphs. *Formalized Mathematics*, 14(3):79–92, 2006.
- [2] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(2):377–382, 1990.
- [3] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [4] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [5] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [6] Czesław Byliński. A classical first order language. *Formalized Mathematics*, 1(4):669–676, 1990.
- [7] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [8] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [9] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Formalized Mathematics*, 1(3):521–527, 1990.
- [10] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [11] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [12] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(1):165–167, 1990.
- [13] M. Ch. Golumbic. *Algorithmic Graph Theory and Perfect Graphs*. Academic Press, New York, 1980.
- [14] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Formalized Mathematics*, 1(3):471–475, 1990.
- [15] Gilbert Lee. Walks in Graphs. *Formalized Mathematics*, 13(2):253–269, 2005.
- [16] Gilbert Lee. Weighted and Labeled Graphs. *Formalized Mathematics*, 13(2):279–293, 2005.
- [17] Gilbert Lee and Piotr Rudnicki. On ordering of bags. *Formalized Mathematics*, 10(1):39–46, 2002.
- [18] Gilbert Lee and Piotr Rudnicki. Alternative graph structures. *Formalized Mathematics*, 13(2):235–252, 2005.
- [19] Yatsuka Nakamura, Piotr Rudnicki, Andrzej Trybulec, and Pauline N. Kawamoto. Preliminaries to circuits, I. *Formalized Mathematics*, 5(2):167–172, 1996.
- [20] Takaya Nishiyama and Yasuho Mizuhara. Binary arithmetics. *Formalized Mathematics*, 4(1):83–86, 1993.
- [21] Piotr Rudnicki. Little Bezout theorem (factor theorem). *Formalized Mathematics*, 12(1):49–58, 2004.
- [22] Piotr Rudnicki and Andrzej Trybulec. Abian’s fixed point theorem. *Formalized Mathematics*, 6(3):335–338, 1997.
- [23] Piotr Rudnicki and Andrzej Trybulec. Multivariate polynomials with arbitrary number of variables. *Formalized Mathematics*, 9(1):95–110, 2001.
- [24] Christoph Schwarzweller. Term orders. *Formalized Mathematics*, 11(1):105–111, 2003.
- [25] R. E. Tarjan and M. Yannakakis. Simple linear-time algorithms to test chordality of graphs, test acyclicity of hypergraphs, and selectively reduce acyclic hypergraphs. *SIAM J. Comput.*, 13(3):566–579, 1984.
- [26] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics*.
- [27] Andrzej Trybulec. Domains and their Cartesian products. *Formalized Mathematics*, 1(1):115–122, 1990.
- [28] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [29] Andrzej Trybulec. Many-sorted sets. *Formalized Mathematics*, 4(1):15–22, 1993.
- [30] Andrzej Trybulec. On the sets inhabited by numbers. *Formalized Mathematics*, 11(4):341–347, 2003.
- [31] Wojciech A. Trybulec. Pigeon hole principle. *Formalized Mathematics*, 1(3):575–579, 1990.
- [32] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [33] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.

- [34] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.
- [35] Edmund Woronowicz and Anna Zalewska. Properties of binary relations. *Formalized Mathematics*, 1(1):85–89, 1990.

*Received November 17, 2006*

---