

# The Rank+Nullity Theorem

Jesse Alama  
Department of Philosophy  
Stanford University  
USA

**Summary.** The rank+nullity theorem states that, if  $T$  is a linear transformation from a finite-dimensional vector space  $V$  to a finite-dimensional vector space  $W$ , then  $\dim(V) = \text{rank}(T) + \text{nullity}(T)$ , where  $\text{rank}(T) = \dim(\text{im}(T))$  and  $\text{nullity}(T) = \dim(\text{ker}(T))$ . The proof treated here is standard; see, for example, [14]: take a basis  $A$  of  $\text{ker}(T)$  and extend it to a basis  $B$  of  $V$ , and then show that  $\dim(\text{im}(T))$  is equal to  $|B - A|$ , and that  $T$  is one-to-one on  $B - A$ .

MML identifier: RANKNULL, version: 7.8.05 4.87.985

The articles [21], [11], [32], [22], [19], [33], [34], [7], [2], [17], [10], [18], [8], [9], [20], [1], [12], [3], [5], [6], [27], [29], [24], [31], [25], [13], [4], [30], [28], [26], [23], [15], [16], and [35] provide the notation and terminology for this paper.

## 1. PRELIMINARIES

One can prove the following three propositions:

- (1) For all functions  $f, g$  such that  $g$  is one-to-one and  $f \upharpoonright \text{rng } g$  is one-to-one and  $\text{rng } g \subseteq \text{dom } f$  holds  $f \cdot g$  is one-to-one.
- (2) For every function  $f$  and for all sets  $X, Y$  such that  $X \subseteq Y$  and  $f \upharpoonright Y$  is one-to-one holds  $f \upharpoonright X$  is one-to-one.
- (3) Let  $V$  be a 1-sorted structure and  $X, Y$  be subsets of  $V$ . Then  $X$  meets  $Y$  if and only if there exists an element  $v$  of  $V$  such that  $v \in X$  and  $v \in Y$ .

In the sequel  $F$  is a field and  $V, W$  are vector spaces over  $F$ .

Let  $F$  be a field and let  $V$  be a finite dimensional vector space over  $F$ . One can verify that there exists a basis of  $V$  which is finite.

Let  $F$  be a field and let  $V, W$  be vector spaces over  $F$ . Note that there exists a function from  $V$  into  $W$  which is linear.

Next we state three propositions:

- (4) If  $\Omega_V$  is finite, then  $V$  is finite dimensional.
- (5) For every finite dimensional vector space  $V$  over  $F$  such that  $\overline{\overline{\Omega_V}} = 1$  holds  $\dim(V) = 0$ .
- (6) If  $\overline{\overline{\Omega_V}} = 2$ , then  $\dim(V) = 1$ .

## 2. BASIC FACTS OF LINEAR TRANSFORMATIONS

Let  $F$  be a field and let  $V, W$  be vector spaces over  $F$ . A linear transformation from  $V$  to  $W$  is a linear function from  $V$  into  $W$ .

In the sequel  $T$  is a linear transformation from  $V$  to  $W$ .

One can prove the following propositions:

- (7) For all non empty 1-sorted structures  $V, W$  and for every function  $T$  from  $V$  into  $W$  holds  $\text{dom } T = \Omega_V$  and  $\text{rng } T \subseteq \Omega_W$ .
- (8) For all elements  $x, y$  of  $V$  holds  $T(x) - T(y) = T(x - y)$ .
- (9)  $T(0_V) = 0_W$ .

Let  $F$  be a field, let  $V, W$  be vector spaces over  $F$ , and let  $T$  be a linear transformation from  $V$  to  $W$ . The functor  $\ker T$  yielding a strict subspace of  $V$  is defined as follows:

(Def. 1)  $\Omega_{\ker T} = \{u; u \text{ ranges over elements of } V: T(u) = 0_W\}$ .

We now state the proposition

- (10) For every element  $x$  of  $V$  holds  $x \in \ker T$  iff  $T(x) = 0_W$ .

Let  $V, W$  be non empty 1-sorted structures, let  $T$  be a function from  $V$  into  $W$ , and let  $X$  be a subset of  $V$ . Then  $T^\circ X$  is a subset of  $W$ .

Let  $F$  be a field, let  $V, W$  be vector spaces over  $F$ , and let  $T$  be a linear transformation from  $V$  to  $W$ . The functor  $\text{im } T$  yielding a strict subspace of  $W$  is defined as follows:

(Def. 2)  $\Omega_{\text{im } T} = T^\circ(\Omega_V)$ .

The following propositions are true:

- (11)  $0_V \in \ker T$ .
- (12) For every subset  $X$  of  $V$  holds  $T^\circ X$  is a subset of  $\text{im } T$ .
- (13) For every element  $y$  of  $W$  holds  $y \in \text{im } T$  iff there exists an element  $x$  of  $V$  such that  $y = T(x)$ .
- (14) For every element  $x$  of  $\ker T$  holds  $T(x) = 0_W$ .
- (15) If  $T$  is one-to-one, then  $\ker T = \mathbf{0}_V$ .
- (16) For every finite dimensional vector space  $V$  over  $F$  holds  $\dim(\mathbf{0}_V) = 0$ .

- (17) For all elements  $x, y$  of  $V$  such that  $T(x) = T(y)$  holds  $x - y \in \ker T$ .
- (18) For every subset  $A$  of  $V$  and for all elements  $x, y$  of  $V$  such that  $x - y \in \text{Lin}(A)$  holds  $x \in \text{Lin}(A \cup \{y\})$ .

3. SOME LEMMAS ON LINEARLY INDEPENDENT SUBSETS, LINEAR COMBINATIONS, AND LINEAR TRANSFORMATIONS

One can prove the following propositions:

- (19) For every subset  $X$  of  $V$  such that  $V$  is a subspace of  $W$  holds  $X$  is a subset of  $W$ .
- (20) For every subset  $A$  of  $V$  such that  $A$  is linearly independent holds  $A$  is a basis of  $\text{Lin}(A)$ .
- (21) For every subset  $A$  of  $V$  and for every element  $x$  of  $V$  such that  $x \in \text{Lin}(A)$  and  $x \notin A$  holds  $A \cup \{x\}$  is linearly dependent.
- (22) For every subset  $A$  of  $V$  and for every basis  $B$  of  $V$  such that  $A$  is a basis of  $\ker T$  and  $A \subseteq B$  holds  $T \upharpoonright (B \setminus A)$  is one-to-one.
- (23) Let  $A$  be a subset of  $V$ ,  $l$  be a linear combination of  $A$ ,  $x$  be an element of  $V$ , and  $a$  be an element of  $F$ . Then  $l + \cdot (x, a)$  is a linear combination of  $A \cup \{x\}$ .

Let  $V$  be a 1-sorted structure and let  $X$  be a subset of  $V$ . The functor  $V \setminus X$  yields a subset of  $V$  and is defined by:

(Def. 3)  $V \setminus X = \Omega_V \setminus X$ .

Let  $F$  be a field, let  $V$  be a vector space over  $F$ , let  $l$  be a linear combination of  $V$ , and let  $X$  be a subset of  $V$ . Then  $l^\circ X$  is a subset of  $F$ .

In the sequel  $l$  is a linear combination of  $V$ .

Let  $F$  be a field and let  $V$  be a vector space over  $F$ . Note that there exists a subset of  $V$  which is linearly dependent.

Let  $F$  be a field, let  $V$  be a vector space over  $F$ , let  $l$  be a linear combination of  $V$ , and let  $A$  be a subset of  $V$ . The functor  $l[A]$  yields a linear combination of  $A$  and is defined by:

(Def. 4)  $l[A] = l \upharpoonright A + \cdot (V \setminus A \mapsto 0_F)$ .

The following propositions are true:

- (24)  $l = l[\text{the support of } l]$ .
- (25) For every subset  $A$  of  $V$  and for every element  $v$  of  $V$  such that  $v \in A$  holds  $l[A](v) = l(v)$ .
- (26) For every subset  $A$  of  $V$  and for every element  $v$  of  $V$  such that  $v \notin A$  holds  $l[A](v) = 0_F$ .
- (27) For all subsets  $A, B$  of  $V$  and for every linear combination  $l$  of  $B$  such that  $A \subseteq B$  holds  $l = l[A] + l[B \setminus A]$ .

Let  $F$  be a field, let  $V$  be a vector space over  $F$ , let  $l$  be a linear combination of  $V$ , and let  $X$  be a subset of  $V$ . Observe that  $l^\circ X$  is finite.

Let  $V, W$  be non empty 1-sorted structures, let  $T$  be a function from  $V$  into  $W$ , and let  $X$  be a subset of  $W$ . Then  $T^{-1}(X)$  is a subset of  $V$ .

We now state the proposition

- (28) For every subset  $X$  of  $V$  such that  $X$  misses the support of  $l$  holds  $l^\circ X \subseteq \{0_F\}$ .

Let  $F$  be a field, let  $V, W$  be vector spaces over  $F$ , let  $l$  be a linear combination of  $V$ , and let  $T$  be a linear transformation from  $V$  to  $W$ . The functor  $T^\circ l$  yielding a linear combination of  $W$  is defined by:

(Def. 5) For every element  $w$  of  $W$  holds  $(T^\circ l)(w) = \sum(l^\circ T^{-1}(\{w\}))$ .

One can prove the following propositions:

- (29)  $T^\circ l$  is a linear combination of  $T^\circ(\text{the support of } l)$ .
- (30) The support of  $T^\circ l \subseteq T^\circ(\text{the support of } l)$ .
- (31) Let  $l, m$  be linear combinations of  $V$ . Suppose the support of  $l$  misses the support of  $m$ . Then the support of  $l + m = (\text{the support of } l) \cup (\text{the support of } m)$ .
- (32) Let  $l, m$  be linear combinations of  $V$ . Suppose the support of  $l$  misses the support of  $m$ . Then the support of  $l - m = (\text{the support of } l) \cup (\text{the support of } m)$ .
- (33) For all subsets  $A, B$  of  $V$  such that  $A \subseteq B$  and  $B$  is a basis of  $V$  holds  $V$  is the direct sum of  $\text{Lin}(A)$  and  $\text{Lin}(B \setminus A)$ .
- (34) Let  $A$  be a subset of  $V$ ,  $l$  be a linear combination of  $A$ , and  $v$  be an element of  $V$ . Suppose  $T \upharpoonright A$  is one-to-one and  $v \in A$ . Then there exists a subset  $X$  of  $V$  such that  $X$  misses  $A$  and  $T^{-1}(\{T(v)\}) = \{v\} \cup X$ .
- (35) For every subset  $X$  of  $V$  such that  $X$  misses the support of  $l$  and  $X \neq \emptyset$  holds  $l^\circ X = \{0_F\}$ .
- (36) For every element  $w$  of  $W$  such that  $w \in$  the support of  $T^\circ l$  holds  $T^{-1}(\{w\})$  meets the support of  $l$ .
- (37) Let  $v$  be an element of  $V$ . Suppose  $T \upharpoonright (\text{the support of } l)$  is one-to-one and  $v \in$  the support of  $l$ . Then  $(T^\circ l)(T(v)) = l(v)$ .
- (38) Let  $G$  be a finite sequence of elements of  $V$ . Suppose  $\text{rng } G =$  the support of  $l$  and  $T \upharpoonright (\text{the support of } l)$  is one-to-one. Then  $T \cdot (lG) = (T^\circ l)(T \cdot G)$ .
- (39) If  $T \upharpoonright (\text{the support of } l)$  is one-to-one, then  $T^\circ(\text{the support of } l) =$  the support of  $T^\circ l$ .
- (40) Let  $A$  be a subset of  $V$ ,  $B$  be a basis of  $V$ , and  $l$  be a linear combination of  $B \setminus A$ . If  $A$  is a basis of  $\ker T$  and  $A \subseteq B$ , then  $T(\sum l) = \sum(T^\circ l)$ .
- (41) Let  $X$  be a subset of  $V$ . Suppose  $X$  is linearly dependent. Then there exists a linear combination  $l$  of  $X$  such that the support of  $l \neq \emptyset$  and

$$\sum l = 0_V.$$

Let  $F$  be a field, let  $V, W$  be vector spaces over  $F$ , let  $X$  be a subset of  $V$ , let  $T$  be a linear transformation from  $V$  to  $W$ , and let  $l$  be a linear combination of  $T^\circ X$ . Let us assume that  $T \upharpoonright X$  is one-to-one. The functor  $T \# l$  yields a linear combination of  $X$  and is defined as follows:

(Def. 6)  $T \# l = l \cdot T + \cdot (V \setminus X \mapsto 0_F)$ .

We now state two propositions:

- (42) Let  $X$  be a subset of  $V$ ,  $l$  be a linear combination of  $T^\circ X$ , and  $v$  be an element of  $V$ . If  $v \in X$  and  $T \upharpoonright X$  is one-to-one, then  $(T \# l)(v) = l(T(v))$ .
- (43) For every subset  $X$  of  $V$  and for every linear combination  $l$  of  $T^\circ X$  such that  $T \upharpoonright X$  is one-to-one holds  $T^\circ T \# l = l$ .

#### 4. THE RANK+NULLITY THEOREM

Let  $F$  be a field, let  $V, W$  be finite dimensional vector spaces over  $F$ , and let  $T$  be a linear transformation from  $V$  to  $W$ . The functor  $\text{rank } T$  yielding a natural number is defined by:

(Def. 7)  $\text{rank } T = \dim(\text{im } T)$ .

The functor  $\text{nullity } T$  yields a natural number and is defined by:

(Def. 8)  $\text{nullity } T = \dim(\text{ker } T)$ .

Next we state two propositions:

- (44) Let  $V, W$  be finite dimensional vector spaces over  $F$  and  $T$  be a linear transformation from  $V$  to  $W$ . Then  $\dim(V) = \text{rank } T + \text{nullity } T$ .
- (45) Let  $V, W$  be finite dimensional vector spaces over  $F$  and  $T$  be a linear transformation from  $V$  to  $W$ . If  $T$  is one-to-one, then  $\dim(V) = \text{rank } T$ .

#### REFERENCES

- [1] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(2):377–382, 1990.
- [2] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [4] Grzegorz Bancerek and Andrzej Trybulec. Miscellaneous facts about functions. *Formalized Mathematics*, 5(4):485–492, 1996.
- [5] Czesław Byliński. Binary operations applied to finite sequences. *Formalized Mathematics*, 1(4):643–649, 1990.
- [6] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [7] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [8] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [9] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Formalized Mathematics*, 1(3):521–527, 1990.
- [10] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.

- [11] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [12] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(1):165–167, 1990.
- [13] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(2):335–342, 1990.
- [14] Serge Lang. *Algebra*. Springer, 3rd edition, 2005.
- [15] Robert Milewski. Associated matrix of linear map. *Formalized Mathematics*, 5(3):339–345, 1996.
- [16] Michał Muzalewski. Rings and modules – part II. *Formalized Mathematics*, 2(4):579–585, 1991.
- [17] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics*.
- [18] Andrzej Trybulec. Binary operations applied to functions. *Formalized Mathematics*, 1(2):329–334, 1990.
- [19] Andrzej Trybulec. Domains and their Cartesian products. *Formalized Mathematics*, 1(1):115–122, 1990.
- [20] Andrzej Trybulec. Function domains and Frænkel operator. *Formalized Mathematics*, 1(3):495–500, 1990.
- [21] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [22] Andrzej Trybulec. Tuples, projections and Cartesian products. *Formalized Mathematics*, 1(1):97–105, 1990.
- [23] Wojciech A. Trybulec. Basis of vector space. *Formalized Mathematics*, 1(5):883–885, 1990.
- [24] Wojciech A. Trybulec. Groups. *Formalized Mathematics*, 1(5):821–827, 1990.
- [25] Wojciech A. Trybulec. Linear combinations in real linear space. *Formalized Mathematics*, 1(3):581–588, 1990.
- [26] Wojciech A. Trybulec. Linear combinations in vector space. *Formalized Mathematics*, 1(5):877–882, 1990.
- [27] Wojciech A. Trybulec. Non-contiguous substrings and one-to-one finite sequences. *Formalized Mathematics*, 1(3):569–573, 1990.
- [28] Wojciech A. Trybulec. Operations on subspaces in vector space. *Formalized Mathematics*, 1(5):871–876, 1990.
- [29] Wojciech A. Trybulec. Pigeon hole principle. *Formalized Mathematics*, 1(3):575–579, 1990.
- [30] Wojciech A. Trybulec. Subspaces and cosets of subspaces in vector space. *Formalized Mathematics*, 1(5):865–870, 1990.
- [31] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(2):291–296, 1990.
- [32] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [33] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [34] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.
- [35] Mariusz Żynel. The Steinitz theorem and the dimension of a vector space. *Formalized Mathematics*, 5(3):423–428, 1996.

*Received July 31, 2007*

---