

# Basic Properties of Even and Odd Functions

Bo Li  
Qingdao University of Science  
and Technology  
China

Yanhong Men  
Qingdao University of Science  
and Technology  
China

**Summary.** In this article we present definitions, basic properties and some examples of even and odd functions [6].

MML identifier: FUNCT\_8, version: 7.11.02 4.125.1059

The articles [2], [5], [1], [8], [14], [12], [15], [7], [17], [3], [4], [11], [19], [13], [10], [18], [16], and [9] provide the notation and terminology for this paper.

## 1. EVEN AND ODD FUNCTIONS

In this paper  $x, r$  denote real numbers.

Let  $A$  be a set. We say that  $A$  is symmetrical if and only if:

(Def. 1) For every complex number  $x$  such that  $x \in A$  holds  $-x \in A$ .

One can check that there exists a subset of  $\mathbb{C}$  which is symmetrical.

Let us note that there exists a subset of  $\mathbb{R}$  which is symmetrical.

In the sequel  $A$  is a symmetrical subset of  $\mathbb{C}$ .

Let  $R$  be a binary relation. We say that  $R$  has symmetrical domain if and only if:

(Def. 2)  $\text{dom } R$  is symmetrical.

Let us observe that every binary relation which is empty has also symmetrical domain and there exists a binary relation which has symmetrical domain.

Let  $R$  be a binary relation with symmetrical domain. One can check that  $\text{dom } R$  is symmetrical.

Let  $X, Y$  be complex-membered sets and let  $F$  be a partial function from  $X$  to  $Y$ . We say that  $F$  is quasi even if and only if:

(Def. 3) For every  $x$  such that  $x, -x \in \text{dom } F$  holds  $F(-x) = F(x)$ .

Let  $X, Y$  be complex-membered sets and let  $F$  be a partial function from  $X$  to  $Y$ . We say that  $F$  is even if and only if:

(Def. 4)  $F$  is quasi even and has symmetrical domain.

Let  $X, Y$  be complex-membered sets. Note that every partial function from  $X$  to  $Y$  which is quasi even and has symmetrical domain is also even and every partial function from  $X$  to  $Y$  which is even is also quasi even and has symmetrical domain.

Let  $A$  be a set, let  $X, Y$  be complex-membered sets, and let  $F$  be a partial function from  $X$  to  $Y$ . We say that  $F$  is even on  $A$  if and only if:

(Def. 5)  $A \subseteq \text{dom } F$  and  $F \upharpoonright A$  is even.

Let  $X, Y$  be complex-membered sets and let  $F$  be a partial function from  $X$  to  $Y$ . We say that  $F$  is quasi odd if and only if:

(Def. 6) For every  $x$  such that  $x, -x \in \text{dom } F$  holds  $F(-x) = -F(x)$ .

Let  $X, Y$  be complex-membered sets and let  $F$  be a partial function from  $X$  to  $Y$ . We say that  $F$  is odd if and only if:

(Def. 7)  $F$  is quasi odd and has symmetrical domain.

Let  $X, Y$  be complex-membered sets. Note that every partial function from  $X$  to  $Y$  which is quasi odd and has symmetrical domain is also odd and every partial function from  $X$  to  $Y$  which is odd is also quasi odd and has symmetrical domain.

Let  $A$  be a set, let  $X, Y$  be complex-membered sets, and let  $F$  be a partial function from  $X$  to  $Y$ . We say that  $F$  is odd on  $A$  if and only if:

(Def. 8)  $A \subseteq \text{dom } F$  and  $F \upharpoonright A$  is odd.

In the sequel  $F, G$  denote partial functions from  $\mathbb{R}$  to  $\mathbb{R}$ .

One can prove the following propositions:

- (1)  $F$  is odd on  $A$  iff  $A \subseteq \text{dom } F$  and for every  $x$  such that  $x \in A$  holds  $F(x) + F(-x) = 0$ .
- (2)  $F$  is even on  $A$  iff  $A \subseteq \text{dom } F$  and for every  $x$  such that  $x \in A$  holds  $F(x) - F(-x) = 0$ .
- (3) If  $F$  is odd on  $A$  and for every  $x$  such that  $x \in A$  holds  $F(x) \neq 0$ , then  $A \subseteq \text{dom } F$  and for every  $x$  such that  $x \in A$  holds  $\frac{F(x)}{F(-x)} = -1$ .
- (4) If  $A \subseteq \text{dom } F$  and for every  $x$  such that  $x \in A$  holds  $\frac{F(x)}{F(-x)} = -1$ , then  $F$  is odd on  $A$ .
- (5) If  $F$  is even on  $A$  and for every  $x$  such that  $x \in A$  holds  $F(x) \neq 0$ , then  $A \subseteq \text{dom } F$  and for every  $x$  such that  $x \in A$  holds  $\frac{F(x)}{F(-x)} = 1$ .
- (6) If  $A \subseteq \text{dom } F$  and for every  $x$  such that  $x \in A$  holds  $\frac{F(x)}{F(-x)} = 1$ , then  $F$  is even on  $A$ .

- (7) If  $F$  is even on  $A$  and odd on  $A$ , then for every  $x$  such that  $x \in A$  holds  $F(x) = 0$ .
- (8) If  $F$  is even on  $A$ , then for every  $x$  such that  $x \in A$  holds  $F(x) = F(|x|)$ .
- (9) If  $A \subseteq \text{dom } F$  and for every  $x$  such that  $x \in A$  holds  $F(x) = F(|x|)$ , then  $F$  is even on  $A$ .
- (10) If  $F$  is odd on  $A$  and  $G$  is odd on  $A$ , then  $F + G$  is odd on  $A$ .
- (11) If  $F$  is even on  $A$  and  $G$  is even on  $A$ , then  $F + G$  is even on  $A$ .
- (12) If  $F$  is odd on  $A$  and  $G$  is odd on  $A$ , then  $F - G$  is odd on  $A$ .
- (13) If  $F$  is even on  $A$  and  $G$  is even on  $A$ , then  $F - G$  is even on  $A$ .
- (14) If  $F$  is odd on  $A$ , then  $rF$  is odd on  $A$ .
- (15) If  $F$  is even on  $A$ , then  $rF$  is even on  $A$ .
- (16) If  $F$  is odd on  $A$ , then  $-F$  is odd on  $A$ .
- (17) If  $F$  is even on  $A$ , then  $-F$  is even on  $A$ .
- (18) If  $F$  is odd on  $A$ , then  $F^{-1}$  is odd on  $A$ .
- (19) If  $F$  is even on  $A$ , then  $F^{-1}$  is even on  $A$ .
- (20) If  $F$  is odd on  $A$ , then  $|F|$  is even on  $A$ .
- (21) If  $F$  is even on  $A$ , then  $|F|$  is even on  $A$ .
- (22) If  $F$  is odd on  $A$  and  $G$  is odd on  $A$ , then  $FG$  is even on  $A$ .
- (23) If  $F$  is even on  $A$  and  $G$  is even on  $A$ , then  $FG$  is even on  $A$ .
- (24) If  $F$  is even on  $A$  and  $G$  is odd on  $A$ , then  $FG$  is odd on  $A$ .
- (25) If  $F$  is even on  $A$ , then  $r + F$  is even on  $A$ .
- (26) If  $F$  is even on  $A$ , then  $F - r$  is even on  $A$ .
- (27) If  $F$  is even on  $A$ , then  $F^2$  is even on  $A$ .
- (28) If  $F$  is odd on  $A$ , then  $F^2$  is even on  $A$ .
- (29) If  $F$  is odd on  $A$  and  $G$  is odd on  $A$ , then  $F/G$  is even on  $A$ .
- (30) If  $F$  is even on  $A$  and  $G$  is even on  $A$ , then  $F/G$  is even on  $A$ .
- (31) If  $F$  is odd on  $A$  and  $G$  is even on  $A$ , then  $F/G$  is odd on  $A$ .
- (32) If  $F$  is even on  $A$  and  $G$  is odd on  $A$ , then  $F/G$  is odd on  $A$ .
- (33) If  $F$  is odd, then  $-F$  is odd.
- (34) If  $F$  is even, then  $-F$  is even.
- (35) If  $F$  is odd, then  $F^{-1}$  is odd.
- (36) If  $F$  is even, then  $F^{-1}$  is even.
- (37) If  $F$  is odd, then  $|F|$  is even.
- (38) If  $F$  is even, then  $|F|$  is even.
- (39) If  $F$  is odd, then  $F^2$  is even.
- (40) If  $F$  is even, then  $F^2$  is even.
- (41) If  $F$  is even, then  $r + F$  is even.

- (42) If  $F$  is even, then  $F - r$  is even.
- (43) If  $F$  is odd, then  $r F$  is odd.
- (44) If  $F$  is even, then  $r F$  is even.
- (45) If  $F$  is odd and  $G$  is odd and  $\text{dom } F \cap \text{dom } G$  is symmetrical, then  $F + G$  is odd.
- (46) If  $F$  is even and  $G$  is even and  $\text{dom } F \cap \text{dom } G$  is symmetrical, then  $F + G$  is even.
- (47) If  $F$  is odd and  $G$  is odd and  $\text{dom } F \cap \text{dom } G$  is symmetrical, then  $F - G$  is odd.
- (48) If  $F$  is even and  $G$  is even and  $\text{dom } F \cap \text{dom } G$  is symmetrical, then  $F - G$  is even.
- (49) If  $F$  is odd and  $G$  is odd and  $\text{dom } F \cap \text{dom } G$  is symmetrical, then  $F G$  is even.
- (50) If  $F$  is even and  $G$  is even and  $\text{dom } F \cap \text{dom } G$  is symmetrical, then  $F G$  is even.
- (51) If  $F$  is even and  $G$  is odd and  $\text{dom } F \cap \text{dom } G$  is symmetrical, then  $F G$  is odd.
- (52) If  $F$  is odd and  $G$  is odd and  $\text{dom } F \cap \text{dom } G$  is symmetrical, then  $F/G$  is even.
- (53) If  $F$  is even and  $G$  is even and  $\text{dom } F \cap \text{dom } G$  is symmetrical, then  $F/G$  is even.
- (54) If  $F$  is odd and  $G$  is even and  $\text{dom } F \cap \text{dom } G$  is symmetrical, then  $F/G$  is odd.
- (55) If  $F$  is even and  $G$  is odd and  $\text{dom } F \cap \text{dom } G$  is symmetrical, then  $F/G$  is odd.

## 2. SOME EXAMPLES

The function signum from  $\mathbb{R}$  into  $\mathbb{R}$  is defined by:

(Def. 9) For every real number  $x$  holds  $\text{signum}(x) = \text{sgn } x$ .

Let  $x$  be a real number. One can verify that  $\text{signum}(x)$  is real.

Next we state a number of propositions:

- (56) For every real number  $x$  such that  $x > 0$  holds  $\text{signum}(x) = 1$ .
- (57) For every real number  $x$  such that  $x < 0$  holds  $\text{signum}(x) = -1$ .
- (58)  $\text{signum}(0) = 0$ .
- (59) For every real number  $x$  holds  $\text{signum}(-x) = -\text{signum}(x)$ .
- (60) For every symmetrical subset  $A$  of  $\mathbb{R}$  holds signum is odd on  $A$ .
- (61) For every real number  $x$  such that  $x \geq 0$  holds  $|\square|_{\mathbb{R}}(x) = x$ .

- (62) For every real number  $x$  such that  $x < 0$  holds  $|\square|_{\mathbb{R}}(x) = -x$ .
- (63) For every real number  $x$  holds  $|\square|_{\mathbb{R}}(-x) = |\square|_{\mathbb{R}}(x)$ .
- (64) For every symmetrical subset  $A$  of  $\mathbb{R}$  holds  $|\square|_{\mathbb{R}}$  is even on  $A$ .
- (65) For every symmetrical subset  $A$  of  $\mathbb{R}$  holds the function  $\sin$  is odd on  $A$ .
- (66) For every symmetrical subset  $A$  of  $\mathbb{R}$  holds the function  $\cos$  is even on  $A$ .

Let us observe that the function  $\sin$  is odd.

Let us observe that the function  $\cos$  is even.

We now state two propositions:

- (67) For every symmetrical subset  $A$  of  $\mathbb{R}$  holds the function  $\sinh$  is odd on  $A$ .
- (68) For every symmetrical subset  $A$  of  $\mathbb{R}$  holds the function  $\cosh$  is even on  $A$ .

Let us note that the function  $\sinh$  is odd.

Let us mention that the function  $\cosh$  is even.

The following propositions are true:

- (69) If  $A \subseteq ]-\frac{\pi}{2}, \frac{\pi}{2}[$ , then the function  $\tan$  is odd on  $A$ .
- (70) Suppose  $A \subseteq \text{dom}(\text{the function } \tan)$  and for every  $x$  such that  $x \in A$  holds  $(\text{the function } \cos)(x) \neq 0$ . Then the function  $\tan$  is odd on  $A$ .
- (71) Suppose  $A \subseteq \text{dom}(\text{the function } \cot)$  and for every  $x$  such that  $x \in A$  holds  $(\text{the function } \sin)(x) \neq 0$ . Then the function  $\cot$  is odd on  $A$ .
- (72) If  $A \subseteq [-1, 1]$ , then the function  $\arctan$  is odd on  $A$ .
- (73) For every symmetrical subset  $A$  of  $\mathbb{R}$  holds  $|\text{the function } \sin|$  is even on  $A$ .
- (74) For every symmetrical subset  $A$  of  $\mathbb{R}$  holds  $|\text{the function } \cos|$  is even on  $A$ .
- (75) For every symmetrical subset  $A$  of  $\mathbb{R}$  holds  $(\text{the function } \sin)^{-1}$  is odd on  $A$ .
- (76) For every symmetrical subset  $A$  of  $\mathbb{R}$  holds  $(\text{the function } \cos)^{-1}$  is even on  $A$ .
- (77) For every symmetrical subset  $A$  of  $\mathbb{R}$  holds  $-\text{the function } \sin$  is odd on  $A$ .
- (78) For every symmetrical subset  $A$  of  $\mathbb{R}$  holds  $-\text{the function } \cos$  is even on  $A$ .
- (79) For every symmetrical subset  $A$  of  $\mathbb{R}$  holds  $(\text{the function } \sin)^2$  is even on  $A$ .
- (80) For every symmetrical subset  $A$  of  $\mathbb{R}$  holds  $(\text{the function } \cos)^2$  is even on  $A$ .

In the sequel  $B$  denotes a symmetrical subset of  $\mathbb{R}$ .

One can prove the following propositions:

- (81) If  $B \subseteq \text{dom}(\text{the function sec})$ , then the function sec is even on  $B$ .
- (82) If for every real number  $x$  such that  $x \in B$  holds  $(\text{the function cos})(x) \neq 0$ , then the function sec is even on  $B$ .
- (83) If  $B \subseteq \text{dom}(\text{the function cosec})$ , then the function cosec is odd on  $B$ .
- (84) If for every real number  $x$  such that  $x \in B$  holds  $(\text{the function sin})(x) \neq 0$ , then the function cosec is odd on  $B$ .

#### REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [2] Czesław Byliński. The complex numbers. *Formalized Mathematics*, 1(3):507–513, 1990.
- [3] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [4] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [5] Pacharapokin Chanapat, Kanchun, and Hiroshi Yamazaki. Formulas and identities of trigonometric functions. *Formalized Mathematics*, 12(2):139–141, 2004.
- [6] Chuanshang Chen. *Mathematical Analysis*. Higher Education Press, Beijing, 1978.
- [7] Agata Darmochwał. The Euclidean space. *Formalized Mathematics*, 2(4):599–603, 1991.
- [8] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [9] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(2):269–272, 1990.
- [10] Xiquan Liang and Bing Xie. Inverse trigonometric functions arctan and arccot. *Formalized Mathematics*, 16(2):147–158, 2008, doi:10.2478/v10037-008-0021-3.
- [11] Takashi Mitsuishi and Yuguang Yang. Properties of the trigonometric function. *Formalized Mathematics*, 8(1):103–106, 1999.
- [12] Jan Popiołek. Some properties of functions modul and signum. *Formalized Mathematics*, 1(2):263–264, 1990.
- [13] Konrad Raczkowski and Paweł Sadowski. Topological properties of subsets in real numbers. *Formalized Mathematics*, 1(4):777–780, 1990.
- [14] Andrzej Trybulec. On the sets inhabited by numbers. *Formalized Mathematics*, 11(4):341–347, 2003.
- [15] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [16] Peng Wang and Bo Li. Several differentiation formulas of special functions. Part V. *Formalized Mathematics*, 15(3):73–79, 2007, doi:10.2478/v10037-007-0009-4.
- [17] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [18] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.
- [19] Yuguang Yang and Yasunari Shidama. Trigonometric functions and existence of circle ratio. *Formalized Mathematics*, 7(2):255–263, 1998.

Received May 25, 2009