

# Fixpoint Theorem for Continuous Functions on Chain-Complete Posets

Kazuhisa Ishida  
Neyagawa-shi  
Osaka, Japan

Yasunari Shidama  
Shinshu University  
Nagano, Japan

**Summary.** This text includes the definition of chain-complete poset, fixpoint theorem on it, and the definition of the function space of continuous functions on chain-complete posets [10].

MML identifier: POSET\_1, version: 7.11.04 4.130.1076

The papers [8], [4], [5], [3], [1], [9], [7], [11], [13], [12], [2], [14], and [6] provide the notation and terminology for this paper.

## 1. MONOTONE FUNCTIONS, CHAIN AND CHAIN-COMPLETE POSETS

Let  $P$  be a non empty poset. Observe that there exists a chain of  $P$  which is non empty.

Let  $I_1$  be a relational structure. We say that  $I_1$  is chain-complete if and only if:

- (Def. 1)  $I_1$  is lower-bounded and for every chain  $L$  of  $I_1$  such that  $L$  is non empty holds  $\sup L$  exists in  $I_1$ .

One can prove the following proposition

- (1) Let  $P_1, P_2$  be non empty posets,  $K$  be a non empty chain of  $P_1$ , and  $h$  be a monotone function from  $P_1$  into  $P_2$ . Then  $h \circ K$  is a non empty chain of  $P_2$ .

Let us note that there exists a poset which is strict, chain-complete, and non empty.

Let us mention that every relational structure which is chain-complete is also lower-bounded.

For simplicity, we adopt the following rules:  $x, y$  denote sets,  $P, Q$  denote strict chain-complete non empty posets,  $L$  denotes a non empty chain of  $P$ ,  $M$  denotes a non empty chain of  $Q$ ,  $p$  denotes an element of  $P$ ,  $f$  denotes a monotone function from  $P$  into  $Q$ , and  $g, g_1, g_2$  denote monotone functions from  $P$  into  $P$ .

We now state the proposition

$$(2) \quad \sup(f^\circ L) \leq f(\sup L).$$

## 2. FIXPOINT THEOREM FOR CONTINUOUS FUNCTIONS ON CHAIN-COMPLETE POSETS

Let  $P$  be a non empty poset, let  $g$  be a monotone function from  $P$  into  $P$ , and let  $p$  be an element of  $P$ . The functor  $\text{iterSet}(g, p)$  yields a non empty set and is defined by:

$$(\text{Def. 2}) \quad \text{iterSet}(g, p) = \{x \in P: \forall_{n: \text{natural number}} x = g^n(p)\}.$$

Next we state the proposition

$$(3) \quad \text{iterSet}(g, \perp_P) \text{ is a non empty chain of } P.$$

Let us consider  $P$  and let  $g$  be a monotone function from  $P$  into  $P$ . The functor  $\text{iter-min } g$  yields a non empty chain of  $P$  and is defined by:

$$(\text{Def. 3}) \quad \text{iter-min } g = \text{iterSet}(g, \perp_P).$$

The following propositions are true:

$$(4) \quad \sup \text{iter-min } g = \sup(g^\circ \text{iter-min } g).$$

$$(5) \quad \text{If } g_1 \leq g_2, \text{ then } \sup \text{iter-min } g_1 \leq \sup \text{iter-min } g_2.$$

Let  $P, Q$  be non empty posets and let  $f$  be a function from  $P$  into  $Q$ . We say that  $f$  is continuous if and only if:

$$(\text{Def. 4}) \quad f \text{ is monotone and for every non empty chain } L \text{ of } P \text{ holds } f \text{ preserves } \sup \text{ of } L.$$

We now state two propositions:

$$(6) \quad \text{For every function } f \text{ from } P \text{ into } Q \text{ holds } f \text{ is continuous iff } f \text{ is monotone and for every } L \text{ holds } f(\sup L) = \sup(f^\circ L).$$

$$(7) \quad \text{For every element } z \text{ of } Q \text{ holds } P \longmapsto z \text{ is continuous.}$$

Let us consider  $P, Q$ . Observe that there exists a function from  $P$  into  $Q$  which is continuous.

Let us consider  $P, Q$ . One can verify that every function from  $P$  into  $Q$  which is continuous is also monotone.

The following proposition is true

$$(8) \quad \text{For every monotone function } f \text{ from } P \text{ into } Q \text{ such that for every } L \text{ holds } f(\sup L) \leq \sup(f^\circ L) \text{ holds } f \text{ is continuous.}$$

Let us consider  $P$  and let  $g$  be a monotone function from  $P$  into  $P$ . Let us assume that  $g$  is continuous. The least fixpoint of  $g$  yields an element of  $P$  and is defined by the conditions (Def. 5).

- (Def. 5)(i) The least fixpoint of  $g$  is a fixpoint of  $g$ , and  
(ii) for every  $p$  such that  $p$  is a fixpoint of  $g$  holds the least fixpoint of  $g \leq p$ .

One can prove the following propositions:

- (9) For every continuous function  $g$  from  $P$  into  $P$  holds the least fixpoint of  $g = \text{sup iter-min } g$ .  
(10) Let  $g_1, g_2$  be continuous functions from  $P$  into  $P$ . If  $g_1 \leq g_2$ , then the least fixpoint of  $g_1 \leq$  the least fixpoint of  $g_2$ .

### 3. FUNCTION SPACE OF CONTINUOUS FUNCTIONS ON CHAIN-COMPLETE POSETS

Let us consider  $P, Q$ . The functor  $\text{ConFuncs}(P, Q)$  yields a non empty set and is defined by the condition (Def. 6).

- (Def. 6)  $\text{ConFuncs}(P, Q) = \{x; x \text{ ranges over elements of (the carrier of } Q)\text{the carrier of } P: \forall f: \text{continuous function from } P \text{ into } Q \ f = x\}$ .

Let us consider  $P, Q$ . The functor  $\text{ConRelat}(P, Q)$  yielding a binary relation on  $\text{ConFuncs}(P, Q)$  is defined by the condition (Def. 7).

- (Def. 7) Let given  $x, y$ . Then  $\langle x, y \rangle \in \text{ConRelat}(P, Q)$  if and only if the following conditions are satisfied:  
(i)  $x \in \text{ConFuncs}(P, Q)$ ,  
(ii)  $y \in \text{ConFuncs}(P, Q)$ , and  
(iii) there exist functions  $f, g$  from  $P$  into  $Q$  such that  $x = f$  and  $y = g$  and  $f \leq g$ .

Let us consider  $P, Q$ . One can verify the following observations:

- \*  $\text{ConRelat}(P, Q)$  is reflexive,
- \*  $\text{ConRelat}(P, Q)$  is transitive, and
- \*  $\text{ConRelat}(P, Q)$  is antisymmetric.

Let us consider  $P, Q$ . The functor  $\text{ConPoset}(P, Q)$  yielding a strict non empty poset is defined as follows:

- (Def. 8)  $\text{ConPoset}(P, Q) = \langle \text{ConFuncs}(P, Q), \text{ConRelat}(P, Q) \rangle$ .

In the sequel  $F$  is a non empty chain of  $\text{ConPoset}(P, Q)$ .

Let us consider  $P, Q, F, p$ . The functor  $F\text{-image}(p)$  yielding a non empty chain of  $Q$  is defined as follows:

- (Def. 9)  $F\text{-image}(p) = \{x \in Q: \forall f: \text{continuous function from } P \text{ into } Q \ (f \in F \wedge x = f(p))\}$ .

Let us consider  $P, Q, F$ . The functor  $\text{sup-func } F$  yields a function from  $P$  into  $Q$  and is defined as follows:

(Def. 10) For all  $p, M$  such that  $M = F\text{-image}(p)$  holds  $(\text{sup-func } F)(p) = \text{sup } M$ .

Let us consider  $P, Q, F$ . One can check that  $\text{sup-func } F$  is continuous.

The following proposition is true

(11)  $\text{Sup } F$  exists in  $\text{ConPoset}(P, Q)$  and  $\text{sup-func } F = \bigsqcup_{\text{ConPoset}(P, Q)} F$ .

Let us consider  $P, Q$ . The functor  $\text{min-func}(P, Q)$  yielding a function from  $P$  into  $Q$  is defined as follows:

(Def. 11)  $\text{min-func}(P, Q) = P \mapsto \perp_Q$ .

Let us consider  $P, Q$ . One can check that  $\text{min-func}(P, Q)$  is continuous.

The following proposition is true

(12) For every element  $f$  of  $\text{ConPoset}(P, Q)$  such that  $f = \text{min-func}(P, Q)$  holds  $f \leq$  the carrier of  $\text{ConPoset}(P, Q)$ .

Let us consider  $P, Q$ . Note that  $\text{ConPoset}(P, Q)$  is chain-complete.

#### 4. CONTINUITY OF FIXPOINT FUNCTION FROM $\text{CONPOSET}(P, P)$ INTO $P$

Let us consider  $P$ . The functor  $\text{fix-func } P$  yielding a function from  $\text{ConPoset}(P, P)$  into  $P$  is defined by the condition (Def. 12).

(Def. 12) Let  $g$  be an element of  $\text{ConPoset}(P, P)$  and  $h$  be a continuous function from  $P$  into  $P$ . If  $g = h$ , then  $(\text{fix-func } P)(g) =$  the least fixpoint of  $h$ .

Let us consider  $P$ . One can check that  $\text{fix-func } P$  is continuous.

#### REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [2] Grzegorz Bancerek. Bounds in posets and relational substructures. *Formalized Mathematics*, 6(1):81–91, 1997.
- [3] Grzegorz Bancerek and Andrzej Trybulec. Miscellaneous facts about functions. *Formalized Mathematics*, 5(4):485–492, 1996.
- [4] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [5] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [6] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [7] Adam Grabowski. On the category of posets. *Formalized Mathematics*, 5(4):501–505, 1996.
- [8] Piotr Rudnicki and Andrzej Trybulec. Abian’s fixed point theorem. *Formalized Mathematics*, 6(3):335–338, 1997.
- [9] Wojciech A. Trybulec and Grzegorz Bancerek. Kuratowski – Zorn lemma. *Formalized Mathematics*, 1(2):387–393, 1990.
- [10] Glynn Winskel. *The Formal Semantics of Programming Languages*. The MIT Press, 1993.
- [11] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [12] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.

- [13] Edmund Woronowicz and Anna Zalewska. Properties of binary relations. *Formalized Mathematics*, 1(1):85–89, 1990.
- [14] Mariusz Żynel and Czesław Byliński. Properties of relational structures, posets, lattices and maps. *Formalized Mathematics*, 6(1):123–130, 1997.

*Received November 10, 2009*

---