

More on the Continuity of Real Functions¹

Keiko Narita
Hirosaki-city
Aomori, Japan

Artur Kornilowicz
Institute of Informatics
University of Białystok
Sosnowa 64, 15-887 Białystok, Poland

Yasunari Shidama
Shinshu University
Nagano, Japan

Summary. In this article we demonstrate basic properties of the continuous functions from \mathbb{R} to \mathcal{R}^n which correspond to state space equations in control engineering.

MML identifier: NFCONT_4, version: 7.11.07 4.160.1126

The terminology and notation used here have been introduced in the following articles: [3], [7], [17], [2], [4], [12], [13], [14], [16], [1], [5], [9], [15], [18], [10], [8], [20], [21], [19], [11], [22], and [6].

For simplicity, we use the following convention: n, i denote elements of \mathbb{N} , X, X_1 denote sets, r, p, s, x_0, x_1, x_2 denote real numbers, f, f_1, f_2 denote partial functions from \mathbb{R} to \mathcal{R}^n , and h denotes a partial function from \mathbb{R} to the carrier of $\langle \mathcal{E}^n, \|\cdot\| \rangle$.

Let us consider n, f, x_0 . We say that f is continuous in x_0 if and only if:

(Def. 1) There exists a partial function g from \mathbb{R} to the carrier of $\langle \mathcal{E}^n, \|\cdot\| \rangle$ such that $f = g$ and g is continuous in x_0 .

We now state four propositions:

- (1) If $h = f$, then f is continuous in x_0 iff h is continuous in x_0 .
- (2) If $x_0 \in X$ and f is continuous in x_0 , then $f|X$ is continuous in x_0 .

¹This work was supported by JSPS KAKENHI 22300285.

- (3) f is continuous in x_0 if and only if the following conditions are satisfied:
- (i) $x_0 \in \text{dom } f$, and
 - (ii) for every r such that $0 < r$ there exists s such that $0 < s$ and for every x_1 such that $x_1 \in \text{dom } f$ and $|x_1 - x_0| < s$ holds $|f_{x_1} - f_{x_0}| < r$.
- (4) Let r be a real number, z be an element of \mathcal{R}^n , and w be a point of $\langle \mathcal{E}^n, \|\cdot\| \rangle$. Suppose $z = w$. Then $\{y \in \mathcal{R}^n: |y - z| < r\} = \{y; y \text{ ranges over points of } \langle \mathcal{E}^n, \|\cdot\| \rangle: \|y - w\| < r\}$.

Let n be an element of \mathbb{N} , let Z be a set, and let f be a partial function from Z to \mathcal{R}^n . The functor $|f|$ yielding a partial function from Z to \mathbb{R} is defined by:

(Def. 2) $\text{dom } |f| = \text{dom } f$ and for every set x such that $x \in \text{dom } |f|$ holds $|f|_x = |f_x|$.

Let n be an element of \mathbb{N} , let Z be a non empty set, and let f be a partial function from Z to \mathcal{R}^n . The functor $-f$ yields a partial function from Z to \mathcal{R}^n and is defined by:

(Def. 3) $\text{dom}(-f) = \text{dom } f$ and for every set c such that $c \in \text{dom}(-f)$ holds $(-f)_c = -f_c$.

One can prove the following propositions:

- (5) Let f_1, f_2 be partial functions from \mathbb{R} to the carrier of $\langle \mathcal{E}^n, \|\cdot\| \rangle$ and g_1, g_2 be partial functions from \mathbb{R} to \mathcal{R}^n . If $f_1 = g_1$ and $f_2 = g_2$, then $f_1 + f_2 = g_1 + g_2$.
- (6) Let f_1 be a partial function from \mathbb{R} to the carrier of $\langle \mathcal{E}^n, \|\cdot\| \rangle$, g_1 be a partial function from \mathbb{R} to \mathcal{R}^n , and a be a real number. If $f_1 = g_1$, then $a \cdot f_1 = a \cdot g_1$.
- (7) For every partial function f_1 from \mathbb{R} to \mathcal{R}^n holds $(-1) \cdot f_1 = -f_1$.
- (8) Let f_1 be a partial function from \mathbb{R} to the carrier of $\langle \mathcal{E}^n, \|\cdot\| \rangle$ and g_1 be a partial function from \mathbb{R} to \mathcal{R}^n . If $f_1 = g_1$, then $-f_1 = -g_1$.
- (9) Let f_1 be a partial function from \mathbb{R} to the carrier of $\langle \mathcal{E}^n, \|\cdot\| \rangle$ and g_1 be a partial function from \mathbb{R} to \mathcal{R}^n . If $f_1 = g_1$, then $\|f_1\| = \|g_1\|$.
- (10) Let f_1, f_2 be partial functions from \mathbb{R} to the carrier of $\langle \mathcal{E}^n, \|\cdot\| \rangle$ and g_1, g_2 be partial functions from \mathbb{R} to \mathcal{R}^n . If $f_1 = g_1$ and $f_2 = g_2$, then $f_1 - f_2 = g_1 - g_2$.
- (11) f is continuous in x_0 if and only if the following conditions are satisfied:
 - (i) $x_0 \in \text{dom } f$, and
 - (ii) for every subset N_1 of \mathcal{R}^n such that there exists a real number r such that $0 < r$ and $\{y \in \mathcal{R}^n: |y - f_{x_0}| < r\} = N_1$ there exists a neighbourhood N of x_0 such that for every x_1 such that $x_1 \in \text{dom } f$ and $x_1 \in N$ holds $f_{x_1} \in N_1$.
- (12) f is continuous in x_0 if and only if the following conditions are satisfied:
 - (i) $x_0 \in \text{dom } f$, and

- (ii) for every subset N_1 of \mathcal{R}^n such that there exists a real number r such that $0 < r$ and $\{y \in \mathcal{R}^n: |y - f_{x_0}| < r\} = N_1$ there exists a neighbourhood N of x_0 such that $f \circ N \subseteq N_1$.
- (13) If there exists a neighbourhood N of x_0 such that $\text{dom } f \cap N = \{x_0\}$, then f is continuous in x_0 .
- (14) If $x_0 \in \text{dom } f_1 \cap \text{dom } f_2$ and f_1 is continuous in x_0 and f_2 is continuous in x_0 , then $f_1 + f_2$ is continuous in x_0 .
- (15) If $x_0 \in \text{dom } f_1 \cap \text{dom } f_2$ and f_1 is continuous in x_0 and f_2 is continuous in x_0 , then $f_1 - f_2$ is continuous in x_0 .
- (16) If f is continuous in x_0 , then $r \cdot f$ is continuous in x_0 .
- (17) If $x_0 \in \text{dom } f$ and f is continuous in x_0 , then $|f|$ is continuous in x_0 .
- (18) If $x_0 \in \text{dom } f$ and f is continuous in x_0 , then $-f$ is continuous in x_0 .
- (19) Let S be a real normed space, z be a point of $\langle \mathcal{E}^n, \|\cdot\| \rangle$, f_1 be a partial function from \mathbb{R} to \mathcal{R}^n , and f_2 be a partial function from the carrier of $\langle \mathcal{E}^n, \|\cdot\| \rangle$ to the carrier of S . Suppose $x_0 \in \text{dom}(f_2 \cdot f_1)$ and f_1 is continuous in x_0 and $z = (f_1)_{x_0}$ and f_2 is continuous in z . Then $f_2 \cdot f_1$ is continuous in x_0 .
- (20) Let S be a real normed space, f_1 be a partial function from \mathbb{R} to the carrier of S , and f_2 be a partial function from the carrier of S to \mathbb{R} . Suppose $x_0 \in \text{dom}(f_2 \cdot f_1)$ and f_1 is continuous in x_0 and f_2 is continuous in $(f_1)_{x_0}$. Then $f_2 \cdot f_1$ is continuous in x_0 .

Let us consider n , let f be a partial function from \mathcal{R}^n to \mathbb{R} , and let x_0 be an element of \mathcal{R}^n . We say that f is continuous in x_0 if and only if the condition (Def. 4) is satisfied.

- (Def. 4) There exists a point y_0 of $\langle \mathcal{E}^n, \|\cdot\| \rangle$ and there exists a partial function g from the carrier of $\langle \mathcal{E}^n, \|\cdot\| \rangle$ to \mathbb{R} such that $x_0 = y_0$ and $f = g$ and g is continuous in y_0 .

One can prove the following two propositions:

- (21) Let f be a partial function from \mathcal{R}^n to \mathbb{R} , h be a partial function from the carrier of $\langle \mathcal{E}^n, \|\cdot\| \rangle$ to \mathbb{R} , x_0 be an element of \mathcal{R}^n , and y_0 be a point of $\langle \mathcal{E}^n, \|\cdot\| \rangle$. Suppose $f = h$ and $x_0 = y_0$. Then f is continuous in x_0 if and only if h is continuous in y_0 .
- (22) Let f_1 be a partial function from \mathbb{R} to \mathcal{R}^n and f_2 be a partial function from \mathcal{R}^n to \mathbb{R} . Suppose $x_0 \in \text{dom}(f_2 \cdot f_1)$ and f_1 is continuous in x_0 and f_2 is continuous in $(f_1)_{x_0}$. Then $f_2 \cdot f_1$ is continuous in x_0 .

Let us consider n , f . We say that f is continuous if and only if:

- (Def. 5) For every x_0 such that $x_0 \in \text{dom } f$ holds f is continuous in x_0 .

One can prove the following propositions:

- (23) Let g be a partial function from \mathbb{R} to the carrier of $\langle \mathcal{E}^n, \|\cdot\| \rangle$ and f be a partial function from \mathbb{R} to \mathcal{R}^n . If $g = f$, then g is continuous iff f is continuous.
- (24) Suppose $X \subseteq \text{dom } f$. Then $f \upharpoonright X$ is continuous if and only if for all x_0, r such that $x_0 \in X$ and $0 < r$ there exists s such that $0 < s$ and for every x_1 such that $x_1 \in X$ and $|x_1 - x_0| < s$ holds $|f_{x_1} - f_{x_0}| < r$.

Let us consider n . Observe that every partial function from \mathbb{R} to \mathcal{R}^n which is constant is also continuous.

Let us consider n . Observe that there exists a partial function from \mathbb{R} to \mathcal{R}^n which is continuous.

Let us consider n , let f be a continuous partial function from \mathbb{R} to \mathcal{R}^n , and let X be a set. One can verify that $f \upharpoonright X$ is continuous.

One can prove the following proposition

- (25) If $f \upharpoonright X$ is continuous and $X_1 \subseteq X$, then $f \upharpoonright X_1$ is continuous.

Let us consider n . Note that every partial function from \mathbb{R} to \mathcal{R}^n which is empty is also continuous.

Let us consider n, f and let X be a trivial set. One can verify that $f \upharpoonright X$ is continuous.

Let us consider n and let f_1, f_2 be continuous partial functions from \mathbb{R} to \mathcal{R}^n . One can check that $f_1 + f_2$ is continuous.

The following propositions are true:

- (26) If $X \subseteq \text{dom } f_1 \cap \text{dom } f_2$ and $f_1 \upharpoonright X$ is continuous and $f_2 \upharpoonright X$ is continuous, then $(f_1 + f_2) \upharpoonright X$ is continuous and $(f_1 - f_2) \upharpoonright X$ is continuous.
- (27) If $X \subseteq \text{dom } f_1$ and $X_1 \subseteq \text{dom } f_2$ and $f_1 \upharpoonright X$ is continuous and $f_2 \upharpoonright X_1$ is continuous, then $(f_1 + f_2) \upharpoonright (X \cap X_1)$ is continuous and $(f_1 - f_2) \upharpoonright (X \cap X_1)$ is continuous.

Let us consider n , let f be a continuous partial function from \mathbb{R} to \mathcal{R}^n , and let us consider r . Observe that $r \cdot f$ is continuous.

The following propositions are true:

- (28) If $X \subseteq \text{dom } f$ and $f \upharpoonright X$ is continuous, then $(r \cdot f) \upharpoonright X$ is continuous.
- (29) If $X \subseteq \text{dom } f$ and $f \upharpoonright X$ is continuous, then $|f| \upharpoonright X$ is continuous and $(-f) \upharpoonright X$ is continuous.
- (30) If f is total and for all x_1, x_2 holds $f_{x_1+x_2} = f_{x_1} + f_{x_2}$ and there exists x_0 such that f is continuous in x_0 , then $f \upharpoonright \mathbb{R}$ is continuous.
- (31) For every subset Y of $\langle \mathcal{E}^n, \|\cdot\| \rangle$ such that $\text{dom } f$ is compact and $f \upharpoonright \text{dom } f$ is continuous and $Y = \text{rng } f$ holds Y is compact.
- (32) Let Y be a subset of \mathbb{R} and Z be a subset of $\langle \mathcal{E}^n, \|\cdot\| \rangle$. Suppose $Y \subseteq \text{dom } f$ and $Z = f^\circ Y$ and Y is compact and $f \upharpoonright Y$ is continuous. Then Z is compact.

Let us consider n, f . We say that f is Lipschitzian if and only if:

(Def. 6) There exists a partial function g from \mathbb{R} to the carrier of $\langle \mathcal{E}^n, \|\cdot\| \rangle$ such that $g = f$ and g is Lipschitzian.

The following propositions are true:

- (33) f is Lipschitzian if and only if there exists a real number r such that $0 < r$ and for all x_1, x_2 such that $x_1, x_2 \in \text{dom } f$ holds $|f_{x_1} - f_{x_2}| \leq r \cdot |x_1 - x_2|$.
- (34) If $f = h$, then f is Lipschitzian iff h is Lipschitzian.
- (35) $f \upharpoonright X$ is Lipschitzian if and only if there exists a real number r such that $0 < r$ and for all x_1, x_2 such that $x_1, x_2 \in \text{dom}(f \upharpoonright X)$ holds $|f_{x_1} - f_{x_2}| \leq r \cdot |x_1 - x_2|$.

Let us consider n . Note that every partial function from \mathbb{R} to \mathcal{R}^n which is empty is also Lipschitzian.

Let us consider n . Note that there exists a partial function from \mathbb{R} to \mathcal{R}^n which is empty.

Let us consider n , let f be a Lipschitzian partial function from \mathbb{R} to \mathcal{R}^n , and let X be a set. Note that $f \upharpoonright X$ is Lipschitzian.

We now state the proposition

- (36) If $f \upharpoonright X$ is Lipschitzian and $X_1 \subseteq X$, then $f \upharpoonright X_1$ is Lipschitzian.

Let us consider n and let f_1, f_2 be Lipschitzian partial functions from \mathbb{R} to \mathcal{R}^n . Observe that $f_1 + f_2$ is Lipschitzian and $f_1 - f_2$ is Lipschitzian.

We now state two propositions:

- (37) If $f_1 \upharpoonright X$ is Lipschitzian and $f_2 \upharpoonright X_1$ is Lipschitzian, then $(f_1 + f_2) \upharpoonright (X \cap X_1)$ is Lipschitzian.
- (38) If $f_1 \upharpoonright X$ is Lipschitzian and $f_2 \upharpoonright X_1$ is Lipschitzian, then $(f_1 - f_2) \upharpoonright (X \cap X_1)$ is Lipschitzian.

Let us consider n , let f be a Lipschitzian partial function from \mathbb{R} to \mathcal{R}^n , and let us consider p . Observe that $p \cdot f$ is Lipschitzian.

Next we state the proposition

- (39) If $f \upharpoonright X$ is Lipschitzian and $X \subseteq \text{dom } f$, then $(p \cdot f) \upharpoonright X$ is Lipschitzian.

Let us consider n and let f be a Lipschitzian partial function from \mathbb{R} to \mathcal{R}^n . Observe that $|f|$ is Lipschitzian.

Next we state the proposition

- (40) If $f \upharpoonright X$ is Lipschitzian, then $-f \upharpoonright X$ is Lipschitzian and $|f| \upharpoonright X$ is Lipschitzian and $(-f) \upharpoonright X$ is Lipschitzian.

Let us consider n . One can check that every partial function from \mathbb{R} to \mathcal{R}^n which is constant is also Lipschitzian.

Let us consider n . One can verify that every partial function from \mathbb{R} to \mathcal{R}^n which is Lipschitzian is also continuous.

The following propositions are true:

- (41) For all elements r, p of \mathcal{R}^n such that for every x_0 such that $x_0 \in X$ holds $f_{x_0} = x_0 \cdot r + p$ holds $f \upharpoonright X$ is continuous.

- (42) For every element x_0 of \mathcal{R}^n such that $1 \leq i \leq n$ holds $\text{proj}(i, n)$ is continuous in x_0 .
- (43) Let n be a non empty element of \mathbb{N} and h be a partial function from \mathbb{R} to \mathcal{R}^n . Then h is continuous in x_0 if and only if the following conditions are satisfied:
- (i) $x_0 \in \text{dom } h$, and
 - (ii) for every element i of \mathbb{N} such that $i \in \text{Seg } n$ holds $\text{proj}(i, n) \cdot h$ is continuous in x_0 .
- (44) Let n be a non empty element of \mathbb{N} and h be a partial function from \mathbb{R} to \mathcal{R}^n . Then h is continuous if and only if for every element i of \mathbb{N} such that $i \in \text{Seg } n$ holds $\text{proj}(i, n) \cdot h$ is continuous.
- (45) For every point x_0 of $\langle \mathcal{E}^n, \|\cdot\| \rangle$ such that $1 \leq i \leq n$ holds $\text{Proj}(i, n)$ is continuous in x_0 .
- (46) Let n be a non empty element of \mathbb{N} and h be a partial function from \mathbb{R} to the carrier of $\langle \mathcal{E}^n, \|\cdot\| \rangle$. Then h is continuous in x_0 if and only if for every element i of \mathbb{N} such that $i \in \text{Seg } n$ holds $\text{Proj}(i, n) \cdot h$ is continuous in x_0 .
- (47) Let n be a non empty element of \mathbb{N} and h be a partial function from \mathbb{R} to the carrier of $\langle \mathcal{E}^n, \|\cdot\| \rangle$. Then h is continuous if and only if for every element i of \mathbb{N} such that $i \in \text{Seg } n$ holds $\text{Proj}(i, n) \cdot h$ is continuous.

REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [3] Czesław Byliński. The complex numbers. *Formalized Mathematics*, 1(3):507–513, 1990.
- [4] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [5] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [6] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [7] Agata Darmochwał. The Euclidean space. *Formalized Mathematics*, 2(4):599–603, 1991.
- [8] Noboru Endou and Yasunari Shidama. Completeness of the real Euclidean space. *Formalized Mathematics*, 13(4):577–580, 2005.
- [9] Noboru Endou, Yasunari Shidama, and Keiichi Miyajima. Partial differentiation on normed linear spaces \mathcal{R}^n . *Formalized Mathematics*, 15(2):65–72, 2007, doi:10.2478/v10037-007-0008-5.
- [10] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [11] Artur Kornilowicz. Arithmetic operations on functions from sets into functional sets. *Formalized Mathematics*, 17(1):43–60, 2009, doi:10.2478/v10037-009-0005-y.
- [12] Keiichi Miyajima and Yasunari Shidama. Riemann integral of functions from \mathbb{R} into \mathcal{R}^n . *Formalized Mathematics*, 17(2):179–185, 2009, doi: 10.2478/v10037-009-0021-y.
- [13] Takaya Nishiyama, Keiji Ohkubo, and Yasunari Shidama. The continuous functions on normed linear spaces. *Formalized Mathematics*, 12(3):269–275, 2004.
- [14] Hiroyuki Okazaki, Noboru Endou, and Yasunari Shidama. More on continuous functions on normed linear spaces. *Formalized Mathematics*, 19(1):45–49, 2011, doi: 10.2478/v10037-011-0008-3.

- [15] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(1):223–230, 1990.
- [16] Jan Popiołek. Real normed space. *Formalized Mathematics*, 2(1):111–115, 1991.
- [17] Konrad Raczkowski and Paweł Sadowski. Real function continuity. *Formalized Mathematics*, 1(4):787–791, 1990.
- [18] Konrad Raczkowski and Paweł Sadowski. Topological properties of subsets in real numbers. *Formalized Mathematics*, 1(4):777–780, 1990.
- [19] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [20] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [21] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.
- [22] Hiroshi Yamazaki and Yasunari Shidama. Algebra of vector functions. *Formalized Mathematics*, 3(2):171–175, 1992.

Received February 22, 2011
