

# The Linearity of Riemann Integral on Functions from $\mathbb{R}$ into Real Banach Space<sup>1</sup>

Keiko Narita  
Hirosaki-city  
Aomori, Japan

Noboru Endou  
Gifu National College of Technology  
Japan

Yasunari Shidama  
Shinshu University  
Nagano, Japan

**Summary.** In this article, we described basic properties of Riemann integral on functions from  $\mathbb{R}$  into Real Banach Space. We proved mainly the linearity of integral operator about the integral of continuous functions on closed interval of the set of real numbers. These theorems were based on the article [10] and we referred to the former articles about Riemann integral. We applied definitions and theorems introduced in the article [9] and the article [11] to the proof. Using the definition of the article [10], we also proved some theorems on bounded functions.

MSC: 26A42 03B35

Keywords: formalization of Riemann integral

MML identifier: INTEGR21, version: 8.1.02 5.19.1189

The notation and terminology used in this paper have been introduced in the following articles: [2], [12], [3], [4], [9], [10], [7], [8], [16], [1], [17], [13], [14], [5], [15], [20], [21], [18], [19], [22], and [6].

## 1. SOME PROPERTIES OF BOUNDED FUNCTIONS

In this paper  $Z$  denotes a real normed space,  $a, b, c, d, e, r$  denote real numbers, and  $A, B$  denote non empty closed interval subsets of  $\mathbb{R}$ .

Let us consider a partial function  $f$  from  $\mathbb{R}$  to the carrier of  $Z$ . Now we state the propositions:

---

<sup>1</sup>This work was supported by JSPS KAKENHI 22300285, 23500029.

- (1) If  $f$  is bounded and  $A \subseteq \text{dom } f$ , then  $f \upharpoonright A$  is bounded.
- (2) If  $f \upharpoonright A$  is bounded and  $B \subseteq A$  and  $B \subseteq \text{dom}(f \upharpoonright A)$ , then  $f \upharpoonright B$  is bounded.
- (3) If  $a \leq c \leq d \leq b$  and  $f \upharpoonright [a, b]$  is bounded and  $[a, b] \subseteq \text{dom } f$ , then  $f \upharpoonright [c, d]$  is bounded.

Now we state the proposition:

- (4) Let us consider sets  $X, Y$  and partial functions  $f_1, f_2$  from  $\mathbb{R}$  to the carrier of  $Z$ . Suppose
  - (i)  $f_1 \upharpoonright X$  is bounded, and
  - (ii)  $f_2 \upharpoonright Y$  is bounded.

Then

- (iii)  $(f_1 + f_2) \upharpoonright (X \cap Y)$  is bounded, and
- (iv)  $(f_1 - f_2) \upharpoonright (X \cap Y)$  is bounded.

Let us consider a set  $X$  and a partial function  $f$  from  $\mathbb{R}$  to the carrier of  $Z$ . Now we state the propositions:

- (5) If  $f \upharpoonright X$  is bounded, then  $(r \cdot f) \upharpoonright X$  is bounded.
- (6) If  $f \upharpoonright X$  is bounded, then  $(-f) \upharpoonright X$  is bounded.

Now we state the propositions:

- (7) Let us consider a function  $f$  from  $A$  into the carrier of  $Z$ . Then  $f$  is bounded if and only if  $\|f\|$  is bounded.
- (8) Let us consider a partial function  $f$  from  $\mathbb{R}$  to the carrier of  $Z$ . Suppose  $A \subseteq \text{dom } f$ . Then  $\|f \upharpoonright A\| = \|f\| \upharpoonright A$ .
- (9) Let us consider a partial function  $g$  from  $\mathbb{R}$  to the carrier of  $Z$ . Suppose
  - (i)  $A \subseteq \text{dom } g$ , and
  - (ii)  $g \upharpoonright A$  is bounded.

Then  $\|g\| \upharpoonright A$  is bounded. The theorem is a consequence of (8) and (7).

## 2. SOME PROPERTIES OF INTEGRAL OF CONTINUOUS FUNCTIONS

In the sequel  $X, Y$  denote real Banach spaces and  $E$  denotes a point of  $Y$ .

Let us consider a real normed space  $Y$  and a continuous partial function  $f$  from  $\mathbb{R}$  to the carrier of  $Y$ . Now we state the propositions:

- (10) If  $a \leq b$  and  $[a, b] \subseteq \text{dom } f$ , then  $\|f\| \upharpoonright [a, b]$  is bounded.
- (11) If  $a \leq b$  and  $[a, b] \subseteq \text{dom } f$ , then  $f \upharpoonright [a, b]$  is bounded.
- (12) If  $a \leq b$  and  $[a, b] \subseteq \text{dom } f$ , then  $\|f\|$  is integrable on  $[a, b]$ .

Now we state the propositions:

- (13) Let us consider a continuous partial function  $f$  from  $\mathbb{R}$  to the carrier of  $Y$ . Suppose

- (i)  $a \leq c \leq d \leq b$ , and
- (ii)  $[a, b] \subseteq \text{dom } f$ .

Then  $f$  is integrable on  $[c, d]$ .

- (14) Let us consider a partial function  $f$  from  $\mathbb{R}$  to the carrier of  $Y$ . Suppose
- (i)  $a \leq b$ , and
  - (ii)  $[a, b] \subseteq \text{dom } f$ .

$$\text{Then } \int_b^a f(x)dx = - \int_a^b f(x)dx.$$

- (15) Let us consider a continuous partial function  $f$  from  $\mathbb{R}$  to the carrier of  $Y$ . Suppose
- (i)  $a \leq b$ , and
  - (ii)  $[a, b] \subseteq \text{dom } f$ , and
  - (iii)  $c \in [a, b]$ .

Then

- (iv)  $f$  is integrable on  $[a, c]$ , and
- (v)  $f$  is integrable on  $[c, b]$ , and
- (vi)  $\int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx$ .

The theorem is a consequence of (13).

- (16) Let us consider continuous partial functions  $f, g$  from  $\mathbb{R}$  to the carrier of  $Y$ . Suppose
- (i)  $a \leq c \leq d \leq b$ , and
  - (ii)  $[a, b] \subseteq \text{dom } f$ , and
  - (iii)  $[a, b] \subseteq \text{dom } g$ .

Then

- (iv)  $f + g$  is integrable on  $[c, d]$ , and
- (v)  $(f + g)|_{[c, d]}$  is bounded.

The theorem is a consequence of (13), (11), (3), and (4).

Let us consider a continuous partial function  $f$  from  $\mathbb{R}$  to the carrier of  $Y$ . Now we state the propositions:

- (17) If  $a \leq c \leq d \leq b$  and  $[a, b] \subseteq \text{dom } f$ , then  $r \cdot f$  is integrable on  $[c, d]$  and  $(r \cdot f)|_{[c, d]}$  is bounded.
- (18) Suppose  $a \leq c \leq d \leq b$  and  $f$  is integrable on  $[a, b]$  and  $f|_{[a, b]}$  is bounded and  $[a, b] \subseteq \text{dom } f$ . Then

- (i)  $-f$  is integrable on  $[c, d]$ , and
- (ii)  $(-f) \upharpoonright [c, d]$  is bounded.

Now we state the proposition:

- (19) Let us consider continuous partial functions  $f, g$  from  $\mathbb{R}$  to the carrier of  $Y$ . Suppose

- (i)  $a \leq c \leq d \leq b$ , and
- (ii)  $[a, b] \subseteq \text{dom } f$ , and
- (iii)  $[a, b] \subseteq \text{dom } g$ .

Then

- (iv)  $f - g$  is integrable on  $[c, d]$ , and
- (v)  $(f - g) \upharpoonright [c, d]$  is bounded.

The theorem is a consequence of (11), (13), (3), and (4).

Let us consider a partial function  $f$  from  $\mathbb{R}$  to the carrier of  $Y$ . Now we state the propositions:

- (20) Suppose  $A \subseteq \text{dom } f$  and  $f \upharpoonright A$  is bounded and  $f$  is integrable on  $A$  and  $\|f\|$  is integrable on  $A$ . Then  $\|\int_A f(x)dx\| \leq \int_A \|f\|(x)dx$ .

- (21) Suppose  $a \leq b$  and  $[a, b] \subseteq \text{dom } f$  and  $f$  is integrable on  $[a, b]$  and  $\|f\|$  is integrable on  $[a, b]$  and  $f \upharpoonright [a, b]$  is bounded. Then  $\|\int_a^b f(x)dx\| \leq \int_a^b \|f\|(x)dx$ .

Let us consider a continuous partial function  $f$  from  $\mathbb{R}$  to the carrier of  $Y$ . Now we state the propositions:

- (22) Suppose  $a \leq b$  and  $[a, b] \subseteq \text{dom } f$  and  $c, d \in [a, b]$ . Then

- (i)  $\|f\|$  is integrable on  $[\min(c, d), \max(c, d)]$ , and
- (ii)  $\|f\| \upharpoonright [\min(c, d), \max(c, d)]$  is bounded, and

$$(iii) \left\| \int_c^d f(x)dx \right\| \leq \int_{\min(c,d)}^{\max(c,d)} \|f\|(x)dx.$$

- (23) If  $a \leq b$  and  $[a, b] \subseteq \text{dom } f$  and  $c, d \in [a, b]$ , then  $\int_c^d (r \cdot f)(x)dx =$

$$r \cdot \int_c^d f(x)dx.$$

- (24) Suppose  $a \leq b$  and  $[a, b] \subseteq \text{dom } f$  and  $c, d \in [a, b]$ . Then  $\int_c^d -f(x)dx = -\int_c^d f(x)dx$ .
- (25) Suppose  $a \leq b$  and  $[a, b] \subseteq \text{dom } f$  and  $c, d \in [a, b]$  and for every real number  $x$  such that  $x \in [\min(c, d), \max(c, d)]$  holds  $\|f_x\| \leq e$ . Then  $\|\int_c^d f(x)dx\| \leq e \cdot |d - c|$ .

Now we state the propositions:

- (26) Let us consider a real normed space  $Y$ , a non empty closed interval subset  $A$  of  $\mathbb{R}$ , a function  $f$  from  $A$  into the carrier of  $Y$ , and a point  $E$  of  $Y$ . Suppose  $\text{rng } f = \{E\}$ . Then
- (i)  $f$  is integrable, and
  - (ii)  $\text{integral } f = \text{vol}(A) \cdot E$ .
- PROOF: Reconsider  $I = \text{vol}(A) \cdot E$  as a point of  $Y$ . For every division sequence  $T$  of  $A$  and for every middle volume sequence  $S$  of  $f$  and  $T$  such that  $\delta_T$  is convergent and  $\lim \delta_T = 0$  holds middle sum( $f, S$ ) is convergent and  $\lim \text{middle sum}(f, S) = I$  by [11, (6)], [20, (70)], [11, (7)].  $\square$
- (27) Let us consider a partial function  $f$  from  $\mathbb{R}$  to the carrier of  $Y$  and a point  $E$  of  $Y$ . Suppose
- (i)  $a \leq b$ , and
  - (ii)  $[a, b] \subseteq \text{dom } f$ , and
  - (iii) for every real number  $x$  such that  $x \in [a, b]$  holds  $f_x = E$ .

Then

- (iv)  $f$  is integrable on  $[a, b]$ , and
- (v)  $\int_a^b f(x)dx = (b - a) \cdot E$ .

The theorem is a consequence of (26). PROOF: Reconsider  $A = [a, b]$  as a non empty closed interval subset of  $\mathbb{R}$ . Reconsider  $g = f \upharpoonright A$  as a function from  $A$  into the carrier of  $Y$ .  $\{E\} \subseteq \text{rng } g$  by [19, (4)], [3, (49), (3)].  $\text{rng } g \subseteq \{E\}$  by [5, (3)], [3, (49)].  $\square$

- (28) Let us consider a partial function  $f$  from  $\mathbb{R}$  to the carrier of  $Y$ . Suppose
- (i)  $a \leq b$ , and
  - (ii)  $c, d \in [a, b]$ , and
  - (iii)  $[a, b] \subseteq \text{dom } f$ , and

(iv) for every real number  $x$  such that  $x \in [a, b]$  holds  $f_x = E$ .

Then  $\int_c^d f(x)dx = (d - c) \cdot E$ . The theorem is a consequence of (27) and (14).

(29) Let us consider a continuous partial function  $f$  from  $\mathbb{R}$  to the carrier of  $Y$ . Suppose

- (i)  $a \leq b$ , and
- (ii)  $[a, b] \subseteq \text{dom } f$ , and
- (iii)  $c, d \in [a, b]$ .

Then  $\int_a^d f(x)dx = \int_a^c f(x)dx + \int_c^d f(x)dx$ . The theorem is a consequence of (14).

(30) Let us consider continuous partial functions  $f, g$  from  $\mathbb{R}$  to the carrier of  $Y$ . Suppose

- (i)  $a \leq b$ , and
- (ii)  $[a, b] \subseteq \text{dom } f$ , and
- (iii)  $[a, b] \subseteq \text{dom } g$ , and
- (iv)  $c, d \in [a, b]$ .

Then  $\int_c^d (f - g)(x)dx = \int_c^d f(x)dx - \int_c^d g(x)dx$ . The theorem is a consequence of (14).

## REFERENCES

- [1] Józef Białas. Properties of the intervals of real numbers. *Formalized Mathematics*, 3(2):263–269, 1992.
- [2] Czesław Byliński. The complex numbers. *Formalized Mathematics*, 1(3):507–513, 1990.
- [3] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [4] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [5] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [6] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [7] Noboru Endou and Artur Kornilowicz. The definition of the Riemann definite integral and some related lemmas. *Formalized Mathematics*, 8(1):93–102, 1999.
- [8] Noboru Endou, Katsumi Wasaki, and Yasunari Shidama. Definition of integrability for partial functions from  $\mathbb{R}$  to  $\mathbb{R}$  and integrability for continuous functions. *Formalized Mathematics*, 9(2):281–284, 2001.
- [9] Keiichi Miyajima, Takahiro Kato, and Yasunari Shidama. Riemann integral of functions from  $\mathbb{R}$  into real normed space. *Formalized Mathematics*, 19(1):17–22, 2011. doi:10.2478/v10037-011-0003-8.

- [10] Keiichi Miyajima, Artur Kornilowicz, and Yasunari Shidama. Riemann integral of functions from  $\mathbb{R}$  into  $n$ -dimensional real normed space. *Formalized Mathematics*, 20(1):79–86, 2012. doi:10.2478/v10037-012-0011-3.
- [11] Keiko Narita, Noboru Endou, and Yasunari Shidama. Riemann integral of functions from  $\mathbb{R}$  into real Banach space. *Formalized Mathematics*, 21(2):145–152, 2013. doi:10.2478/forma-2013-0016.
- [12] Adam Naumowicz. Conjugate sequences, bounded complex sequences and convergent complex sequences. *Formalized Mathematics*, 6(2):265–268, 1997.
- [13] Hiroyuki Okazaki, Noboru Endou, and Yasunari Shidama. More on continuous functions on normed linear spaces. *Formalized Mathematics*, 19(1):45–49, 2011. doi:10.2478/v10037-011-0008-3.
- [14] Jan Popiolek. Real normed space. *Formalized Mathematics*, 2(1):111–115, 1991.
- [15] Konrad Raczkowski and Paweł Sadowski. Topological properties of subsets in real numbers. *Formalized Mathematics*, 1(4):777–780, 1990.
- [16] Yasunari Shidama. Banach space of bounded linear operators. *Formalized Mathematics*, 12(1):39–48, 2004.
- [17] Andrzej Trybulec. On the sets inhabited by numbers. *Formalized Mathematics*, 11(4):341–347, 2003.
- [18] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(2):291–296, 1990.
- [19] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [20] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [21] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.
- [22] Hiroshi Yamazaki and Yasunari Shidama. Algebra of vector functions. *Formalized Mathematics*, 3(2):171–175, 1992.

*Received October 7, 2013*

---