

# Equivalent Expressions of Direct Sum Decomposition of Groups<sup>1</sup>

Kazuhisa Nakasho  
Shinshu University  
Nagano, Japan

Hiroyuki Okazaki  
Shinshu University  
Nagano, Japan

Hiroshi Yamazaki  
Shinshu University  
Nagano, Japan

Yasunari Shidama  
Shinshu University  
Nagano, Japan

**Summary.** In this article, the equivalent expressions of the direct sum decomposition of groups are mainly discussed. In the first section, we formalize the fact that the internal direct sum decomposition can be defined as normal subgroups and some of their properties. In the second section, we formalize an equivalent form of internal direct sum of commutative groups. In the last section, we formalize that the external direct sum leads an internal direct sum. We referred to [19], [18] [8] and [14] in the formalization.

MSC: 20E34 03B35

Keywords: group theory; direct sum decomposition

MML identifier: GROUP\_20, version: 8.1.04 5.31.1231

The notation and terminology used in this paper have been introduced in the following articles: [1], [20], [6], [9], [10], [7], [22], [17], [16], [23], [24], [25], [26], [13], [3], [5], [11], [15], [28], [29], [27], and [12].

## 1. INTERNAL DIRECT SUM DECOMPOSITION INTO NORMAL SUBGROUPS

Let  $I$  be a set and  $G$  be a group.

A subgroup-family of  $I$  and  $G$  is a group-family of  $I$  and is defined by

(Def. 1) for every object  $i$  such that  $i \in I$  holds  $it(i)$  is a subgroup of  $G$ .

---

<sup>1</sup>This work was supported by JSPS KAKENHI 22300285.

Let  $F$  be a subgroup-family of  $I$  and  $G$ . We say that  $F$  is component commutative if and only if

(Def. 2) for every elements  $i, j$  of  $I$  and for every elements  $g_1, g_2$  of  $G$  such that  $i \neq j$  and  $g_1 \in F(i)$  and  $g_2 \in F(j)$  holds  $g_1 \cdot g_2 = g_2 \cdot g_1$ .

Let  $I$  be a non empty set. One can verify that there exists a subgroup-family of  $I$  and  $G$  which is component commutative.

Now we state the propositions:

- (1) Let us consider a group  $G$ , a normal subgroup  $H$  of  $G$ , and elements  $x, y$  of  $G$ . Suppose  $y \in H$ . Then  $x \cdot y \cdot x^{-1}, x \cdot (y \cdot x^{-1}) \in H$ .
- (2) Let us consider a non empty set  $I$ , a group  $G$ , a group-family  $F$  of  $I$ , and a function  $x$  from  $I$  into  $G$ . Suppose  $x \in \prod F$ . Then  $x$  is a function from  $I$  into  $\bigcup(\text{the support of } F)$ .

PROOF: For every object  $z$  such that  $z \in \text{rng } x$  holds  $z \in \bigcup(\text{the support of } F)$  by [10, (11)], [16, (5), (4)], [9, (3)].  $\square$

- (3) Let us consider a non empty set  $I$ , a group  $G$ , a subgroup  $H$  of  $G$ , a function  $x$  from  $I$  into  $G$ , and a function  $y$  from  $I$  into  $H$ . If  $x = y$ , then  $\text{support } x = \text{support } y$ .

PROOF: For every object  $i, i \in \text{support } x$  iff  $i \in \text{support } y$  by [23, (44)].  $\square$

- (4) Let us consider a non empty set  $I$ , a group  $G$ , and a subgroup  $H$  of  $G$ . Then every finite-support function from  $I$  into  $H$  is a finite-support function from  $I$  into  $G$ . The theorem is a consequence of (3).
- (5) Let us consider a non empty set  $I$ , a group  $G$ , a subgroup  $H$  of  $G$ , and a finite-support function  $x$  from  $I$  into  $G$ . Suppose  $\text{rng } x \subseteq \Omega_H$ . Then  $x$  is a finite-support function from  $I$  into  $H$ . The theorem is a consequence of (3).
- (6) Let us consider a non empty set  $I$ , a group  $G$ , a subgroup  $H$  of  $G$ , a finite-support function  $x$  from  $I$  into  $G$ , and a finite-support function  $y$  from  $I$  into  $H$ . If  $x = y$ , then  $\prod x = \prod y$ . The theorem is a consequence of (3).
- (7) Let us consider a function  $f$ , and sets  $i, x$ . Then  $f = (f + \cdot (i, x)) + \cdot (i, f(i))$ .
- (8) Let us consider a non empty set  $I$ , a group  $G$ , a component commutative subgroup-family  $F$  of  $I$  and  $G$ , finite-support functions  $x, y$  from  $I$  into  $G$ , and an element  $i$  of  $I$ . Suppose  $y = x + \cdot (i, \mathbf{1}_{F(i)})$  and  $x \in \prod F$ . Then  $\prod x = \prod y \cdot x(i) = x(i) \cdot \prod y$ .

PROOF: Reconsider  $p_2 = y$  as an element of  $\prod F$ . Reconsider  $s_1 = p_2$  as an element of  $\text{sum } F$ . Set  $z = \mathbf{1}_{\prod F} + \cdot (i, x(i))$ . Reconsider  $s_2 = z$  as an element of  $\text{sum } F$ .  $x = s_1 \cdot s_2$  by [16, (5), (24)], [23, (40)], [7, (31)].

$s_1 \cdot s_2 = s_2 \cdot s_1$  by [16, (27), (17)], [23, (43)], [16, (32)].  $\square$

- (9) Let us consider a non empty set  $I$ , a group  $G$ , a component commutative subgroup-family  $F$  of  $I$  and  $G$ , a subset  $U$  of  $G$ , an element  $i$  of  $I$ , and finite-support functions  $x, y$  from  $I$  into  $\text{gr}(U)$ . Suppose  $y = x + \cdot (i, \mathbf{1}_{F(i)})$  and  $x \in \prod F$ . Then  $\prod x = \prod y \cdot x(i) = x(i) \cdot \prod y$ . The theorem is a consequence of (4), (6), and (8).
- (10) Let us consider a non empty set  $I$ , a group  $G$ , a component commutative subgroup-family  $F$  of  $I$  and  $G$ , a subset  $U$  of  $G$ , a finite-support function  $y$  from  $I$  into  $\text{gr}(U)$ , an element  $i$  of  $I$ , and an element  $g$  of  $\text{gr}(U)$ . Suppose  $y \in \prod F$  and  $y(i) = \mathbf{1}_{F(i)}$  and  $g \in F(i)$ . Then  $\prod y \cdot g = g \cdot \prod y$ . The theorem is a consequence of (7) and (9).
- (11) Let us consider a non empty set  $I$ , a group  $G$ , a component commutative subgroup-family  $F$  of  $I$  and  $G$ , and a subset  $U$  of  $G$ . Suppose  $U = \bigcup(\text{the support of } F)$ . Let us consider an element  $g$  of  $G$ , a finite sequence  $H$  of elements of  $G$ , and a finite sequence  $K$  of elements of  $\mathbb{Z}$ . Suppose  $\text{len } H = \text{len } K$  and  $\text{rng } H \subseteq U$  and  $\prod H^K = g$ . Then there exists a finite-support function  $f$  from  $I$  into  $G$  such that

(i)  $f \in \prod F$ , and

(ii)  $g = \prod f$ .

PROOF: Define  $\mathcal{P}[\text{natural number}] \equiv$  for every element  $g$  of  $G$  for every finite sequence  $H$  of elements of  $G$  for every finite sequence  $K$  of elements of  $\mathbb{Z}$  such that  $\text{len } H = \mathbb{N}$  and  $\text{len } H = \text{len } K$  and  $\text{rng } H \subseteq U$  and  $\prod H^K = g$  there exists a finite-support function  $f$  from  $I$  into  $G$  such that  $f \in \prod F$  and  $g = \prod f$ .  $\mathcal{P}[0]$  by [25, (21)], [16, (12), (13), (16)]. For every natural number  $n$  such that  $\mathcal{P}[n]$  holds  $\mathcal{P}[n+1]$  by [28, (70)], [6, (4)], [21, (55)], [9, (3)]. For every natural number  $n$ ,  $\mathcal{P}[n]$  from [4, Sch. 2].  $\square$

- (12) Let us consider a non empty set  $I$ , a group  $G$ , a subgroup-family  $F$  of  $I$  and  $G$ , finite-support functions  $h, h_0$  from  $I$  into  $G$ , an element  $i$  of  $I$ , and a subset  $U_1$  of  $G$ . Suppose  $U_1 = \bigcup((\text{the support of } F) \setminus \{i\})$  and  $h_0 = h + \cdot (i, \mathbf{1}_{F(i)})$  and  $h \in \prod F$ . Then  $\prod h_0 \in \text{gr}(U_1)$ .

PROOF: For every object  $y$  such that  $y \in \text{rng } h_0$  holds  $y \in \Omega_{\text{gr}(U_1)}$  by [10, (113)], [7, (32)], [16, (5), (4)]. Reconsider  $x_0 = h_0$  as a finite-support function from  $I$  into  $\text{gr}(U_1)$ .  $\prod x_0 = \prod h_0$ .  $\square$

- (13) Let us consider a non empty set  $I$ , a group  $G$ , a component commutative subgroup-family  $F$  of  $I$  and  $G$ , and a subset  $U$  of  $G$ . Suppose  $U = \bigcup(\text{the support of } F)$ . Let us consider an element  $g$  of  $G$ . Suppose  $g \in \text{gr}(U)$ . Then there exists a finite-support function  $f$  from  $I$  into  $\text{gr}(U)$  such that

- (i)  $f \in \text{sum } F$ , and
- (ii)  $g = \prod f$ .

The theorem is a consequence of (11), (2), (5), and (6).

- (14) Let us consider a non empty set  $I$ , a group  $G$ , a component commutative subgroup-family  $F$  of  $I$  and  $G$ , and a subset  $U$  of  $G$ . Suppose  $U = \bigcup(\text{the support of } F)$ . Let us consider an element  $i$  of  $I$ . Then  $F(i)$  is a normal subgroup of  $\text{gr}(U)$ .

PROOF: Reconsider  $F_1 = F(i)$  as a subgroup of  $\text{gr}(U)$ . For every element  $a$  of  $\text{gr}(U)$ ,  $a \cdot F_1 \subseteq F_1 \cdot a$  by [23, (103), (42)], (13), [23, (40)].  $\square$

- (15) Let us consider a non empty set  $I$ , a group  $G$ , and a component commutative subgroup-family  $F$  of  $I$  and  $G$ . Suppose for every element  $i$  of  $I$ , there exists a subset  $U_1$  of  $G$  such that  $U_1 = \bigcup((\text{the support of } F) \upharpoonright (I \setminus \{i\}))$  and  $\Omega_{\text{gr}(U_1)} \cap \Omega_{F(i)} = \{\mathbf{1}_G\}$ . Let us consider finite-support functions  $x_1, x_2$  from  $I$  into  $G$ . If  $x_1, x_2 \in \text{sum } F$  and  $\prod x_1 = \prod x_2$ , then  $x_1 = x_2$ .

PROOF: Define  $\mathcal{P}[\text{natural number}] \equiv$  for every finite-support functions  $x_1, x_2$  from  $I$  into  $G$  such that  $\overline{\text{support } x_1} = \mathbb{S}_1$  and  $x_1, x_2 \in \text{sum } F$  and  $\prod x_1 = \prod x_2$  holds  $x_1 = x_2$ .  $\mathcal{P}[0]$  by [16, (15), (14)], [23, (42)], [16, (26)]. For every natural number  $k$  such that  $\mathcal{P}[k]$  holds  $\mathcal{P}[k+1]$  by [23, (42)], [16, (26)], [23, (44)], [16, (30), (25)]. For every natural number  $k$ ,  $\mathcal{P}[k]$  from [4, Sch. 2].  $\square$

- (16) Let us consider a non empty set  $I$ , a strict group  $G$ , and a group-family  $F$  of  $I$ . Then  $F$  is an internal direct sum components of  $G$  and  $I$  if and only if for every element  $i$  of  $I$ ,  $F(i)$  is a normal subgroup of  $G$  and there exists a subset  $U$  of  $G$  such that  $U = \bigcup(\text{the support of } F)$  and  $\text{gr}(U) = G$  and for every element  $i$  of  $I$ , there exists a subset  $U_1$  of  $G$  such that  $U_1 = \bigcup((\text{the support of } F) \upharpoonright (I \setminus \{i\}))$  and  $\Omega_{\text{gr}(U_1)} \cap \Omega_{F(i)} = \{\mathbf{1}_G\}$ .

PROOF: Consider  $U$  being a subset of  $G$  such that  $U = \bigcup(\text{the support of } F)$  and  $\text{gr}(U) = G$ . For every elements  $i, j$  of  $I$  such that  $i \neq j$  holds  $\Omega_{F(i)} \cap \Omega_{F(j)} = \{\mathbf{1}_G\}$  by [23, (46)], [12, (31)], [28, (62)], [9, (49)]. For every elements  $i, j$  of  $I$  and for every elements  $g_1, g_2$  of  $G$  such that  $i \neq j$  and  $g_1 \in F(i)$  and  $g_2 \in F(j)$  holds  $g_1 \cdot g_2 = g_2 \cdot g_1$  by [23, (51)], (1), [22, (17)], [23, (50)]. For every element  $y$  of  $G$ , there exists a finite-support function  $x$  from  $I$  into  $G$  such that  $x \in \text{sum } F$  and  $y = \prod x$ . For every finite-support functions  $x_1, x_2$  from  $I$  into  $G$  such that  $x_1, x_2 \in \text{sum } F$  and  $\prod x_1 = \prod x_2$  holds  $x_1 = x_2$ .  $\square$

## 2. INTERNAL DIRECT SUM DECOMPOSITION FOR COMMUTATIVE GROUP

Now we state the proposition:

- (17) Let us consider a non empty set  $I$ , a commutative group  $G$ , and a group-family  $F$  of  $I$ . Then  $F$  is an internal direct sum components of  $G$  and  $I$  if and only if for every element  $i$  of  $I$ ,  $F(i)$  is a subgroup of  $G$  and for every elements  $i, j$  of  $I$  such that  $i \neq j$  holds  $\Omega_{F(i)} \cap \Omega_{F(j)} = \{1_G\}$  and for every element  $y$  of  $G$ , there exists a finite-support function  $x$  from  $I$  into  $G$  such that  $x \in \text{sum } F$  and  $y = \prod x$  and for every finite-support functions  $x_1, x_2$  from  $I$  into  $G$  such that  $x_1, x_2 \in \text{sum } F$  and  $\prod x_1 = \prod x_2$  holds  $x_1 = x_2$ .

## 3. EQUIVALENCE BETWEEN INTERNAL AND EXTERNAL DIRECT SUM

Now we state the propositions:

- (18) Let us consider a non empty set  $I$ , a group  $G$ , a subgroup-family  $F$  of  $I$  and  $G$ , a homomorphism  $h$  from  $\text{sum } F$  to  $G$ , and a finite-support function  $a$  from  $I$  into  $G$ . Suppose  $a \in \text{sum } F$  and for every element  $i$  of  $I$  and for every element  $x$  of  $F(i)$ ,  $h((1\text{ProdHom}(F, i))(x)) = x$ . Then  $h(a) = \prod a$ .  
 PROOF: Define  $\mathcal{P}[\text{natural number}] \equiv$  for every finite-support function  $b$  from  $I$  into  $G$  such that  $b \in \text{sum } F$  holds if  $\overline{\text{support } b} = \$1$ , then  $h(b) = \prod b$ .  $\mathcal{P}[0]$  by [16, (14)], [23, (44)], [26, (31)], [16, (15)]. For every natural number  $k$  such that  $\mathcal{P}[k]$  holds  $\mathcal{P}[k + 1]$  by [16, (25)], [23, (44)], [16, (26)], [23, (40)]. For every natural number  $k$ ,  $\mathcal{P}[k]$  from [4, Sch. 2]. Consider  $k$  being a natural number such that  $\overline{\text{support } a} = k$ .  $\square$
- (19) Let us consider a non empty set  $I$ , a group  $G$ , and a direct sum components  $M$  of  $G$  and  $I$ . Then there exists a homomorphism  $f$  from  $\text{sum } M$  to  $G$  and there exists an internal direct sum components  $N$  of  $G$  and  $I$  such that  $f$  is bijective and for every element  $i$  of  $I$ , there exists a homomorphism  $q_1$  from  $M(i)$  to  $N(i)$  such that  $q_1 = f \cdot 1\text{ProdHom}(M, i)$  and  $q_1$  is bijective.  
 PROOF: Consider  $f$  being a homomorphism from  $\text{sum } M$  to  $G$  such that  $f$  is bijective. Define  $\mathcal{D}(\text{element of } I) = f^\circ(\text{ProjGroup}(M, \$1))$ . Consider  $N$  being a function such that  $\text{dom } N = I$  and for every element  $i$  of  $I$  such that  $i \in I$  holds  $N(i) = \mathcal{D}(i)$  from [2, Sch. 2]. For every object  $i$  such that  $i \in I$  holds  $N(i)$  is a strict subgroup of  $G$ . Define  $\mathcal{E}(\text{element of } I) = f \cdot 1\text{ProdHom}(M, \$1)$ . Consider  $q$  being a function such that  $\text{dom } q = I$  and for every element  $i$  of  $I$  such that  $i \in I$  holds  $q(i) = \mathcal{E}(i)$  from [2, Sch. 2]. Reconsider  $r = \text{SumMap}(M, N, q)$  as a homomorphism from  $\text{sum } M$  to  $\text{sum } N$ . Reconsider  $s = r^{-1}$  as a homomorphism from  $\text{sum } N$  to  $\text{sum } M$ .

Reconsider  $g = f \cdot s$  as a function. For every element  $i$  of  $I$  and for every element  $n$  of  $N(i)$ ,  $g((1\text{ProdHom}(N, i))(n)) = n$  by [16, (42)], [23, (40)], [9, (13), (34)]. For every finite-support function  $a$  from  $I$  into  $G$  such that  $a \in \text{sum } N$  holds  $g(a) = \prod a$ . For every element  $i$  of  $I$ , there exists a homomorphism  $q_1$  from  $M(i)$  to  $N(i)$  such that  $q_1 = f \cdot 1\text{ProdHom}(M, i)$  and  $q_1$  is bijective.  $\square$

## REFERENCES

- [1] Grzegorz Bancerek. König's theorem. *Formalized Mathematics*, 1(3):589–593, 1990.
- [2] Grzegorz Bancerek. Tarski's classes and ranks. *Formalized Mathematics*, 1(3):563–567, 1990.
- [3] Grzegorz Bancerek. Monoids. *Formalized Mathematics*, 3(2):213–225, 1992.
- [4] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [5] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [6] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [7] Grzegorz Bancerek and Andrzej Trybulec. Miscellaneous facts about functions. *Formalized Mathematics*, 5(4):485–492, 1996.
- [8] Nicolas Bourbaki. *Elements of Mathematics. Algebra I. Chapters 1-3*. Springer-Verlag, Berlin, Heidelberg, New York, London, Paris, Tokyo, 1989.
- [9] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [10] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [11] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [12] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [13] Artur Kornilowicz. The product of the families of the groups. *Formalized Mathematics*, 7(1):127–134, 1998.
- [14] Serge Lang. *Algebra*. Springer, 3rd edition, 2005.
- [15] Beata Madras. Product of family of universal algebras. *Formalized Mathematics*, 4(1):103–108, 1993.
- [16] Kazuhisa Nakasho, Hiroshi Yamazaki, Hiroyuki Okazaki, and Yasunari Shidama. Definition and properties of direct sum decomposition of groups. *Formalized Mathematics*, 23(1):15–27, 2015. doi:10.2478/forma-2015-0002.
- [17] Hiroyuki Okazaki, Kenichi Arai, and Yasunari Shidama. Normal subgroup of product of groups. *Formalized Mathematics*, 19(1):23–26, 2011. doi:10.2478/v10037-011-0004-7.
- [18] D. Robinson. *A Course in the Theory of Groups*. Springer New York, 2012.
- [19] J.J. Rotman. *An Introduction to the Theory of Groups*. Springer, 1995.
- [20] Andrzej Trybulec. Domains and their Cartesian products. *Formalized Mathematics*, 1(1):115–122, 1990.
- [21] Wojciech A. Trybulec. Non-contiguous substrings and one-to-one finite sequences. *Formalized Mathematics*, 1(3):569–573, 1990.
- [22] Wojciech A. Trybulec. Groups. *Formalized Mathematics*, 1(5):821–827, 1990.
- [23] Wojciech A. Trybulec. Subgroup and cosets of subgroups. *Formalized Mathematics*, 1(5):855–864, 1990.
- [24] Wojciech A. Trybulec. Classes of conjugation. Normal subgroups. *Formalized Mathematics*, 1(5):955–962, 1990.
- [25] Wojciech A. Trybulec. Lattice of subgroups of a group. Frattini subgroup. *Formalized Mathematics*, 2(1):41–47, 1991.

- [26] Wojciech A. Trybulec and Michał J. Trybulec. Homomorphisms and isomorphisms of groups. Quotient group. *Formalized Mathematics*, 2(4):573–578, 1991.
- [27] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [28] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [29] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.

*Received February 26, 2015*

---