

# Groups – Additive Notation

Roland Coghetto  
Rue de la Brasserie 5  
7100 La Louvière, Belgium

**Summary.** We translate the articles covering group theory already available in the Mizar Mathematical Library from multiplicative into additive notation. We adapt the works of Wojciech A. Trybulec [41, 42, 43] and Artur Korniłowicz [25].

In particular, these authors have defined the notions of group, abelian group, power of an element of a group, order of a group and order of an element, subgroup, coset of a subgroup, index of a subgroup, conjugation, normal subgroup, topological group, dense subset and basis of a topological group. Lagrange’s theorem and some other theorems concerning these notions [9, 24, 22] are presented.

Note that “The term  $\mathbb{Z}$ -module is simply another name for an additive abelian group” [27]. We take an approach different than that used by Futa et al. [21] to use in a future article the results obtained by Artur Korniłowicz [25]. Indeed, Hölzl et al. showed that it was possible to build “a generic theory of limits based on filters” in Isabelle/HOL [23, 10]. Our goal is to define the convergence of a sequence and the convergence of a series in an abelian topological group [11] using the notion of filters.

MSC: 20A05 20K00 03B35

Keywords: additive group; subgroup; Lagrange theorem; conjugation; normal subgroup; index; additive topological group; basis; neighborhood; additive abelian group;  $\mathbb{Z}$ -module

MML identifier: GROUP\_1A, version: 8.1.04 5.32.1240

The notation and terminology used in this paper have been introduced in the following articles: [12], [32], [31], [2], [18], [28], [33], [13], [19], [39], [14], [15], [1], [40], [26], [35], [36], [5], [6], [16], [30], [8], [46], [47], [44], [29], [37], [45], [25], [48], [20], [7], [38], and [17].

## 1. ADDITIVE NOTATION FOR GROUPS – GROUP\_1

From now on  $m, n$  denote natural numbers,  $i, j$  denote integers,  $S$  denotes a non empty additive magma, and  $r, r_1, r_2, s, s_1, s_2, t, t_1, t_2$  denote elements of  $S$ .

The scheme *SeqEx2Dbis* deals with non empty sets  $\mathcal{X}, \mathcal{Z}$  and a ternary predicate  $\mathcal{P}$  and states that

(Sch. 1) There exists a function  $f$  from  $\mathbb{N} \times \mathcal{X}$  into  $\mathcal{Z}$  such that for every natural number  $x$  for every element  $y$  of  $\mathcal{X}$ ,  $\mathcal{P}[x, y, f(x, y)]$

provided

- for every natural number  $x$  and for every element  $y$  of  $\mathcal{X}$ , there exists an element  $z$  of  $\mathcal{Z}$  such that  $\mathcal{P}[x, y, z]$ .

Let  $I_1$  be an additive magma. We say that  $I_1$  is add-unital if and only if

(Def. 1) there exists an element  $e$  of  $I_1$  such that for every element  $h$  of  $I_1$ ,  $h + e = h$  and  $e + h = h$ .

We say that  $I_1$  is additive group-like if and only if

(Def. 2) there exists an element  $e$  of  $I_1$  such that for every element  $h$  of  $I_1$ ,  $h + e = h$  and  $e + h = h$  and there exists an element  $g$  of  $I_1$  such that  $h + g = e$  and  $g + h = e$ .

Let us note that every additive magma which is additive group-like is also add-unital and there exists an additive magma which is strict, additive group-like, add-associative, and non empty.

An additive group is an additive group-like, add-associative, non empty additive magma. Now we state the propositions:

- (1) Suppose for every  $r, s$ , and  $t$ ,  $(r + s) + t = r + (s + t)$  and there exists  $t$  such that for every  $s_1$ ,  $s_1 + t = s_1$  and  $t + s_1 = s_1$  and there exists  $s_2$  such that  $s_1 + s_2 = t$  and  $s_2 + s_1 = t$ . Then  $S$  is an additive group.
- (2) Suppose for every  $r, s$ , and  $t$ ,  $(r + s) + t = r + (s + t)$  and for every  $r$  and  $s$ , there exists  $t$  such that  $r + t = s$  and there exists  $t$  such that  $t + r = s$ . Then  $S$  is add-associative and additive group-like.
- (3)  $\langle \mathbb{R}, +_{\mathbb{R}} \rangle$  is add-associative and additive group-like.

From now on  $G$  denotes an additive group-like, non empty additive magma and  $e, h$  denote elements of  $G$ .

Let  $G$  be an additive magma. Assume  $G$  is add-unital. The functor  $0_G$  yielding an element of  $G$  is defined by

(Def. 3) for every element  $h$  of  $G$ ,  $h + 0_G = h$  and  $0_G + h = h$ .

Now we state the proposition:

(4) If for every  $h$ ,  $h + e = h$  and  $e + h = h$ , then  $e = 0_G$ .

From now on  $G$  denotes an additive group and  $f, g, h$  denote elements of  $G$ .

Let us consider  $G$  and  $h$ . The functor  $-h$  yielding an element of  $G$  is defined

by

(Def. 4)  $h + it = 0_G$  and  $it + h = 0_G$ .

Let us note that the functor is involutive.

Now we state the propositions:

(5) If  $h + g = 0_G$  and  $g + h = 0_G$ , then  $g = -h$ .

(6) If  $h + g = h + f$  or  $g + h = f + h$ , then  $g = f$ .

(7) If  $h + g = h$  or  $g + h = h$ , then  $g = 0_G$ . The theorem is a consequence of (6).

(8)  $-0_G = 0_G$ .

(9) If  $-h = -g$ , then  $h = g$ . The theorem is a consequence of (6).

(10) If  $-h = 0_G$ , then  $h = 0_G$ . The theorem is a consequence of (8).

(11) If  $h + g = 0_G$ , then  $h = -g$  and  $g = -h$ . The theorem is a consequence of (6).

(12)  $h + f = g$  if and only if  $f = -h + g$ . The theorem is a consequence of (6).

(13)  $f + h = g$  if and only if  $f = g + -h$ . The theorem is a consequence of (6).

(14) There exists  $f$  such that  $g + f = h$ . The theorem is a consequence of (12).

(15) There exists  $f$  such that  $f + g = h$ . The theorem is a consequence of (13).

(16)  $-(h + g) = -g + -h$ . The theorem is a consequence of (11).

(17)  $g + h = h + g$  if and only if  $-(g + h) = -g + -h$ . The theorem is a consequence of (16) and (6).

(18)  $g + h = h + g$  if and only if  $-g + -h = -h + -g$ . The theorem is a consequence of (16) and (17).

(19)  $g + h = h + g$  if and only if  $g + -h = -h + g$ . The theorem is a consequence of (18), (11), and (16).

From now on  $u$  denotes a unary operation on  $G$ .

Let us consider  $G$ . The functor add inverse  $G$  yielding a unary operation on  $G$  is defined by

(Def. 5)  $it(h) = -h$ .

Let  $G$  be an add-associative, non empty additive magma. Let us note that the addition of  $G$  is associative.

Let us consider an add-unital, non empty additive magma  $G$ . Now we state the propositions:

(20)  $0_G$  is a unity w.r.t. the addition of  $G$ .

(21)  $1_\alpha = 0_G$ , where  $\alpha$  is the addition of  $G$ . The theorem is a consequence of (20).

Let  $G$  be an add-unital, non empty additive magma. Let us note that the addition of  $G$  is unital.

Now we state the proposition:

(22) add inverse  $G$  is an inverse operation w.r.t. the addition of  $G$ . The theorem is a consequence of (21).

Let us consider  $G$ . One can verify that the addition of  $G$  has inverse operation.

Now we state the proposition:

(23) The inverse operation w.r.t. the addition of  $G = \text{add inverse } G$ . The theorem is a consequence of (22).

Let  $G$  be a non empty additive magma. The functor  $\text{mult } G$  yielding a function from  $\mathbb{N} \times (\text{the carrier of } G)$  into the carrier of  $G$  is defined by

(Def. 6) for every element  $h$  of  $G$ ,  $it(0, h) = 0_G$  and for every natural number  $n$ ,  
 $it(n + 1, h) = it(n, h) + h$ .

Let us consider  $G$ ,  $i$ , and  $h$ . The functor  $i \cdot h$  yielding an element of  $G$  is defined by the term

(Def. 7) 
$$\begin{cases} (\text{mult } G)(|i|, h), & \text{if } 0 \leq i, \\ -(\text{mult } G)(|i|, h), & \text{otherwise.} \end{cases}$$

Let us consider  $n$ . One can check that the functor  $n \cdot h$  is defined by the term

(Def. 8)  $(\text{mult } G)(n, h)$ .

Now we state the propositions:

(24)  $0 \cdot h = 0_G$ .

(25)  $1 \cdot h = h$ .

(26)  $2 \cdot h = h + h$ . The theorem is a consequence of (25).

(27)  $3 \cdot h = h + h + h$ . The theorem is a consequence of (26).

(28)  $2 \cdot h = 0_G$  if and only if  $-h = h$ . The theorem is a consequence of (26) and (11).

(29) If  $i \leq 0$ , then  $i \cdot h = -|i| \cdot h$ . The theorem is a consequence of (8).

(30)  $i \cdot 0_G = 0_G$ . The theorem is a consequence of (8).

(31)  $(-1) \cdot h = -h$ . The theorem is a consequence of (25).

(32)  $(i + j) \cdot h = i \cdot h + j \cdot h$ .

PROOF: Define  $\mathcal{P}[\text{integer}] \equiv$  for every  $i$ ,  $(i + \$_1) \cdot h = i \cdot h + \$_1 \cdot h$ . For every  $j$  such that  $\mathcal{P}[j]$  holds  $\mathcal{P}[j - 1]$  and  $\mathcal{P}[j + 1]$ .  $\mathcal{P}[0]$ . For every  $j$ ,  $\mathcal{P}[j]$  from [40, Sch. 4].  $\square$

(33) (i)  $(i + 1) \cdot h = i \cdot h + h$ , and

(ii)  $(i + 1) \cdot h = h + i \cdot h$ .

The theorem is a consequence of (25) and (32).

(34)  $(-i) \cdot h = -i \cdot h$ .

Let us assume that  $g + h = h + g$ . Now we state the propositions:

(35)  $i \cdot (g + h) = i \cdot g + i \cdot h$ . The theorem is a consequence of (16).

(36)  $i \cdot g + j \cdot h = j \cdot h + i \cdot g$ . The theorem is a consequence of (19) and (16).

(37)  $g + i \cdot h = i \cdot h + g$ . The theorem is a consequence of (25) and (36).

Let us consider  $G$  and  $h$ . We say that  $h$  is of order 0 if and only if

(Def. 9) if  $n \cdot h = 0_G$ , then  $n = 0$ .

One can check that  $0_G$  is non of order 0.

Let us consider  $h$ . The functor  $\text{ord}(h)$  yielding an element of  $\mathbb{N}$  is defined by

(Def. 10) (i)  $it = 0$ , if  $h$  is of order 0,

(ii)  $it \cdot h = 0_G$  and  $it \neq 0$  and for every  $m$  such that  $m \cdot h = 0_G$  and  $m \neq 0$  holds  $it \leq m$ , **otherwise**.

Now we state the propositions:

(38)  $\text{ord}(h) \cdot h = 0_G$ .

(39)  $\text{ord}(0_G) = 1$ .

(40) If  $\text{ord}(h) = 1$ , then  $h = 0_G$ . The theorem is a consequence of (25).

Observe that there exists an additive group which is strict and Abelian.

Now we state the proposition:

(41)  $\langle \mathbb{R}, +_{\mathbb{R}} \rangle$  is an Abelian additive group. The theorem is a consequence of (3).

In the sequel  $A$  denotes an Abelian additive group and  $a, b$  denote elements of  $A$ .

Now we state the propositions:

(42)  $-(a + b) = -a + -b$ .

(43)  $i \cdot (a + b) = i \cdot a + i \cdot b$ .

(44)  $\langle$ the carrier of  $A$ , the addition of  $A, 0_A \rangle$  is Abelian, add-associative, right zeroed, and right complementable.

Let us consider an add-unital, non empty additive magma  $L$  and an element  $x$  of  $L$ . Now we state the propositions:

(45)  $(\text{mult } L)(1, x) = x$ .

(46)  $(\text{mult } L)(2, x) = x + x$ . The theorem is a consequence of (45).

Now we state the proposition:

(47) Let us consider an add-associative, Abelian, add-unital, non empty additive magma  $L$ , elements  $x, y$  of  $L$ , and a natural number  $n$ . Then  $(\text{mult } L)(n, x + y) = (\text{mult } L)(n, x) + (\text{mult } L)(n, y)$ .

PROOF: Define  $\mathcal{P}[\text{natural number}] \equiv (\text{mult } L)(\$_1, x+y) = (\text{mult } L)(\$_1, x) + (\text{mult } L)(\$_1, y)$ . For every natural number  $n$ ,  $\mathcal{P}[n]$  from [5, Sch. 2].  $\square$

Let  $G, H$  be additive magmas and  $I_1$  be a function from  $G$  into  $H$ . We say that  $I_1$  preserves zero if and only if

(Def. 11)  $I_1(0_G) = 0_H$ .

## 2. SUBGROUPS AND LAGRANGE THEOREM – GROUP\_2

In the sequel  $x$  denotes an object,  $y, y_1, y_2, Y, Z$  denote sets,  $k$  denotes a natural number,  $G$  denotes an additive group,  $a, g, h$  denote elements of  $G$ , and  $A$  denotes a subset of  $G$ .

Let us consider  $G$  and  $A$ . The functor  $-A$  yielding a subset of  $G$  is defined by the term

(Def. 12)  $\{-g : g \in A\}$ .

One can check that the functor is involutive.

Now we state the propositions:

(48)  $x \in -A$  if and only if there exists  $g$  such that  $x = -g$  and  $g \in A$ .

(49)  $-\{g\} = \{-g\}$ .

(50)  $-\{g, h\} = \{-g, -h\}$ .

(51)  $-\emptyset_\alpha = \emptyset$ , where  $\alpha$  is the carrier of  $G$ .

(52)  $-\Omega_\alpha =$  the carrier of  $G$ , where  $\alpha$  is the carrier of  $G$ .

(53)  $A \neq \emptyset$  if and only if  $-A \neq \emptyset$ . The theorem is a consequence of (48).

Let us consider  $G$ . Let  $A$  be an empty subset of  $G$ . Observe that  $-A$  is empty.

Let  $A$  be a non empty subset of  $G$ . One can check that  $-A$  is non empty.

In the sequel  $G$  denotes a non empty additive magma,  $A, B, C$  denote subsets of  $G$ , and  $a, b, g, g_1, g_2, h, h_1, h_2$  denote elements of  $G$ .

Let  $G$  be an Abelian, non empty additive magma and  $A, B$  be subsets of  $G$ . One can check that the functor  $A + B$  is commutative.

(54)  $x \in A + B$  if and only if there exists  $g$  and there exists  $h$  such that  $x = g + h$  and  $g \in A$  and  $h \in B$ .

(55)  $A \neq \emptyset$  and  $B \neq \emptyset$  if and only if  $A + B \neq \emptyset$ . The theorem is a consequence of (54).

(56) If  $G$  is add-associative, then  $(A + B) + C = A + (B + C)$ .

(57) Let us consider an additive group  $G$ , and subsets  $A, B$  of  $G$ . Then  $-(A + B) = -B + -A$ . The theorem is a consequence of (16).

(58)  $A + (B \cup C) = A + B \cup (A + C)$ .

(59)  $(A \cup B) + C = A + C \cup (B + C)$ .

(60)  $A + B \cap C \subseteq (A + B) \cap (A + C)$ .

(61)  $A \cap B + C \subseteq (A + C) \cap (B + C)$ .

(62) (i)  $\emptyset_\alpha + A = \emptyset$ , and

(ii)  $A + \emptyset_\alpha = \emptyset$ ,

where  $\alpha$  is the carrier of  $G$ . The theorem is a consequence of (54).

(63) Let us consider an additive group  $G$ , and a subset  $A$  of  $G$ . Suppose  $A \neq \emptyset$ . Then

(i)  $\Omega_\alpha + A =$  the carrier of  $G$ , and

(ii)  $A + \Omega_\alpha =$  the carrier of  $G$ ,

where  $\alpha$  is the carrier of  $G$ .

(64)  $\{g\} + \{h\} = \{g + h\}$ .

(65)  $\{g\} + \{g_1, g_2\} = \{g + g_1, g + g_2\}$ .

(66)  $\{g_1, g_2\} + \{g\} = \{g_1 + g, g_2 + g\}$ .

(67)  $\{g, h\} + \{g_1, g_2\} = \{g + g_1, g + g_2, h + g_1, h + g_2\}$ .

Let us consider an additive group  $G$  and a subset  $A$  of  $G$ . Now we state the propositions:

(68) Suppose for every elements  $g_1, g_2$  of  $G$  such that  $g_1, g_2 \in A$  holds  $g_1 + g_2 \in A$  and for every element  $g$  of  $G$  such that  $g \in A$  holds  $-g \in A$ . Then  $A + A = A$ .

(69) If for every element  $g$  of  $G$  such that  $g \in A$  holds  $-g \in A$ , then  $-A = A$ .

(70) If for every  $a$  and  $b$  such that  $a \in A$  and  $b \in B$  holds  $a + b = b + a$ , then  $A + B = B + A$ .

(71) If  $G$  is an Abelian additive group, then  $A + B = B + A$ .

(72) Let us consider an Abelian additive group  $G$ , and subsets  $A, B$  of  $G$ . Then  $-(A + B) = -A + -B$ . The theorem is a consequence of (42).

Let us consider  $G, g$ , and  $A$ . The functors:  $g + A$  and  $A + g$  yielding subsets of  $G$  are defined by terms,

(Def. 13)  $\{g\} + A$ ,

(Def. 14)  $A + \{g\}$ ,

respectively. Now we state the propositions:

(73)  $x \in g + A$  if and only if there exists  $h$  such that  $x = g + h$  and  $h \in A$ .

(74)  $x \in A + g$  if and only if there exists  $h$  such that  $x = h + g$  and  $h \in A$ .

Let us assume that  $G$  is add-associative. Now we state the propositions:

(75)  $(g + A) + B = g + (A + B)$ .

(76)  $(A + g) + B = A + (g + B)$ .

(77)  $(A + B) + g = A + (B + g)$ .

(78)  $(g + h) + A = g + (h + A)$ . The theorem is a consequence of (64) and (56).

(79)  $(g + A) + h = g + (A + h)$ .

(80)  $(A + g) + h = A + (g + h)$ . The theorem is a consequence of (56) and (64).

(81) (i)  $\emptyset_\alpha + a = \emptyset$ , and

(ii)  $a + \emptyset_\alpha = \emptyset$ ,

where  $\alpha$  is the carrier of  $G$ .

From now on  $G$  denotes an additive group-like, non empty additive magma,  $h, g, g_1, g_2$  denote elements of  $G$ , and  $A$  denotes a subset of  $G$ .

(82) Let us consider an additive group  $G$ , and an element  $a$  of  $G$ . Then

(i)  $\Omega_\alpha + a =$  the carrier of  $G$ , and

(ii)  $a + \Omega_\alpha =$  the carrier of  $G$ ,

where  $\alpha$  is the carrier of  $G$ .

(83) (i)  $0_G + A = A$ , and

(ii)  $A + 0_G = A$ .

The theorem is a consequence of (73) and (74).

(84) If  $G$  is an Abelian additive group, then  $g + A = A + g$ .

Let  $G$  be an additive group-like, non empty additive magma.

A subgroup of  $G$  is an additive group-like, non empty additive magma and is defined by

(Def. 15) the carrier of  $it \subseteq$  the carrier of  $G$  and the addition of  $it =$  (the addition of  $G$ )  $\upharpoonright$  (the carrier of  $it$ ).

In the sequel  $H$  denotes a subgroup of  $G$  and  $h, h_1, h_2$  denote elements of  $H$ .

Now we state the propositions:

(85) If  $G$  is finite, then  $H$  is finite.

(86) If  $x \in H$ , then  $x \in G$ .

(87)  $h \in G$ .

(88)  $h$  is an element of  $G$ .

(89) If  $h_1 = g_1$  and  $h_2 = g_2$ , then  $h_1 + h_2 = g_1 + g_2$ .



Let  $G$  be an additive group. Let us observe that every subgroup of  $G$  is add-associative.

In the sequel  $G, G_1, G_2, G_3$  denote additive groups,  $a, a_1, a_2, b, b_1, b_2, g, g_1, g_2$  denote elements of  $G$ ,  $A, B$  denote subsets of  $G$ ,  $H, H_1, H_2, H_3$  denote subgroups of  $G$ , and  $h, h_1, h_2$  denote elements of  $H$ .

- (90)  $0_H = 0_G$ . The theorem is a consequence of (87), (89), and (7).
- (91)  $0_{H_1} = 0_{H_2}$ . The theorem is a consequence of (90).
- (92)  $0_G \in H$ . The theorem is a consequence of (90).
- (93)  $0_{H_1} \in H_2$ . The theorem is a consequence of (90) and (92).
- (94) If  $h = g$ , then  $-h = -g$ . The theorem is a consequence of (87), (89), (90), and (11).
- (95) add inverse  $H =$  add inverse  $G \upharpoonright$  (the carrier of  $H$ ). The theorem is a consequence of (87) and (94).
- (96) If  $g_1, g_2 \in H$ , then  $g_1 + g_2 \in H$ . The theorem is a consequence of (89).
- (97) If  $g \in H$ , then  $-g \in H$ . The theorem is a consequence of (94).

Let us consider  $G$ . Observe that there exists a subgroup of  $G$  which is strict.

- (98) Suppose  $A \neq \emptyset$  and for every  $g_1$  and  $g_2$  such that  $g_1, g_2 \in A$  holds  $g_1 + g_2 \in A$  and for every  $g$  such that  $g \in A$  holds  $-g \in A$ . Then there exists a strict subgroup  $H$  of  $G$  such that the carrier of  $H = A$ .

PROOF: Reconsider  $D = A$  as a non empty set. Set  $o =$  (the addition of  $G$ )  $\upharpoonright A$ .  $\text{rng } o \subseteq A$  by [17, (87)], [14, (47)]. Set  $H = \langle D, o \rangle$ .  $H$  is additive group-like.  $\square$

- (99) If  $G$  is an Abelian additive group, then  $H$  is Abelian. The theorem is a consequence of (87) and (89).

Let  $G$  be an Abelian additive group. One can check that every subgroup of  $G$  is Abelian.

- (100)  $G$  is a subgroup of  $G$ .
- (101) Suppose  $G_1$  is a subgroup of  $G_2$  and  $G_2$  is a subgroup of  $G_1$ . Then the additive magma of  $G_1 =$  the additive magma of  $G_2$ .
- (102) If  $G_1$  is a subgroup of  $G_2$  and  $G_2$  is a subgroup of  $G_3$ , then  $G_1$  is a subgroup of  $G_3$ .
- (103) If the carrier of  $H_1 \subseteq$  the carrier of  $H_2$ , then  $H_1$  is a subgroup of  $H_2$ .
- (104) If for every  $g$  such that  $g \in H_1$  holds  $g \in H_2$ , then  $H_1$  is a subgroup of  $H_2$ . The theorem is a consequence of (87) and (103).
- (105) Suppose the carrier of  $H_1 =$  the carrier of  $H_2$ . Then the additive magma of  $H_1 =$  the additive magma of  $H_2$ . The theorem is a consequence of (103) and (101).

(106) Suppose for every  $g$ ,  $g \in H_1$  iff  $g \in H_2$ . Then the additive magma of  $H_1 =$  the additive magma of  $H_2$ . The theorem is a consequence of (104) and (101).

Let us consider  $G$ . Let  $H_1, H_2$  be strict subgroups of  $G$ . One can check that  $H_1 = H_2$  if and only if the condition (Def. 16) is satisfied.

(Def. 16) for every  $g$ ,  $g \in H_1$  iff  $g \in H_2$ .

Now we state the propositions:

(107) Let us consider an additive group  $G$ , and a subgroup  $H$  of  $G$ . Suppose the carrier of  $G \subseteq$  the carrier of  $H$ . Then the additive magma of  $H =$  the additive magma of  $G$ . The theorem is a consequence of (100) and (105).

(108) Suppose for every element  $g$  of  $G$ ,  $g \in H$ . Then the additive magma of  $H =$  the additive magma of  $G$ . The theorem is a consequence of (100) and (106).

Let us consider  $G$ . The functor  $\mathbf{0}_G$  yielding a strict subgroup of  $G$  is defined by

(Def. 17) the carrier of  $it = \{0_G\}$ .

The functor  $\Omega_G$  yielding a strict subgroup of  $G$  is defined by the term

(Def. 18) the additive magma of  $G$ .

Note that the functor is projective.

Now we state the propositions:

(109)  $\mathbf{0}_H = \mathbf{0}_G$ . The theorem is a consequence of (90) and (102).

(110)  $\mathbf{0}_{H_1} = \mathbf{0}_{H_2}$ . The theorem is a consequence of (109).

(111)  $\mathbf{0}_G$  is a subgroup of  $H$ . The theorem is a consequence of (109).

(112) Let us consider a strict additive group  $G$ . Then every subgroup of  $G$  is a subgroup of  $\Omega_G$ .

(113) Every strict additive group is a subgroup of  $\Omega_G$ .

(114)  $\mathbf{0}_G$  is finite.

Let us consider  $G$ . Note that  $\mathbf{0}_G$  is finite and there exists a subgroup of  $G$  which is strict and finite and there exists an additive group which is strict and finite.

Let  $G$  be a finite additive group. One can verify that every subgroup of  $G$  is finite.

Now we state the propositions:

(115)  $\overline{\mathbf{0}_G} = 1$ .

(116) Let us consider a strict, finite subgroup  $H$  of  $G$ . If  $\overline{H} = 1$ , then  $H = \mathbf{0}_G$ . The theorem is a consequence of (92).

(117)  $\overline{\overline{H}} \subseteq \overline{\overline{G}}$ .

Let us consider a finite additive group  $G$  and a subgroup  $H$  of  $G$ . Now we state the propositions:

(118)  $\overline{\overline{H}} \leq \overline{\overline{G}}$ .

(119) Suppose  $\overline{\overline{G}} = \overline{\overline{H}}$ . Then the additive magma of  $H =$  the additive magma of  $G$ .

PROOF: The carrier of  $H =$  the carrier of  $G$  by [3, (48)].  $\square$

Let us consider  $G$  and  $H$ . The functor  $\overline{\overline{H}}$  yielding a subset of  $G$  is defined by the term

(Def. 19) the carrier of  $H$ .

Now we state the propositions:

(120) If  $g_1, g_2 \in \overline{\overline{H}}$ , then  $g_1 + g_2 \in \overline{\overline{H}}$ . The theorem is a consequence of (96).

(121) If  $g \in \overline{\overline{H}}$ , then  $-g \in \overline{\overline{H}}$ . The theorem is a consequence of (97).

(122)  $\overline{\overline{H}} + \overline{\overline{H}} = \overline{\overline{H}}$ . The theorem is a consequence of (121), (120), and (68).

(123)  $-\overline{\overline{H}} = \overline{\overline{H}}$ . The theorem is a consequence of (121) and (69).

(124) (i) if  $\overline{\overline{H_1}} + \overline{\overline{H_2}} = \overline{\overline{H_2}} + \overline{\overline{H_1}}$ , then there exists a strict subgroup  $H$  of  $G$  such that the carrier of  $H = \overline{\overline{H_1}} + \overline{\overline{H_2}}$ , and

(ii) if there exists  $H$  such that the carrier of  $H = \overline{\overline{H_1}} + \overline{\overline{H_2}}$ , then  $\overline{\overline{H_1}} + \overline{\overline{H_2}} = \overline{\overline{H_2}} + \overline{\overline{H_1}}$ .

The theorem is a consequence of (121), (16), (120), (55), and (98).

(125) Suppose  $G$  is an Abelian additive group. Then there exists a strict subgroup  $H$  of  $G$  such that the carrier of  $H = \overline{\overline{H_1}} + \overline{\overline{H_2}}$ . The theorem is a consequence of (71) and (124).

Let us consider  $G, H_1,$  and  $H_2$ . The functor  $H_1 \cap H_2$  yielding a strict subgroup of  $G$  is defined by

(Def. 20) the carrier of  $it = \overline{\overline{H_1}} \cap \overline{\overline{H_2}}$ .

Now we state the propositions:

(126) (i) for every subgroup  $H$  of  $G$  such that  $H = H_1 \cap H_2$  holds the carrier of  $H =$  (the carrier of  $H_1$ )  $\cap$  (the carrier of  $H_2$ ), and

(ii) for every strict subgroup  $H$  of  $G$  such that the carrier of  $H =$  (the carrier of  $H_1$ )  $\cap$  (the carrier of  $H_2$ ) holds  $H = H_1 \cap H_2$ .

(127)  $\overline{\overline{H_1 \cap H_2}} = \overline{\overline{H_1}} \cap \overline{\overline{H_2}}$ .

(128)  $x \in H_1 \cap H_2$  if and only if  $x \in H_1$  and  $x \in H_2$ .

(129) Let us consider a strict subgroup  $H$  of  $G$ . Then  $H \cap H = H$ . The theorem is a consequence of (105).

Let us consider  $G, H_1,$  and  $H_2$ . Note that the functor  $H_1 \cap H_2$  is commutative.

(130)  $(H_1 \cap H_2) \cap H_3 = H_1 \cap (H_2 \cap H_3)$ . The theorem is a consequence of (105).

(131) (i)  $\mathbf{0}_G \cap H = \mathbf{0}_G$ , and

(ii)  $H \cap \mathbf{0}_G = \mathbf{0}_G$ .

The theorem is a consequence of (111).

(132) Let us consider a strict additive group  $G$ , and a strict subgroup  $H$  of  $G$ . Then

(i)  $H \cap \Omega_G = H$ , and

(ii)  $\Omega_G \cap H = H$ .

(133) Let us consider a strict additive group  $G$ . Then  $\Omega_G \cap \Omega_G = G$ .

(134)  $H_1 \cap H_2$  is subgroup of  $H_1$  and subgroup of  $H_2$ .

(135) Let us consider a subgroup  $H_1$  of  $G$ . Then  $H_1$  is a subgroup of  $H_2$  if and only if the additive magma of  $H_1 \cap H_2 =$  the additive magma of  $H_1$ .

(136) If  $H_1$  is a subgroup of  $H_2$ , then  $H_1 \cap H_3$  is a subgroup of  $H_2$ . The theorem is a consequence of (102).

(137) If  $H_1$  is subgroup of  $H_2$  and subgroup of  $H_3$ , then  $H_1$  is a subgroup of  $H_2 \cap H_3$ . The theorem is a consequence of (86), (128), and (104).

(138) If  $H_1$  is a subgroup of  $H_2$ , then  $H_1 \cap H_3$  is a subgroup of  $H_2 \cap H_3$ . The theorem is a consequence of (126) and (103).

(139) If  $H_1$  is finite or  $H_2$  is finite, then  $H_1 \cap H_2$  is finite.

Let us consider  $G$ ,  $H$ , and  $A$ . The functors:  $A + H$  and  $H + A$  yielding subsets of  $G$  are defined by terms,

(Def. 21)  $A + \overline{H}$ ,

(Def. 22)  $\overline{H} + A$ ,

respectively. Now we state the propositions:

(140)  $x \in A + H$  if and only if there exists  $g_1$  and there exists  $g_2$  such that  $x = g_1 + g_2$  and  $g_1 \in A$  and  $g_2 \in H$ .

(141)  $x \in H + A$  if and only if there exists  $g_1$  and there exists  $g_2$  such that  $x = g_1 + g_2$  and  $g_1 \in H$  and  $g_2 \in A$ .

(142)  $(A + B) + H = A + (B + H)$ .

(143)  $(A + H) + B = A + (H + B)$ .

(144)  $(H + A) + B = H + (A + B)$ .

(145)  $(A + H_1) + H_2 = A + (H_1 + \overline{H_2})$ .

(146)  $(H_1 + A) + H_2 = H_1 + (A + H_2)$ .

(147)  $(H_1 + \overline{H_2}) + A = H_1 + (H_2 + A)$ .

(148) If  $G$  is an Abelian additive group, then  $A + H = H + A$ .

Let us consider  $G$ ,  $H$ , and  $a$ . The functors:  $a + H$  and  $H + a$  yielding subsets of  $G$  are defined by terms,

(Def. 23)  $a + \overline{H}$ ,

(Def. 24)  $\overline{H} + a$ ,

respectively. Now we state the propositions:

(149)  $x \in a + H$  if and only if there exists  $g$  such that  $x = a + g$  and  $g \in H$ .

The theorem is a consequence of (73).

(150)  $x \in H + a$  if and only if there exists  $g$  such that  $x = g + a$  and  $g \in H$ .

The theorem is a consequence of (74).

(151)  $(a + b) + H = a + (b + H)$ .

(152)  $(a + H) + b = a + (H + b)$ .

(153)  $(H + a) + b = H + (a + b)$ .

(154) (i)  $a \in a + H$ , and

(ii)  $a \in H + a$ .

The theorem is a consequence of (92), (149), and (150).

(155) (i)  $0_G + H = \overline{H}$ , and

(ii)  $H + 0_G = \overline{H}$ .

(156) (i)  $\mathbf{0}_G + a = \{a\}$ , and

(ii)  $a + \mathbf{0}_G = \{a\}$ .

The theorem is a consequence of (64).

(157) (i)  $a + \Omega_G =$  the carrier of  $G$ , and

(ii)  $\Omega_G + a =$  the carrier of  $G$ .

The theorem is a consequence of (63).

(158) If  $G$  is an Abelian additive group, then  $a + H = H + a$ .

(159)  $a \in H$  if and only if  $a + H = \overline{H}$ . The theorem is a consequence of (149), (96), (97), and (92).

(160)  $a + H = b + H$  if and only if  $-b + a \in H$ . The theorem is a consequence of (78), (83), and (159).

(161)  $a + H = b + H$  if and only if  $a + H$  meets  $b + H$ . The theorem is a consequence of (154), (149), (97), (13), (12), (96), and (160).

(162)  $(a + b) + H \subseteq a + H + (b + H)$ . The theorem is a consequence of (149) and (92).

(163) (i)  $\overline{H} \subseteq a + H + (-a + H)$ , and

(ii)  $\overline{H} \subseteq -a + H + (a + H)$ .

The theorem is a consequence of (83) and (162).

(164)  $2 \cdot a + H \subseteq a + H + (a + H)$ . The theorem is a consequence of (26) and (162).

(165)  $a \in H$  if and only if  $H + a = \overline{H}$ . The theorem is a consequence of (150), (96), (97), and (92).

(166)  $H + a = H + b$  if and only if  $b + -a \in H$ . The theorem is a consequence of (83), (80), and (165).

(167)  $H + a = H + b$  if and only if  $H + a$  meets  $H + b$ . The theorem is a consequence of (154), (150), (97), (12), (13), (96), and (166).

(168)  $(H + a) + b \subseteq H + a + (H + b)$ . The theorem is a consequence of (92), (150), and (80).

(169) (i)  $\overline{H} \subseteq H + a + (H + -a)$ , and

(ii)  $\overline{H} \subseteq H + -a + (H + a)$ .

The theorem is a consequence of (80), (83), and (168).

(170)  $H + 2 \cdot a \subseteq H + a + (H + a)$ . The theorem is a consequence of (80), (26), and (168).

(171)  $a + H_1 \cap H_2 = (a + H_1) \cap (a + H_2)$ . The theorem is a consequence of (149), (128), and (6).

(172)  $H_1 \cap H_2 + a = (H_1 + a) \cap (H_2 + a)$ . The theorem is a consequence of (150), (128), and (6).

(173) There exists a strict subgroup  $H_1$  of  $G$  such that the carrier of  $H_1 = a + H_2 + -a$ . The theorem is a consequence of (154), (74), (149), (97), (150), (16), (73), (56), (96), and (98).

(174)  $a + H \approx b + H$ .

PROOF: Define  $\mathcal{P}[\text{object}, \text{object}] \equiv$  there exists  $g_1$  such that  $\$1 = g_1$  and  $\$2 = b + -a + g_1$ . For every object  $x$  such that  $x \in a + H$  there exists an object  $y$  such that  $\mathcal{P}[x, y]$ . Consider  $f$  being a function such that  $\text{dom } f = a + H$  and for every object  $x$  such that  $x \in a + H$  holds  $\mathcal{P}[x, f(x)]$  from [4, Sch. 1].  $\text{rng } f = b + H$ .  $f$  is one-to-one.  $\square$

(175)  $a + H \approx H + b$ .

PROOF: Define  $\mathcal{P}[\text{object}, \text{object}] \equiv$  there exists  $g_1$  such that  $\$1 = g_1$  and  $\$2 = -a + g_1 + b$ . For every object  $x$  such that  $x \in a + H$  there exists an object  $y$  such that  $\mathcal{P}[x, y]$ . Consider  $f$  being a function such that  $\text{dom } f = a + H$  and for every object  $x$  such that  $x \in a + H$  holds  $\mathcal{P}[x, f(x)]$  from [4, Sch. 1].  $\text{rng } f = H + b$ .  $f$  is one-to-one.  $\square$

(176)  $H + a \approx H + b$ . The theorem is a consequence of (175).

(177) (i)  $\overline{H} \approx a + H$ , and

(ii)  $\overline{H} \approx H + a$ .

The theorem is a consequence of (83), (174), and (176).

- (178) (i)  $\overline{\overline{H}} = \overline{a + H}$ , and  
 (ii)  $\overline{\overline{H}} = \overline{H + a}$ .

(179) Let us consider a finite subgroup  $H$  of  $G$ . Then there exist finite sets  $B, C$  such that

- (i)  $B = a + H$ , and  
 (ii)  $C = H + a$ , and  
 (iii)  $\overline{\overline{H}} = \overline{\overline{B}}$ , and  
 (iv)  $\overline{\overline{H}} = \overline{\overline{C}}$ .

The theorem is a consequence of (177).

Let us consider  $G$  and  $H$ . The functors: the left cosets of  $H$  and the right cosets of  $H$  yielding families of subsets of  $G$  are defined by conditions,

(Def. 25)  $A \in$  the left cosets of  $H$  iff there exists  $a$  such that  $A = a + H$ ,

(Def. 26)  $A \in$  the right cosets of  $H$  iff there exists  $a$  such that  $A = H + a$ ,

respectively. Now we state the propositions:

(180) If  $G$  is finite, then the right cosets of  $H$  is finite and the left cosets of  $H$  is finite.

- (181) (i)  $\overline{\overline{H}} \in$  the left cosets of  $H$ , and  
 (ii)  $\overline{\overline{H}} \in$  the right cosets of  $H$ .

The theorem is a consequence of (83).

(182) The left cosets of  $H \approx$  the right cosets of  $H$ .

PROOF: Define  $\mathcal{P}[\text{object}, \text{object}] \equiv$  there exists  $g$  such that  $\$1 = g + H$  and  $\$2 = H + -g$ . For every object  $x$  such that  $x \in$  the left cosets of  $H$  there exists an object  $y$  such that  $\mathcal{P}[x, y]$ . Consider  $f$  being a function such that  $\text{dom } f =$  the left cosets of  $H$  and for every object  $x$  such that  $x \in$  the left cosets of  $H$  holds  $\mathcal{P}[x, f(x)]$  from [4, Sch. 1].  $\text{rng } f =$  the right cosets of  $H$ .  $f$  is one-to-one.  $\square$

- (183) (i)  $\bigcup(\text{the left cosets of } H) =$  the carrier of  $G$ , and  
 (ii)  $\bigcup(\text{the right cosets of } H) =$  the carrier of  $G$ .

The theorem is a consequence of (87), (149), and (150).

(184) The left cosets of  $\mathbf{0}_G =$  the set of all  $\{a\}$ . The theorem is a consequence of (156).

(185) The right cosets of  $\mathbf{0}_G =$  the set of all  $\{a\}$ . The theorem is a consequence of (156).

Let us consider a strict subgroup  $H$  of  $G$ . Now we state the propositions:

(186) If the left cosets of  $H =$  the set of all  $\{a\}$ , then  $H = \mathbf{0}_G$ . The theorem is a consequence of (87), (149), (92), and (6).

(187) If the right cosets of  $H =$  the set of all  $\{a\}$ , then  $H = \mathbf{0}_G$ . The theorem is a consequence of (87), (150), (92), and (6).

(188) (i) the left cosets of  $\Omega_G = \{\text{the carrier of } G\}$ , and

(ii) the right cosets of  $\Omega_G = \{\text{the carrier of } G\}$ .

The theorem is a consequence of (157).

Let us consider a strict additive group  $G$  and a strict subgroup  $H$  of  $G$ . Now we state the propositions:

(189) If the left cosets of  $H = \{\text{the carrier of } G\}$ , then  $H = G$ . The theorem is a consequence of (149), (6), and (108).

(190) If the right cosets of  $H = \{\text{the carrier of } G\}$ , then  $H = G$ . The theorem is a consequence of (150), (6), and (108).

Let us consider  $G$  and  $H$ . The functor  $|\bullet : H|$  yielding a cardinal number is defined by the term

(Def. 27)  $\overline{\alpha}$ , where  $\alpha$  is the left cosets of  $H$ .

Now we state the proposition:

(191) (i)  $|\bullet : H| = \overline{\alpha}$ , and

(ii)  $|\bullet : H| = \overline{\beta}$ ,

where  $\alpha$  is the left cosets of  $H$  and  $\beta$  is the right cosets of  $H$ .

Let us consider  $G$  and  $H$ . Assume the left cosets of  $H$  is finite. The functor  $|\bullet : H|_{\mathbb{N}}$  yielding an element of  $\mathbb{N}$  is defined by

(Def. 28) there exists a finite set  $B$  such that  $B =$  the left cosets of  $H$  and  $it = \overline{B}$ .

Now we state the proposition:

(192) Suppose the left cosets of  $H$  is finite. Then

(i) there exists a finite set  $B$  such that  $B =$  the left cosets of  $H$  and  $|\bullet : H|_{\mathbb{N}} = \overline{B}$ , and

(ii) there exists a finite set  $C$  such that  $C =$  the right cosets of  $H$  and  $|\bullet : H|_{\mathbb{N}} = \overline{C}$ .

The theorem is a consequence of (182).

Let us consider a finite additive group  $G$  and a subgroup  $H$  of  $G$ . Now we state the propositions:

(193) LAGRANGE THEOREM FOR ADDITIVE GROUPS:

$\overline{G} = \overline{H} \cdot |\bullet : H|_{\mathbb{N}}$ . The theorem is a consequence of (179), (174), (161), and (183).

(194)  $\overline{H} | \overline{G}$ . The theorem is a consequence of (193).



- (195) Let us consider a finite additive group  $G$ , subgroups  $I, H$  of  $G$ , and a subgroup  $J$  of  $H$ . Suppose  $I = J$ . Then  $|\bullet : I|_{\mathbb{N}} = |\bullet : J|_{\mathbb{N}} \cdot |\bullet : H|_{\mathbb{N}}$ . The theorem is a consequence of (193).
- (196)  $|\bullet : \Omega_G|_{\mathbb{N}} = 1$ . The theorem is a consequence of (188).
- (197) Let us consider a strict additive group  $G$ , and a strict subgroup  $H$  of  $G$ . Suppose the left cosets of  $H$  is finite and  $|\bullet : H|_{\mathbb{N}} = 1$ . Then  $H = G$ . The theorem is a consequence of (183) and (189).
- (198)  $|\bullet : \mathbf{0}_G| = \overline{G}$ .  
 PROOF: Define  $\mathcal{F}(\text{object}) = \{\$1\}$ . Consider  $f$  being a function such that  $\text{dom } f = \text{the carrier of } G$  and for every object  $x$  such that  $x \in \text{the carrier of } G$  holds  $f(x) = \mathcal{F}(x)$  from [14, Sch. 3].  $\text{rng } f = \text{the left cosets of } \mathbf{0}_G$ .  $f$  is one-to-one by [17, (3)].  $\square$
- (199) Let us consider a finite additive group  $G$ . Then  $|\bullet : \mathbf{0}_G|_{\mathbb{N}} = \overline{G}$ . The theorem is a consequence of (193) and (115).
- (200) Let us consider a finite additive group  $G$ , and a strict subgroup  $H$  of  $G$ . Suppose  $|\bullet : H|_{\mathbb{N}} = \overline{G}$ . Then  $H = \mathbf{0}_G$ . The theorem is a consequence of (193) and (116).
- (201) Let us consider a strict subgroup  $H$  of  $G$ . Suppose the left cosets of  $H$  is finite and  $|\bullet : H| = \overline{G}$ . Then
- (i)  $G$  is finite, and
  - (ii)  $H = \mathbf{0}_G$ .
- The theorem is a consequence of (200).

### 3. CLASSES OF CONJUGATION AND NORMAL SUBGROUPS – GROUP\_3

From now on  $x, y, y_1, y_2$  denote sets,  $G$  denotes an additive group,  $a, b, c, d, g, h$  denote elements of  $G$ ,  $A, B, C, D$  denote subsets of  $G$ ,  $H, H_1, H_2, H_3$  denote subgroups of  $G$ ,  $n$  denotes a natural number, and  $i$  denotes an integer.

Now we state the propositions:

- (202) (i)  $a + b + -b = a$ , and
- (ii)  $a + -b + b = a$ , and
  - (iii)  $-b + b + a = a$ , and
  - (iv)  $b + -b + a = a$ , and
  - (v)  $a + (b + -b) = a$ , and
  - (vi)  $a + (-b + b) = a$ , and
  - (vii)  $-b + (b + a) = a$ , and

$$(viii) \quad b + (-b + a) = a.$$

(203)  $G$  is an Abelian additive group if and only if the addition of  $G$  is commutative.

(204)  $0_G$  is Abelian.

(205) If  $A \subseteq B$  and  $C \subseteq D$ , then  $A + C \subseteq B + D$ .

(206) If  $A \subseteq B$ , then  $a + A \subseteq a + B$  and  $A + a \subseteq B + a$ .

(207) If  $H_1$  is a subgroup of  $H_2$ , then  $a + H_1 \subseteq a + H_2$  and  $H_1 + a \subseteq H_2 + a$ .  
The theorem is a consequence of (205).

(208)  $a + H = \{a\} + H$ .

(209)  $H + a = H + \{a\}$ .

(210)  $(A + a) + H = A + (a + H)$ . The theorem is a consequence of (142).

(211)  $(a + H) + A = a + (H + A)$ . The theorem is a consequence of (143).

(212)  $(A + H) + a = A + (H + a)$ . The theorem is a consequence of (143).

(213)  $(H + a) + A = H + (a + A)$ . The theorem is a consequence of (144).

(214)  $(H_1 + a) + H_2 = H_1 + (a + H_2)$ .

Let us consider  $G$ . The functor  $\text{SubGr } G$  yielding a set is defined by

(Def. 29) for every object  $x$ ,  $x \in \text{it}$  iff  $x$  is a strict subgroup of  $G$ .

Note that  $\text{SubGr } G$  is non empty.

Now we state the propositions:

(215) Let us consider a strict additive group  $G$ . Then  $G \in \text{SubGr } G$ . The theorem is a consequence of (100).

(216) If  $G$  is finite, then  $\text{SubGr } G$  is finite.

PROOF: Define  $\mathcal{P}[\text{object}, \text{object}] \equiv$  there exists a strict subgroup  $H$  of  $G$  such that  $\$1 = H$  and  $\$2 =$  the carrier of  $H$ . For every object  $x$  such that  $x \in \text{SubGr } G$  there exists an object  $y$  such that  $\mathcal{P}[x, y]$ . Consider  $f$  being a function such that  $\text{dom } f = \text{SubGr } G$  and for every object  $x$  such that  $x \in \text{SubGr } G$  holds  $\mathcal{P}[x, f(x)]$  from [4, Sch. 1].  $\text{rng } f \subseteq 2^\alpha$ , where  $\alpha$  is the carrier of  $G$ .  $f$  is one-to-one.  $\square$

Let us consider  $G$ ,  $a$ , and  $b$ . The functor  $a \cdot b$  yielding an element of  $G$  is defined by the term

(Def. 30)  $-b + a + b$ .

Now we state the propositions:

(217) If  $a \cdot g = b \cdot g$ , then  $a = b$ . The theorem is a consequence of (6).

(218)  $0_G \cdot a = 0_G$ .

(219) If  $a \cdot b = 0_G$ , then  $a = 0_G$ . The theorem is a consequence of (11) and (7).

(220)  $a \cdot 0_G = a$ . The theorem is a consequence of (8).

(221)  $a \cdot a = a.$

(222) (i)  $a \cdot (-a) = a,$  and

(ii)  $(-a) \cdot a = -a.$

(223)  $a \cdot b = a$  if and only if  $a + b = b + a.$  The theorem is a consequence of (12).

(224)  $(a + b) \cdot g = a \cdot g + b \cdot g.$

(225)  $a \cdot g \cdot h = a \cdot (g + h).$  The theorem is a consequence of (16).

(226) (i)  $a \cdot b \cdot (-b) = a,$  and

(ii)  $a \cdot (-b) \cdot b = a.$

The theorem is a consequence of (225) and (220).

(227)  $(-a) \cdot b = -a \cdot b.$  The theorem is a consequence of (16).

(228)  $(n \cdot a) \cdot b = n \cdot (a \cdot b).$

(229)  $(i \cdot a) \cdot b = i \cdot (a \cdot b).$  The theorem is a consequence of (29) and (227).

(230) If  $G$  is an Abelian additive group, then  $a \cdot b = a.$  The theorem is a consequence of (202).

(231) If for every  $a$  and  $b,$   $a \cdot b = a,$  then  $G$  is Abelian. The theorem is a consequence of (223).

Let us consider  $G, A,$  and  $B.$  The functor  $A \cdot B$  yielding a subset of  $G$  is defined by the term

(Def. 31)  $\{g \cdot h : g \in A \text{ and } h \in B\}.$

Now we state the propositions:

(232)  $x \in A \cdot B$  if and only if there exists  $g$  and there exists  $h$  such that  $x = g \cdot h$  and  $g \in A$  and  $h \in B.$

(233)  $A \cdot B \neq \emptyset$  if and only if  $A \neq \emptyset$  and  $B \neq \emptyset.$  The theorem is a consequence of (232).

(234)  $A \cdot B \subseteq -B + A + B.$

(235)  $(A + B) \cdot C \subseteq A \cdot C + B \cdot C.$  The theorem is a consequence of (224).

(236)  $A \cdot B \cdot C = A \cdot (B + C).$  The theorem is a consequence of (225).

(237)  $(-A) \cdot B = -A \cdot B.$  The theorem is a consequence of (227).

(238)  $\{a\} \cdot \{b\} = \{a \cdot b\}.$  The theorem is a consequence of (49), (64), (233), and (234).

(239)  $\{a\} \cdot \{b, c\} = \{a \cdot b, a \cdot c\}.$

(240)  $\{a, b\} \cdot \{c\} = \{a \cdot c, b \cdot c\}.$

(241)  $\{a, b\} \cdot \{c, d\} = \{a \cdot c, a \cdot d, b \cdot c, b \cdot d\}.$

Let us consider  $G, A,$  and  $g.$  The functors:  $A \cdot g$  and  $g \cdot A$  yielding subsets of  $G$  are defined by terms,

(Def. 32)  $A \cdot \{g\}$ ,

(Def. 33)  $\{g\} \cdot A$ ,

respectively. Now we state the propositions:

(242)  $x \in A \cdot g$  if and only if there exists  $h$  such that  $x = h \cdot g$  and  $h \in A$ .

(243)  $x \in g \cdot A$  if and only if there exists  $h$  such that  $x = g \cdot h$  and  $h \in A$ .

(244)  $g \cdot A \subseteq -A + g + A$ . The theorem is a consequence of (243) and (74).

(245)  $A \cdot B \cdot g = A \cdot (B + g)$ .

(246)  $A \cdot g \cdot B = A \cdot (g + B)$ .

(247)  $g \cdot A \cdot B = g \cdot (A + B)$ .

(248)  $A \cdot a \cdot b = A \cdot (a + b)$ . The theorem is a consequence of (236) and (64).

(249)  $a \cdot A \cdot b = a \cdot (A + b)$ .

(250)  $a \cdot b \cdot A = a \cdot (b + A)$ . The theorem is a consequence of (238) and (236).

(251)  $A \cdot g = -g + A + g$ . The theorem is a consequence of (234), (49), (74), (73), and (242).

(252)  $(A + B) \cdot a \subseteq A \cdot a + B \cdot a$ .

(253)  $A \cdot 0_G = A$ . The theorem is a consequence of (251), (83), and (8).

(254) If  $A \neq \emptyset$ , then  $0_G \cdot A = \{0_G\}$ . The theorem is a consequence of (243) and (218).

(255) (i)  $A \cdot a \cdot (-a) = A$ , and

(ii)  $A \cdot (-a) \cdot a = A$ .

The theorem is a consequence of (248) and (253).

(256)  $G$  is an Abelian additive group if and only if for every  $A$  and  $B$  such that  $B \neq \emptyset$  holds  $A \cdot B = A$ . The theorem is a consequence of (230), (238), and (231).

(257)  $G$  is an Abelian additive group if and only if for every  $A$  and  $g$ ,  $A \cdot g = A$ . The theorem is a consequence of (256), (238), and (231).

(258)  $G$  is an Abelian additive group if and only if for every  $A$  and  $g$  such that  $A \neq \emptyset$  holds  $g \cdot A = \{g\}$ . The theorem is a consequence of (256), (238), and (231).

Let us consider  $G$ ,  $H$ , and  $a$ . The functor  $H \cdot a$  yielding a strict subgroup of  $G$  is defined by

(Def. 34) the carrier of  $it = \overline{H} \cdot a$ .

Now we state the propositions:

(259)  $x \in H \cdot a$  if and only if there exists  $g$  such that  $x = g \cdot a$  and  $g \in H$ . The theorem is a consequence of (242).

(260) The carrier of  $H \cdot a = -a + H + a$ . The theorem is a consequence of (251).

(261)  $H \cdot a \cdot b = H \cdot (a + b)$ . The theorem is a consequence of (248) and (105).

Let us consider a strict subgroup  $H$  of  $G$ . Now we state the propositions:

(262)  $H \cdot 0_G = H$ . The theorem is a consequence of (253) and (105).

(263) (i)  $H \cdot a \cdot (-a) = H$ , and

(ii)  $H \cdot (-a) \cdot a = H$ .

The theorem is a consequence of (261) and (262).

Now we state the propositions:

(264)  $(H_1 \cap H_2) \cdot a = H_1 \cdot a \cap (H_2 \cdot a)$ . The theorem is a consequence of (259), (128), and (217).

(265)  $\overline{H} = \overline{H \cdot a}$ .

PROOF: Define  $\mathcal{F}$ (element of  $G$ ) =  $\$1 \cdot a$ . Consider  $f$  being a function from the carrier of  $G$  into the carrier of  $G$  such that for every  $g$ ,  $f(g) = \mathcal{F}(g)$  from [15, Sch. 4]. Set  $g = f \upharpoonright$ (the carrier of  $H$ ).  $\text{rng } g =$  the carrier of  $H \cdot a$  by [46, (62)], (88), (242), [14, (47)].  $g$  is one-to-one by [46, (62)], (88), [14, (47)], (217).  $\square$

(266)  $H$  is finite if and only if  $H \cdot a$  is finite. The theorem is a consequence of (265).

Let us consider  $G$  and  $a$ . Let  $H$  be a finite subgroup of  $G$ . Observe that  $H \cdot a$  is finite.

Now we state the propositions:

(267) Let us consider a finite subgroup  $H$  of  $G$ . Then  $\overline{H} = \overline{H \cdot a}$ .

(268)  $\mathbf{0}_G \cdot a = \mathbf{0}_G$ . The theorem is a consequence of (238) and (218).

(269) Let us consider a strict subgroup  $H$  of  $G$ . If  $H \cdot a = \mathbf{0}_G$ , then  $H = \mathbf{0}_G$ . The theorem is a consequence of (266), (115), (265), and (116).

(270) Let us consider an additive group  $G$ , and an element  $a$  of  $G$ . Then  $\Omega_G \cdot a = \Omega_G$ . The theorem is a consequence of (225), (220), and (259).

(271) Let us consider a strict subgroup  $H$  of  $G$ . If  $H \cdot a = G$ , then  $H = G$ . The theorem is a consequence of (259), (217), and (108).

(272)  $|\bullet : H| = |\bullet : H \cdot a|$ .

PROOF: Define  $\mathcal{P}$ [object, object]  $\equiv$  there exists  $b$  such that  $\$1 = b + H$  and  $\$2 = b \cdot a + H \cdot a$ . For every object  $x$  such that  $x \in$  the left cosets of  $H$  there exists an object  $y$  such that  $\mathcal{P}[x, y]$ . Consider  $f$  being a function such that  $\text{dom } f =$  the left cosets of  $H$  and for every object  $x$  such that  $x \in$  the left cosets of  $H$  holds  $\mathcal{P}[x, f(x)]$  from [4, Sch. 1]. For every  $x, y_1$ , and  $y_2$  such that  $x \in$  the left cosets of  $H$  and  $\mathcal{P}[x, y_1]$  and  $\mathcal{P}[x, y_2]$  holds  $y_1 = y_2$ .  $\text{rng } f =$  the left cosets of  $H \cdot a$ .  $f$  is one-to-one.  $\square$

(273) If the left cosets of  $H$  is finite, then  $|\bullet : H|_{\mathbb{N}} = |\bullet : H \cdot a|_{\mathbb{N}}$ . The theorem is a consequence of (272).

(274) If  $G$  is an Abelian additive group, then for every strict subgroup  $H$  of  $G$  and for every  $a$ ,  $H \cdot a = H$ . The theorem is a consequence of (260), (158), (153), (155), and (105).

Let us consider  $G$ ,  $a$ , and  $b$ . We say that  $a$  and  $b$  are conjugated if and only if

(Def. 35) there exists  $g$  such that  $a = b \cdot g$ .

Now we state the proposition:

(275)  $a$  and  $b$  are conjugated if and only if there exists  $g$  such that  $b = a \cdot g$ . The theorem is a consequence of (226).

Let us consider  $G$ ,  $a$ , and  $b$ . Observe that  $a$  and  $b$  are conjugated is reflexive and symmetric.

Now we state the propositions:

(276) If  $a$  and  $b$  are conjugated and  $b$  and  $c$  are conjugated, then  $a$  and  $c$  are conjugated. The theorem is a consequence of (225).

(277) If  $a$  and  $0_G$  are conjugated or  $0_G$  and  $a$  are conjugated, then  $a = 0_G$ . The theorem is a consequence of (275) and (219).

(278)  $a \cdot \overline{\Omega_G} = \{b : a \text{ and } b \text{ are conjugated}\}$ . The theorem is a consequence of (243).

Let us consider  $G$  and  $a$ . The functor  $a^\bullet$  yielding a subset of  $G$  is defined by the term

(Def. 36)  $a \cdot \overline{\Omega_G}$ .

Now we state the propositions:

(279)  $x \in a^\bullet$  if and only if there exists  $b$  such that  $b = x$  and  $a$  and  $b$  are conjugated. The theorem is a consequence of (278).

(280)  $a \in b^\bullet$  if and only if  $a$  and  $b$  are conjugated. The theorem is a consequence of (279).

(281)  $a \cdot g \in a^\bullet$ .

(282)  $a \in a^\bullet$ .

(283) If  $a \in b^\bullet$ , then  $b \in a^\bullet$ . The theorem is a consequence of (280).

(284)  $a^\bullet = b^\bullet$  if and only if  $a^\bullet$  meets  $b^\bullet$ . The theorem is a consequence of (280), (279), and (276).

(285)  $a^\bullet = \{0_G\}$  if and only if  $a = 0_G$ . The theorem is a consequence of (280), (279), and (277).

(286)  $a^\bullet + A = A + a^\bullet$ . The theorem is a consequence of (280), (202), (226), (224), (221), (225), (279), and (275).

Let us consider  $G$ ,  $A$ , and  $B$ . We say that  $A$  and  $B$  are conjugated if and only if

(Def. 37) there exists  $g$  such that  $A = B \cdot g$ .

Now we state the propositions:

- (287)  $A$  and  $B$  are conjugated if and only if there exists  $g$  such that  $B = A \cdot g$ .  
The theorem is a consequence of (255).
- (288)  $A$  and  $A$  are conjugated. The theorem is a consequence of (253).
- (289) If  $A$  and  $B$  are conjugated, then  $B$  and  $A$  are conjugated. The theorem is a consequence of (255).

Let us consider  $G$ ,  $A$ , and  $B$ . Let us observe that  $A$  and  $B$  are conjugated is reflexive and symmetric.

Now we state the propositions:

- (290) If  $A$  and  $B$  are conjugated and  $B$  and  $C$  are conjugated, then  $A$  and  $C$  are conjugated. The theorem is a consequence of (248).
- (291)  $\{a\}$  and  $\{b\}$  are conjugated if and only if  $a$  and  $b$  are conjugated.  
PROOF: If  $\{a\}$  and  $\{b\}$  are conjugated, then  $a$  and  $b$  are conjugated by (287), (238), (275), [17, (3)]. Consider  $g$  such that  $a \cdot g = b$ .  $\{b\} = \{a\} \cdot g$ .  
 $\square$
- (292) If  $A$  and  $\overline{H_1}$  are conjugated, then there exists a strict subgroup  $H_2$  of  $G$  such that the carrier of  $H_2 = A$ .

Let us consider  $G$  and  $A$ . The functor  $A^\bullet$  yielding a family of subsets of  $G$  is defined by the term

(Def. 38)  $\{B : A \text{ and } B \text{ are conjugated}\}$ .

Now we state the propositions:

- (293)  $x \in A^\bullet$  if and only if there exists  $B$  such that  $x = B$  and  $A$  and  $B$  are conjugated.
- (294)  $A \in B^\bullet$  if and only if  $A$  and  $B$  are conjugated.
- (295)  $A \cdot g \in A^\bullet$ . The theorem is a consequence of (287).
- (296)  $A \in A^\bullet$ .
- (297) If  $A \in B^\bullet$ , then  $B \in A^\bullet$ . The theorem is a consequence of (294).
- (298)  $A^\bullet = B^\bullet$  if and only if  $A^\bullet$  meets  $B^\bullet$ . The theorem is a consequence of (294) and (290).
- (299)  $\{a\}^\bullet = \{\{b\} : b \in a^\bullet\}$ . The theorem is a consequence of (287), (275), (280), (238), and (291).
- (300) If  $G$  is finite, then  $A^\bullet$  is finite.

Let us consider  $G$ ,  $H_1$ , and  $H_2$ . We say that  $H_1$  and  $H_2$  are conjugated if and only if

(Def. 39) there exists  $g$  such that the additive magma of  $H_1 = H_2 \cdot g$ .

Now we state the propositions:

(301) Let us consider strict subgroups  $H_1, H_2$  of  $G$ . Then  $H_1$  and  $H_2$  are conjugated if and only if there exists  $g$  such that  $H_2 = H_1 \cdot g$ . The theorem is a consequence of (263).

(302) Let us consider a strict subgroup  $H_1$  of  $G$ . Then  $H_1$  and  $H_1$  are conjugated. The theorem is a consequence of (262).

(303) Let us consider strict subgroups  $H_1, H_2$  of  $G$ . If  $H_1$  and  $H_2$  are conjugated, then  $H_2$  and  $H_1$  are conjugated. The theorem is a consequence of (263).

Let us consider  $G$ . Let  $H_1, H_2$  be strict subgroups of  $G$ . Observe that  $H_1$  and  $H_2$  are conjugated is reflexive and symmetric.

Now we state the proposition:

(304) Let us consider strict subgroups  $H_1, H_2$  of  $G$ . Suppose  $H_1$  and  $H_2$  are conjugated and  $H_2$  and  $H_3$  are conjugated. Then  $H_1$  and  $H_3$  are conjugated. The theorem is a consequence of (261).

In the sequel  $L$  denotes a subset of  $\text{SubGr } G$ .

Let us consider  $G$  and  $H$ . The functor  $H^\bullet$  yielding a subset of  $\text{SubGr } G$  is defined by

(Def. 40) for every object  $x$ ,  $x \in it$  iff there exists a strict subgroup  $H_1$  of  $G$  such that  $x = H_1$  and  $H$  and  $H_1$  are conjugated.

Now we state the propositions:

(305) If  $x \in H^\bullet$ , then  $x$  is a strict subgroup of  $G$ .

(306) Let us consider strict subgroups  $H_1, H_2$  of  $G$ . Then  $H_1 \in H_2^\bullet$  if and only if  $H_1$  and  $H_2$  are conjugated.

Let us consider a strict subgroup  $H$  of  $G$ . Now we state the propositions:

(307)  $H \cdot g \in H^\bullet$ . The theorem is a consequence of (301).

(308)  $H \in H^\bullet$ .

Let us consider strict subgroups  $H_1, H_2$  of  $G$ . Now we state the propositions:

(309) If  $H_1 \in H_2^\bullet$ , then  $H_2 \in H_1^\bullet$ . The theorem is a consequence of (306).

(310)  $H_1^\bullet = H_2^\bullet$  if and only if  $H_1^\bullet$  meets  $H_2^\bullet$ . The theorem is a consequence of (308), (305), (306), and (304).

Now we state the propositions:

(311) If  $G$  is finite, then  $H^\bullet$  is finite.

(312) Let us consider a strict subgroup  $H_1$  of  $G$ . Then  $H_1$  and  $H_2$  are conjugated if and only if  $\overline{H_1}$  and  $\overline{H_2}$  are conjugated.



Let us consider  $G$ . Let  $I_1$  be a subgroup of  $G$ . We say that  $I_1$  is normal if and only if

(Def. 41) for every  $a$ ,  $I_1 \cdot a =$  the additive magma of  $I_1$ .

Let us note that there exists a subgroup of  $G$  which is strict and normal.

From now on  $N_2$  denotes a normal subgroup of  $G$ .

Now we state the propositions:

(313) (i)  $\mathbf{0}_G$  is normal, and

(ii)  $\Omega_G$  is normal.

(314) Let us consider strict, normal subgroups  $N_1, N_2$  of  $G$ . Then  $N_1 \cap N_2$  is normal. The theorem is a consequence of (264).

(315) Let us consider a strict subgroup  $H$  of  $G$ . If  $G$  is an Abelian additive group, then  $H$  is normal.

(316)  $H$  is a normal subgroup of  $G$  if and only if for every  $a$ ,  $a + H = H + a$ . The theorem is a consequence of (260), (79), (151), (83), (153), (155), and (105).

Let us consider a subgroup  $H$  of  $G$ . Now we state the propositions:

(317)  $H$  is a normal subgroup of  $G$  if and only if for every  $a$ ,  $a + H \subseteq H + a$ . The theorem is a consequence of (316), (205), (151), (155), (152), (80), and (83).

(318)  $H$  is a normal subgroup of  $G$  if and only if for every  $a$ ,  $H + a \subseteq a + H$ . The theorem is a consequence of (316), (205), (151), (155), (152), (80), and (83).

(319)  $H$  is a normal subgroup of  $G$  if and only if for every  $A$ ,  $A + H = H + A$ . The theorem is a consequence of (140), (149), (316), (150), and (141).

Let us consider a strict subgroup  $H$  of  $G$ . Now we state the propositions:

(320)  $H$  is a normal subgroup of  $G$  if and only if for every  $a$ ,  $H$  is a subgroup of  $H \cdot a$ . The theorem is a consequence of (100), (260), (80), (83), (207), and (318).

(321)  $H$  is a normal subgroup of  $G$  if and only if for every  $a$ ,  $H \cdot a$  is a subgroup of  $H$ . The theorem is a consequence of (100), (260), (80), (83), (207), and (317).

(322)  $H$  is a normal subgroup of  $G$  if and only if  $H^\bullet = \{H\}$ .

PROOF: If  $H$  is a normal subgroup of  $G$ , then  $H^\bullet = \{H\}$  by (301), (308), [17, (31)].  $H$  is normal.  $\square$

(323)  $H$  is a normal subgroup of  $G$  if and only if for every  $a$  such that  $a \in H$  holds  $a^\bullet \subseteq \overline{H}$ . The theorem is a consequence of (279), (275), (259), and (226).

Let us consider strict, normal subgroups  $N_1, N_2$  of  $G$ . Now we state the propositions:

(324)  $\overline{N_1} + \overline{N_2} = \overline{N_2} + \overline{N_1}$ .

(325) There exists a strict, normal subgroup  $N$  of  $G$  such that the carrier of  $N = \overline{N_1} + \overline{N_2}$ . The theorem is a consequence of (124), (75), (316), (76), and (77).

Now we state the propositions:

(326) Let us consider a normal subgroup  $N$  of  $G$ . Then the left cosets of  $N =$  the right cosets of  $N$ . The theorem is a consequence of (316).

(327) Let us consider a subgroup  $H$  of  $G$ . Suppose the left cosets of  $H$  is finite and  $|\bullet : H|_{\mathbb{N}} = 2$ . Then  $H$  is a normal subgroup of  $G$ .

PROOF: There exists a finite set  $B$  such that  $B =$  the left cosets of  $H$  and  $|\bullet : H|_{\mathbb{N}} = \overline{B}$ . Consider  $x, y$  being objects such that  $x \neq y$  and the left cosets of  $H = \{x, y\}$ .  $\overline{H} \in$  the left cosets of  $H$ . Consider  $z_3$  being an object such that  $\{x, y\} = \{\overline{H}, z_3\}$ .  $\overline{H}$  misses  $z_3$  by (155), (161), [34, (29)], [17, (4)].  $\cup$ (the left cosets of  $H$ ) = the carrier of  $G$  and  $\cup$ (the left cosets of  $H$ ) =  $\overline{H} \cup z_3$ .  $\cup$ (the right cosets of  $H$ ) = the carrier of  $G$  and  $z_3 =$  (the carrier of  $G$ )  $\setminus \overline{H}$ . There exists a finite set  $C$  such that  $C =$  the right cosets of  $H$  and  $|\bullet : H|_{\mathbb{N}} = \overline{C}$ . Consider  $z_1, z_2$  being objects such that  $z_1 \neq z_2$  and the right cosets of  $H = \{z_1, z_2\}$ .  $\overline{H} \in$  the right cosets of  $H$ . Consider  $z_4$  being an object such that  $\{z_1, z_2\} = \{\overline{H}, z_4\}$ .  $\overline{H}$  misses  $z_4$  by (155), (167), [34, (29)], [17, (4)].  $\square$

Let us consider  $G$  and  $A$ . The functor  $N(A)$  yielding a strict subgroup of  $G$  is defined by

(Def. 42) the carrier of  $it = \{h : A \cdot h = A\}$ .

Now we state the propositions:

(328)  $x \in N(A)$  if and only if there exists  $h$  such that  $x = h$  and  $A \cdot h = A$ .

(329)  $\overline{A^\bullet} = |\bullet : N(A)|$ .

PROOF: Define  $\mathcal{P}[\text{object}, \text{object}] \equiv$  there exists  $a$  such that  $\$1 = A \cdot a$  and  $\$2 = N(A) + a$ . For every object  $x$  such that  $x \in A^\bullet$  there exists an object  $y$  such that  $\mathcal{P}[x, y]$ . Consider  $f$  being a function such that  $\text{dom } f = A^\bullet$  and for every object  $x$  such that  $x \in A^\bullet$  holds  $\mathcal{P}[x, f(x)]$  from [4, Sch. 1]. For every  $x, y_1,$  and  $y_2$  such that  $x \in A^\bullet$  and  $\mathcal{P}[x, y_1]$  and  $\mathcal{P}[x, y_2]$  holds  $y_1 = y_2$ .  $\text{rng } f =$  the right cosets of  $N(A)$ .  $f$  is one-to-one.  $\square$

(330) Suppose  $A^\bullet$  is finite or the left cosets of  $N(A)$  is finite. Then there exists a finite set  $C$  such that

(i)  $C = A^\bullet,$  and

(ii)  $\overline{C} = |\bullet : N(A)|_{\mathbb{N}}$ .

The theorem is a consequence of (329).

$$(331) \quad \overline{a^\bullet} = |\bullet : N(\{a\})|.$$

PROOF: Define  $\mathcal{F}(\text{object}) = \{\$1\}$ . Consider  $f$  being a function such that  $\text{dom } f = a^\bullet$  and for every object  $x$  such that  $x \in a^\bullet$  holds  $f(x) = \mathcal{F}(x)$  from [14, Sch. 3].  $\text{rng } f = \{a\}^\bullet$ .  $f$  is one-to-one by [17, (3)].  $\square$

(332) Suppose  $a^\bullet$  is finite or the left cosets of  $N(\{a\})$  is finite. Then there exists a finite set  $C$  such that

$$(i) \quad C = a^\bullet, \text{ and}$$

$$(ii) \quad \overline{C} = |\bullet : N(\{a\})|_{\mathbb{N}}.$$

The theorem is a consequence of (331).

Let us consider  $G$  and  $H$ . The functor  $N(H)$  yielding a strict subgroup of  $G$  is defined by the term

$$(\text{Def. 43}) \quad N(\overline{H}).$$

Let us consider a strict subgroup  $H$  of  $G$ . Now we state the propositions:

(333)  $x \in N(H)$  if and only if there exists  $h$  such that  $x = h$  and  $H \cdot h = H$ . The theorem is a consequence of (328).

$$(334) \quad \overline{H^\bullet} = |\bullet : N(H)|.$$

PROOF: Define  $\mathcal{P}[\text{object}, \text{object}] \equiv$  there exists a strict subgroup  $H_1$  of  $G$  such that  $\$1 = H_1$  and  $\$2 = \overline{H_1}$ . For every object  $x$  such that  $x \in H^\bullet$  there exists an object  $y$  such that  $\mathcal{P}[x, y]$ . Consider  $f$  being a function such that  $\text{dom } f = H^\bullet$  and for every object  $x$  such that  $x \in H^\bullet$  holds  $\mathcal{P}[x, f(x)]$  from [4, Sch. 1].  $\text{rng } f = \overline{H^\bullet}$ .  $f$  is one-to-one.  $\square$

(335) Suppose  $H^\bullet$  is finite or the left cosets of  $N(H)$  is finite. Then there exists a finite set  $C$  such that

$$(i) \quad C = H^\bullet, \text{ and}$$

$$(ii) \quad \overline{C} = |\bullet : N(H)|_{\mathbb{N}}.$$

The theorem is a consequence of (334).

Now we state the proposition:

(336) Let us consider a strict additive group  $G$ , and a strict subgroup  $H$  of  $G$ . Then  $H$  is a normal subgroup of  $G$  if and only if  $N(H) = G$ . The theorem is a consequence of (333) and (108).

Let us consider a strict additive group  $G$ . Now we state the propositions:

(337)  $N(\mathbf{0}_G) = G$ . The theorem is a consequence of (313) and (336).

(338)  $N(\Omega_G) = G$ . The theorem is a consequence of (313) and (336).

## 4. TOPOLOGICAL GROUPS – TOPGRP\_1

In the sequel  $S, R$  denote 1-sorted structures,  $X$  denotes a subset of  $R$ ,  $T$  denotes a topological structure,  $x$  denotes a set,  $H$  denotes a non empty additive magma,  $P, Q, P_1, Q_1$  denote subsets of  $H$ , and  $h$  denotes an element of  $H$ .

Now we state the proposition:

$$(339) \quad \text{If } P \subseteq P_1 \text{ and } Q \subseteq Q_1, \text{ then } P + Q \subseteq P_1 + Q_1.$$

Let us assume that  $P \subseteq Q$ . Now we state the propositions:

$$(340) \quad P + h \subseteq Q + h. \text{ The theorem is a consequence of (74).}$$

$$(341) \quad h + P \subseteq h + Q. \text{ The theorem is a consequence of (73).}$$

From now on  $a$  denotes an element of  $G$ .

Now we state the propositions:

$$(342) \quad a \in -A \text{ if and only if } -a \in A.$$

$$(343) \quad A \subseteq B \text{ if and only if } -A \subseteq -B.$$

$$(344) \quad (\text{add inverse } G)^\circ A = -A.$$

$$(345) \quad (\text{add inverse } G)^{-1}(A) = -A.$$

$$(346) \quad \text{add inverse } G \text{ is one-to-one. The theorem is a consequence of (9).}$$

$$(347) \quad \text{rng add inverse } G = \text{the carrier of } G.$$

Let  $G$  be an additive group. One can verify that add inverse  $G$  is one-to-one and onto.

Now we state the propositions:

$$(348) \quad (\text{add inverse } G)^{-1} = \text{add inverse } G.$$

$$(349) \quad (\text{The addition of } H)^\circ(P \times Q) = P + Q.$$

Let  $G$  be a non empty additive magma and  $a$  be an element of  $G$ . The functors:  $a^+$  and  $^+a$  yielding functions from  $G$  into  $G$  are defined by conditions,

$$(\text{Def. 44}) \quad \text{for every element } x \text{ of } G, a^+(x) = a + x,$$

$$(\text{Def. 45}) \quad \text{for every element } x \text{ of } G, ^+a(x) = x + a,$$

respectively. Let  $G$  be an additive group. One can verify that  $a^+$  is one-to-one and onto and  $^+a$  is one-to-one and onto.

Now we state the propositions:

$$(350) \quad (h^+)^\circ P = h + P. \text{ The theorem is a consequence of (73).}$$

$$(351) \quad (^+h)^\circ P = P + h. \text{ The theorem is a consequence of (74).}$$

$$(352) \quad (a^+)^{-1} = (-a)^+.$$

$$(353) \quad (^+a)^{-1} = ^+(-a).$$

We consider topological additive group structures which extend additive magmas and topological structures and are systems

⟨a carrier, an addition, a topology⟩

where the carrier is a set, the addition is a binary operation on the carrier, the topology is a family of subsets of the carrier.

Let  $A$  be a non empty set,  $R$  be a binary operation on  $A$ , and  $T$  be a family of subsets of  $A$ . Let us observe that  $\langle A, R, T \rangle$  is non empty.

Let  $x$  be a set,  $R$  be a binary operation on  $\{x\}$ , and  $T$  be a family of subsets of  $\{x\}$ . Observe that  $\langle \{x\}, R, T \rangle$  is trivial and every 1-element additive magma is additive group-like, add-associative, and Abelian and there exists a topological additive group structure which is strict and non empty and there exists a topological additive group structure which is strict, topological space-like, and 1-element.

Let  $G$  be an additive group-like, add-associative, non empty topological additive group structure. We say that  $G$  is inverse-continuous if and only if

(Def. 46) add inverse  $G$  is continuous.

Let  $G$  be a topological space-like topological additive group structure. We say that  $G$  is continuous if and only if

(Def. 47) for every function  $f$  from  $G \times G$  into  $G$  such that  $f =$  the addition of  $G$  holds  $f$  is continuous.

One can check that there exists a topological space-like, additive group-like, add-associative, 1-element topological additive group structure which is strict, Abelian, inverse-continuous, and continuous.

A semi additive topological group is a topological space-like, additive group-like, add-associative, non empty topological additive group structure.

A topological additive group is an inverse-continuous, continuous semi additive topological group. Now we state the propositions:

(354) Let us consider a continuous, non empty, topological space-like topological additive group structure  $T$ , elements  $a, b$  of  $T$ , and a neighbourhood  $W$  of  $a + b$ . Then there exists an open neighbourhood  $A$  of  $a$  and there exists an open neighbourhood  $B$  of  $b$  such that  $A + B \subseteq W$ .

(355) Let us consider a topological space-like, non empty topological additive group structure  $T$ . Suppose for every elements  $a, b$  of  $T$  for every neighbourhood  $W$  of  $a + b$ , there exists a neighbourhood  $A$  of  $a$  and there exists a neighbourhood  $B$  of  $b$  such that  $A + B \subseteq W$ . Then  $T$  is continuous.

PROOF: For every point  $W$  of  $T \times T$  and for every neighbourhood  $G$  of  $f(W)$ , there exists a neighbourhood  $H$  of  $W$  such that  $f^\circ H \subseteq G$  by [32, (10)], (349).  $\square$

(356) Let us consider an inverse-continuous semi additive topological group  $T$ , an element  $a$  of  $T$ , and a neighbourhood  $W$  of  $-a$ . Then there exists an open neighbourhood  $A$  of  $a$  such that  $-A \subseteq W$ .

(357) Let us consider a semi additive topological group  $T$ . Suppose for every

element  $a$  of  $T$  for every neighbourhood  $W$  of  $-a$ , there exists a neighbourhood  $A$  of  $a$  such that  $-A \subseteq W$ . Then  $T$  is inverse-continuous. The theorem is a consequence of (344).

(358) Let us consider a topological additive group  $T$ , elements  $a, b$  of  $T$ , and a neighbourhood  $W$  of  $a + -b$ . Then there exists an open neighbourhood  $A$  of  $a$  and there exists an open neighbourhood  $B$  of  $b$  such that  $A + -B \subseteq W$ . The theorem is a consequence of (354) and (356).

(359) Let us consider a semi additive topological group  $T$ . Suppose for every elements  $a, b$  of  $T$  for every neighbourhood  $W$  of  $a + -b$ , there exists a neighbourhood  $A$  of  $a$  and there exists a neighbourhood  $B$  of  $b$  such that  $A + -B \subseteq W$ . Then  $T$  is a topological additive group.

PROOF: For every element  $a$  of  $T$  and for every neighbourhood  $W$  of  $-a$ , there exists a neighbourhood  $A$  of  $a$  such that  $-A \subseteq W$  by [28, (4)]. For every elements  $a, b$  of  $T$  and for every neighbourhood  $W$  of  $a + b$ , there exists a neighbourhood  $A$  of  $a$  and there exists a neighbourhood  $B$  of  $b$  such that  $A + B \subseteq W$ .  $\square$

Let  $G$  be a continuous, non empty, topological space-like topological additive group structure and  $a$  be an element of  $G$ . One can check that  $a^+$  is continuous and  $+a$  is continuous.

Let us consider a continuous semi additive topological group  $G$  and an element  $a$  of  $G$ . Now we state the propositions:

(360)  $a^+$  is a homeomorphism of  $G$ . The theorem is a consequence of (352).

(361)  $+a$  is a homeomorphism of  $G$ . The theorem is a consequence of (353).

Let  $G$  be a continuous semi additive topological group and  $a$  be an element of  $G$ . The functors:  $a^+$  and  $+a$  yield homeomorphisms of  $G$ . Now we state the proposition:

(362) Let us consider an inverse-continuous semi additive topological group  $G$ . Then add inverse  $G$  is a homeomorphism of  $G$ . The theorem is a consequence of (348).

Let  $G$  be an inverse-continuous semi additive topological group. Let us note that the functor add inverse  $G$  yields a homeomorphism of  $G$ . Let us note that every semi additive topological group which is continuous is also homogeneous.

Let us consider a continuous semi additive topological group  $G$ , a closed subset  $F$  of  $G$ , and an element  $a$  of  $G$ . Now we state the propositions:

(363)  $F + a$  is closed. The theorem is a consequence of (351).

(364)  $a + F$  is closed. The theorem is a consequence of (350).

Let  $G$  be a continuous semi additive topological group,  $F$  be a closed subset of  $G$ , and  $a$  be an element of  $G$ . Let us note that  $F + a$  is closed and  $a + F$  is

closed.

Now we state the proposition:

(365) Let us consider an inverse-continuous semi additive topological group  $G$ , and a closed subset  $F$  of  $G$ . Then  $-F$  is closed. The theorem is a consequence of (344).

Let  $G$  be an inverse-continuous semi additive topological group and  $F$  be a closed subset of  $G$ . One can verify that  $-F$  is closed.

Let us consider a continuous semi additive topological group  $G$ , an open subset  $O$  of  $G$ , and an element  $a$  of  $G$ . Now we state the propositions:

(366)  $O + a$  is open. The theorem is a consequence of (351).

(367)  $a + O$  is open. The theorem is a consequence of (350).

Let  $G$  be a continuous semi additive topological group,  $A$  be an open subset of  $G$ , and  $a$  be an element of  $G$ . One can check that  $A + a$  is open and  $a + A$  is open.

Now we state the proposition:

(368) Let us consider an inverse-continuous semi additive topological group  $G$ , and an open subset  $O$  of  $G$ . Then  $-O$  is open. The theorem is a consequence of (344).

Let  $G$  be an inverse-continuous semi additive topological group and  $A$  be an open subset of  $G$ . Observe that  $-A$  is open.

Let us consider a continuous semi additive topological group  $G$  and subsets  $A, O$  of  $G$ .

Let us assume that  $O$  is open. Now we state the propositions:

(369)  $O + A$  is open.

PROOF:  $\text{Int}(O + A) = O + A$  by [48, (16)], (74), [48, (22)].  $\square$

(370)  $A + O$  is open.

PROOF:  $\text{Int}(A + O) = A + O$  by [48, (16)], (73), [48, (22)].  $\square$

Let  $G$  be a continuous semi additive topological group,  $A$  be an open subset of  $G$ , and  $B$  be a subset of  $G$ . Note that  $A + B$  is open and  $B + A$  is open.

Now we state the propositions:

(371) Let us consider an inverse-continuous semi additive topological group  $G$ , a point  $a$  of  $G$ , and a neighbourhood  $A$  of  $a$ . Then  $-A$  is a neighbourhood of  $-a$ . The theorem is a consequence of (343).

(372) Let us consider a topological additive group  $G$ , a point  $a$  of  $G$ , and a neighbourhood  $A$  of  $a + -a$ . Then there exists an open neighbourhood  $B$  of  $a$  such that  $B + -B \subseteq A$ . The theorem is a consequence of (358) and (342).

(373) Let us consider an inverse-continuous semi additive topological group  $G$ , and a dense subset  $A$  of  $G$ . Then  $-A$  is dense. The theorem is a consequence of (345).

Let  $G$  be an inverse-continuous semi additive topological group and  $A$  be a dense subset of  $G$ . Observe that  $-A$  is dense.

Let us consider a continuous semi additive topological group  $G$ , a dense subset  $A$  of  $G$ , and a point  $a$  of  $G$ . Now we state the propositions:

(374)  $a + A$  is dense. The theorem is a consequence of (350).

(375)  $A + a$  is dense. The theorem is a consequence of (351).

Let  $G$  be a continuous semi additive topological group,  $A$  be a dense subset of  $G$ , and  $a$  be a point of  $G$ . Let us observe that  $A + a$  is dense and  $a + A$  is dense.

Now we state the proposition:

(376) Let us consider a topological additive group  $G$ , a basis  $B$  of  $0_G$ , and a dense subset  $M$  of  $G$ . Then  $\{V + x, \text{ where } V \text{ is a subset of } G, x \text{ is a point of } G : V \in B \text{ and } x \in M\}$  is a basis of  $G$ .

PROOF: Set  $Z = \{V + x, \text{ where } V \text{ is a subset of } G, x \text{ is a point of } G : V \in B \text{ and } x \in M\}$ .  $Z \subseteq$  the topology of  $G$  by [38, (12)]. For every subset  $W$  of  $G$  such that  $W$  is open for every point  $a$  of  $G$  such that  $a \in W$  there exists a subset  $V$  of  $G$  such that  $V \in Z$  and  $a \in V$  and  $V \subseteq W$  by (8), [28, (3)], (74), (372).  $Z \subseteq 2^\alpha$ , where  $\alpha$  is the carrier of  $G$ .  $\square$

One can check that every topological additive group is regular.

ACKNOWLEDGEMENT: The author wants to express his gratitude to the anonymous referee for his/her work on merging the three initial articles.

## REFERENCES

- [1] Jonathan Backer, Piotr Rudnicki, and Christoph Schwarzeweller. Ring ideals. *Formalized Mathematics*, 9(3):565–582, 2001.
- [2] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(2):377–382, 1990.
- [3] Grzegorz Bancerek. Cardinal arithmetics. *Formalized Mathematics*, 1(3):543–547, 1990.
- [4] Grzegorz Bancerek. Tarski's classes and ranks. *Formalized Mathematics*, 1(3):563–567, 1990.
- [5] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [6] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [7] Grzegorz Bancerek. Zermelo theorem and axiom of choice. *Formalized Mathematics*, 1(2):265–267, 1990.
- [8] Józef Białas. Group and field definitions. *Formalized Mathematics*, 1(3):433–439, 1990.
- [9] Richard E. Blahut. *Cryptography and Secure Communication*. Cambridge University Press, 2014.
- [10] Sylvie Boldo, Catherine Lelay, and Guillaume Melquiond. Formalization of real analysis:



A survey of proof assistants and libraries. *Mathematical Structures in Computer Science*, pages 1–38, 2014.

- [11] Nicolas Bourbaki. *General Topology: Chapters 1–4*. Springer Science and Business Media, 2013.
- [12] Czesław Byliński. Binary operations. *Formalized Mathematics*, 1(1):175–180, 1990.
- [13] Czesław Byliński. Binary operations applied to finite sequences. *Formalized Mathematics*, 1(4):643–649, 1990.
- [14] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [15] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [16] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [17] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [18] Agata Darmochwał. Compact spaces. *Formalized Mathematics*, 1(2):383–386, 1990.
- [19] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(1):165–167, 1990.
- [20] Agata Darmochwał. Families of subsets, subspaces and mappings in topological spaces. *Formalized Mathematics*, 1(2):257–261, 1990.
- [21] Yuichi Futa, Hiroyuki Okazaki, and Yasunari Shidama.  $\mathbb{Z}$ -modules. *Formalized Mathematics*, 20(1):47–59, 2012. doi:10.2478/v10037-012-0007-z.
- [22] Edwin Hewitt and Kenneth A. Ross. *Abstract Harmonic Analysis: Volume I. Structure of Topological Groups. Integration. Theory Group Representations*, volume 115. Springer Science and Business Media, 2012.
- [23] Johannes Hölzl, Fabian Immler, and Brian Huffman. Type classes and filters for mathematical analysis in Isabelle/HOL. In *Interactive Theorem Proving*, pages 279–294. Springer, 2013.
- [24] Teturo Inui, Yukito Tanabe, and Yositaka Onodera. *Group theory and its applications in physics*, volume 78. Springer Science and Business Media, 2012.
- [25] Artur Kornilowicz. The definition and basic properties of topological groups. *Formalized Mathematics*, 7(2):217–225, 1998.
- [26] Rafał Kwiatek and Grzegorz Zwara. The divisibility of integers and integer relatively primes. *Formalized Mathematics*, 1(5):829–832, 1990.
- [27] Christopher Norman. Basic theory of additive Abelian groups. In *Finitely Generated Abelian Groups and Similarity of Matrices over a Field*, Springer Undergraduate Mathematics Series, pages 47–96. Springer, 2012. ISBN 978-1-4471-2729-1. doi:10.1007/978-1-4471-2730-7\_2.
- [28] Beata Padlewska. Locally connected spaces. *Formalized Mathematics*, 2(1):93–96, 1991.
- [29] Beata Padlewska. Families of sets. *Formalized Mathematics*, 1(1):147–152, 1990.
- [30] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(1):223–230, 1990.
- [31] Alexander Yu. Shibakov and Andrzej Trybulec. The Cantor set. *Formalized Mathematics*, 5(2):233–236, 1996.
- [32] Andrzej Trybulec. A Borsuk theorem on homotopy types. *Formalized Mathematics*, 2(4):535–545, 1991.
- [33] Andrzej Trybulec. Domains and their Cartesian products. *Formalized Mathematics*, 1(1):115–122, 1990.
- [34] Andrzej Trybulec. Enumerated sets. *Formalized Mathematics*, 1(1):25–34, 1990.
- [35] Andrzej Trybulec. Tuples, projections and Cartesian products. *Formalized Mathematics*, 1(1):97–105, 1990.
- [36] Andrzej Trybulec. On the sets inhabited by numbers. *Formalized Mathematics*, 11(4):341–347, 2003.
- [37] Andrzej Trybulec. Semilattice operations on finite subsets. *Formalized Mathematics*, 1(2):369–376, 1990.
- [38] Andrzej Trybulec. Baire spaces, Sober spaces. *Formalized Mathematics*, 6(2):289–294, 1997.

- [39] Andrzej Trybulec and Agata Darmochwał. Boolean domains. *Formalized Mathematics*, 1(1):187–190, 1990.
- [40] Michał J. Trybulec. Integers. *Formalized Mathematics*, 1(3):501–505, 1990.
- [41] Wojciech A. Trybulec. Groups. *Formalized Mathematics*, 1(5):821–827, 1990.
- [42] Wojciech A. Trybulec. Subgroup and cosets of subgroups. *Formalized Mathematics*, 1(5):855–864, 1990.
- [43] Wojciech A. Trybulec. Classes of conjugation. Normal subgroups. *Formalized Mathematics*, 1(5):955–962, 1990.
- [44] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(2):291–296, 1990.
- [45] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [46] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [47] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.
- [48] Mirosław Wysocki and Agata Darmochwał. Subsets of topological spaces. *Formalized Mathematics*, 1(1):231–237, 1990.

*Received April 30, 2015*

---