

# On Multiset Ordering

Grzegorz Bancerek  
Association of Mizar Users  
Białystok, Poland

**Summary.** Formalization of a part of [11]. Unfortunately, not all is possible to be formalized. Namely, in the paper there is a mistake in the proof of Lemma 3. It states that there exists  $x \in M_1$  such that  $M_1(x) > N_1(x)$  and  $(\forall y \in N_1)x \not\prec y$ . It should be  $M_1(x) \geq N_1(x)$ . Nevertheless we do not know whether  $x \in N_1$  or not and cannot prove the contradiction. In the article we referred to [8], [9] and [10].

MSC: 06F05 03B35

Keywords: ordering; Dershowitz-Manna ordering

MML identifier: BAGORD\_2, version: 8.1.04 5.36.1267

## 1. PRELIMINARIES

Now we state the propositions:

- (1) Let us consider natural numbers  $m, n$ . Then  $n = m -' (m -' n) + (n -' m)$ .
- (2) Let us consider natural numbers  $n, m$ . Then  $m -' n \geq m - n$ .

Let us consider natural numbers  $m, n, x, y$ . Now we state the propositions:

- (3) If  $n = m -' x + y$ , then  $m -' n \leq x$  and  $n -' m \leq y$ . The theorem is a consequence of (2).
- (4) If  $x \leq m$  and  $n = m -' x + y$ , then  $x -' (m -' n) = y -' (n -' m)$ . The theorem is a consequence of (3).

Now we state the propositions:

- (5) Let us consider natural numbers  $k, x_1, x_2, y_1, y_2$ . Suppose  $x_2 \leq k$  and  $x_1 \leq k -' x_2 + y_2$ . Then
  - (i)  $x_2 + (x_1 -' y_2) \leq k$ , and

$$(ii) \quad k -' x_2 + y_2 -' x_1 + y_1 = k -' (x_2 + (x_1 -' y_2)) + (y_2 -' x_1 + y_1).$$

PROOF:  $x_2 + (x_1 -' y_2) \leq k$  by [12, (8)].  $\square$

(6) Let us consider natural numbers  $x, y$ . If  $x + y > 0$ , then  $x > 0$  or  $y > 0$ .

From now on  $a, b$  denote objects and  $I, J$  denote sets.

Let us consider  $I$ . Let  $J$  be a non empty set. Let us note that every function from  $I$  into  $J$  is total and there exists a relational structure which is asymmetric, transitive, and non empty.

Let us consider  $I$ . One can verify that there exists a binary relation on  $I$  which is asymmetric and transitive.

Let  $R$  be a transitive relational structure. Observe that the internal relation of  $R$  is transitive.

Let  $R$  be an asymmetric relational structure. Let us observe that the internal relation of  $R$  is asymmetric.

Let us consider  $I$ . Let  $p, q$  be  $I$ -valued finite sequences. Let us observe that  $p \wedge q$  is  $I$ -valued.

Now we state the proposition:

(7) Let us consider finite sequences  $p, q$ . Suppose  $p \wedge q$  is  $I$ -valued. Then

(i)  $p$  is  $I$ -valued, and

(ii)  $q$  is  $I$ -valued.

Let us consider  $I$ . Let  $f$  be an  $I$ -valued finite sequence and  $n$  be a natural number. Let us note that  $f \upharpoonright n$  is  $I$ -valued.

Now we state the propositions:

(8) Let us consider a finite sequence  $p$ . Suppose  $a \in \text{rng } p$ . Then there exist finite sequences  $q, r$  such that  $p = (q \wedge \langle a \rangle) \wedge r$ .

(9) Let us consider finite sequences  $p, q$ . Then  $p \subset q$  if and only if  $\text{len } p < \text{len } q$  and for every natural number  $i$  such that  $i \in \text{dom } p$  holds  $p(i) = q(i)$ .

(10) Let us consider finite sequences  $p, q, r$ . Then  $r \wedge p \subset r \wedge q$  if and only if  $p \subset q$ .

PROOF: If  $r \wedge p \subset r \wedge q$ , then  $p \subset q$  by [4, (22)], (9), [15, (30)], [4, (28)].  $\square$

Let  $R$  be an asymmetric, non empty relational structure and  $x, y$  be elements of  $R$ . Let us observe that the predicate  $x \leq y$  is asymmetric.

Now we state the proposition:

(11) Let us consider an asymmetric, non empty relational structure  $R$ , and elements  $x, y$  of  $R$ . Then  $x \leq y$  if and only if  $x < y$ .

## 2. RELATIONAL EXTENSION

Let us consider  $I$ .

A multiset of  $I$  is an element of  $I^\otimes$ . Observe that every multiset of  $I$  is  $I$ -defined and natural-valued and every multiset of  $I$  is total.

Let  $m$  be a natural-valued function. Let us note that the functor support  $m$  is defined by the term

(Def. 1)  $m^{-1}(\mathbb{N} \setminus \{0\})$ .

Let us consider  $I$ . One can check that every multiset of  $I$  is finite-support.

Now we state the propositions:

(12)  $a$  is a multiset of  $I$  if and only if  $a$  is a bag of  $I$ .

(13)  $1_{I^\otimes} = \text{EmptyBag } I$ .

Let  $R$  be a relational structure and  $x, y$  be elements of  $R$ . We say that  $x \equiv y$  if and only if

(Def. 2)  $x \not\prec y$  and  $y \not\prec x$ .

Observe that the predicate is symmetric.

We consider relational multiplicative magmas which extend multiplicative magmas and relational structures and are systems

$\langle \text{a carrier, a multiplication, an internal relation} \rangle$

where the carrier is a set, the multiplication is a binary operation on the carrier, the internal relation is a binary relation on the carrier.

We consider relational monoids which extend multiplicative loop structures and relational structures and are systems

$\langle \text{a carrier, a multiplication, a one, an internal relation} \rangle$

where the carrier is a set, the multiplication is a binary operation on the carrier, the one is an element of the carrier, the internal relation is a binary relation on the carrier.

Let  $M$  be a multiplicative loop structure.

A relational extension of  $M$  is a relational monoid and is defined by

(Def. 3) the multiplicative loop structure of  $it =$  the multiplicative loop structure of  $M$ .

Let  $M$  be a non empty multiplicative loop structure. Let us observe that every relational extension of  $M$  is non empty.

Let  $M$  be a multiplicative loop structure. One can check that there exists a relational extension of  $M$  which is strict.

Let us consider a multiplicative loop structure  $N$  and a relational extension  $M$  of  $N$ . Now we state the propositions:

(14)  $a$  is an element of  $M$  if and only if  $a$  is an element of  $N$ .

(15)  $1_N = 1_M$ .

Let us consider  $I$ . Let  $M$  be a relational extension of  $I^\otimes$ . Let us observe that every element of  $M$  is function-like and relation-like and every element of  $M$  is  $I$ -defined, natural-valued, and finite-support and every element of  $M$  is total.

Now we state the proposition:

(16) Let us consider a relational extension  $M$  of  $I^\otimes$ . Then the carrier of  $M = \text{Bags } I$ . The theorem is a consequence of (12) and (14).

The scheme *RelEx* deals with a non empty multiplicative loop structure  $\mathcal{M}$  and a binary predicate  $\mathcal{R}$  and states that

(Sch. 1) There exists a strict relational extension  $N$  of  $\mathcal{M}$  such that for every elements  $x, y$  of  $N$ ,  $x \leq y$  iff  $\mathcal{R}[x, y]$ .

Now we state the proposition:

(17) Let us consider a multiplicative loop structure  $N$ , and strict relational extensions  $M_1, M_2$  of  $N$ . Suppose for every elements  $m, n$  of  $M_1$  for every elements  $x, y$  of  $M_2$  such that  $m = x$  and  $n = y$  holds  $m \leq n$  iff  $x \leq y$ . Then  $M_1 = M_2$ .

PROOF: The internal relation of  $M_1 =$  the internal relation of  $M_2$  by [7, (87)].  $\square$

### 3. DERSHOWITZ-MANNA ORDER

Let  $R$  be a non empty relational structure. The Dershowitz-Manna order  $R$  yielding a strict relational extension of  $(\text{the carrier of } R)^\otimes$  is defined by

(Def. 4) for every elements  $m, n$  of  $it$ ,  $m \leq n$  iff there exist elements  $x, y$  of  $it$  such that  $1_{it} \neq x \mid n$  and  $m = n -' x + y$  and for every element  $b$  of  $R$  such that  $y(b) > 0$  there exists an element  $a$  of  $R$  such that  $x(a) > 0$  and  $b \leq a$ .

Now we state the proposition:

(18) Let us consider bags  $m, n$  of  $I$ . Then  $n = m -' (m -' n) + (n -' m)$ . The theorem is a consequence of (1).

Let us consider bags  $m, n, x, y$  of  $I$ . Now we state the propositions:

(19) If  $n = m -' x + y$ , then  $m -' n \mid x$  and  $n -' m \mid y$ . The theorem is a consequence of (3).

(20) If  $x \mid m$  and  $n = m -' x + y$ , then  $x -' (m -' n) = y -' (n -' m)$ . The theorem is a consequence of (4).

Now we state the propositions:

(21) Let us consider bags  $m, x, y$  of  $I$ . If  $x \mid m$  and  $x \neq y$ , then  $m \neq m -' x + y$ .

(22) Let us consider a non empty set  $I$ , a binary relation  $R$  on  $I$ , and a reduction sequence  $r$  w.r.t.  $R$ . If  $\text{len } r > 1$ , then  $r(\text{len } r) \in I$ .

(23) Let us consider an asymmetric, transitive binary relation  $R$  on  $I$ . Then every reduction sequence w.r.t.  $R$  is one-to-one.

PROOF: For every natural numbers  $i, j$  such that  $i > j$  and  $i, j \in \text{dom } r$  holds  $r(i) \neq r(j)$  by [1, (13)], [13, (22)], [1, (11)], [15, (25)].  $\square$

(24) Let us consider an asymmetric, transitive, non empty relational structure  $R$ , and a set  $X$ . Suppose  $X$  is finite and there exists an element  $x$  of  $R$  such that  $x \in X$ . Then there exists an element  $x$  of  $R$  such that  $x$  is maximal in  $X$ .

PROOF: Reconsider  $X_1 = X$  as a finite set. Set  $Y = \{r, \text{ where } r \text{ is an element of } X_1^* : r \text{ is a reduction sequence w.r.t. the internal relation of } R\}$ . Define  $\mathcal{P}[\text{natural number}] \equiv \text{there exists a reduction sequence } r \text{ w.r.t. the internal relation of } R \text{ such that } r \in Y \text{ and } \text{len } r = \$_1$ . For every natural number  $k$  such that  $\mathcal{P}[k]$  holds  $k \leq \overline{X_1}$  by (23), [1, (43)].  $\mathcal{P}[1]$  by [2, (6)], [4, (74), (39)]. Consider  $k$  being a natural number such that  $\mathcal{P}[k]$  and for every natural number  $n$  such that  $\mathcal{P}[n]$  holds  $n \leq k$  from [1, Sch. 6]. Consider  $r$  being a reduction sequence w.r.t. the internal relation of  $R$  such that  $r \in Y$  and  $\text{len } r = k$ . Consider  $q$  being an element of  $X_1^*$  such that  $r = q$  and  $q$  is a reduction sequence w.r.t. the internal relation of  $R$ .  $\square$

(25) Let us consider bags  $m, n$  of  $I$ . Then  $m -' n \mid m$ .

Let us consider  $I$ . Note that every element of Bags  $I$  is function-like and relation-like.

Now we state the proposition:

(26) Let us consider bags  $m, n$  of  $I$ . Then

(i)  $m -' n \neq \text{EmptyBag } I$ , or

(ii)  $m = n$ , or

(iii)  $n -' m \neq \text{EmptyBag } I$ .

Let  $R$  be an asymmetric, transitive, non empty relational structure. Let us observe that the Dershowitz-Manna order  $R$  is defined by

(Def. 5) for every elements  $m, n$  of  $it$ ,  $m \leq n$  iff  $m \neq n$  and for every element  $a$  of  $R$  such that  $m(a) > n(a)$  there exists an element  $b$  of  $R$  such that  $a \leq b$  and  $m(b) < n(b)$ .

Now we state the proposition:

(27) Let us consider bags  $k, x_1, x_2, y_1, y_2$  of  $I$ . Suppose  $x_2 \mid k$  and  $x_1 \mid k -' x_2 + y_2$ . Then

- (i)  $x_2 + (x_1 -' y_2) \mid k$ , and
- (ii)  $k -' x_2 + y_2 -' x_1 + y_1 = k -' (x_2 + (x_1 -' y_2)) + (y_2 -' x_1 + y_1)$ .

The theorem is a consequence of (5).

Let  $R$  be an asymmetric, transitive, non empty relational structure. Let us observe that the Dershowitz-Manna order  $R$  is asymmetric and transitive.

Let us consider  $I$ . The functor  $\text{DivOrder}(I)$  yielding a binary relation on  $\text{Bags } I$  is defined by

(Def. 6) for every bags  $b_1, b_2$  of  $I$ ,  $\langle b_1, b_2 \rangle \in it$  iff  $b_1 \neq b_2$  and  $b_1 \mid b_2$ .

Now we state the proposition:

(28) Let us consider bags  $a, b, c$  of  $I$ . If  $a \mid b \mid c$ , then  $a \mid c$ .

Let us consider  $I$ . Note that  $\text{DivOrder}(I)$  is asymmetric and transitive.

Let us consider an asymmetric, transitive, non empty relational structure  $R$ . Now we state the propositions:

(29)  $\text{DivOrder}(\text{the carrier of } R) \subseteq$  the internal relation of the Dershowitz-Manna order  $R$ . The theorem is a consequence of (12) and (14).

(30) Suppose the internal relation of  $R$  is empty. Then the internal relation of the Dershowitz-Manna order  $R = \text{DivOrder}(\text{the carrier of } R)$ . The theorem is a consequence of (29).

Now we state the proposition:

(31) Let us consider asymmetric, transitive, non empty relational structures  $R_1, R_2$ . Suppose the carrier of  $R_1 =$  the carrier of  $R_2$  and the internal relation of  $R_1 \subseteq$  the internal relation of  $R_2$ . Then the internal relation of the Dershowitz-Manna order  $R_1 \subseteq$  the internal relation of the Dershowitz-Manna order  $R_2$ . The theorem is a consequence of (12) and (14).

#### 4. MONOIDAL ORDER

Let us consider  $I$ . Let  $f$  be a  $(\text{Bags } I)$ -valued finite sequence. The functor  $\sum f$  yielding a bag of  $I$  is defined by

(Def. 7) there exists a function  $F$  from  $\mathbb{N}$  into  $\text{Bags } I$  such that  $it = F(\text{len } f)$  and  $F(0) = \text{EmptyBag } I$  and for every natural number  $i$  and for every bag  $b$  of  $I$  such that  $i < \text{len } f$  and  $b = f(i + 1)$  holds  $F(i + 1) = F(i) + b$ .

Now we state the proposition:

(32)  $\sum \varepsilon_{\text{Bags } I} = \text{EmptyBag } I$ .

Let us consider  $I$ . Let  $b$  be a bag of  $I$ . One can verify that  $\langle b \rangle$  is  $(\text{Bags } I)$ -valued as a finite sequence.

Now we state the proposition:

(33) Let us consider a (Bags  $I$ )-valued finite sequence  $p$ , and a bag  $b$  of  $I$ . Then  $\sum(p \wedge \langle b \rangle) = \sum p + b$ .

PROOF: Set  $f = p \wedge \langle b \rangle$ . Consider  $F$  being a function from  $\mathbb{N}$  into Bags  $I$  such that  $\sum f = F(\text{len } f)$  and  $F(0) = \text{EmptyBag } I$  and for every natural number  $i$  and for every bag  $b$  of  $I$  such that  $i < \text{len } f$  and  $b = f(i + 1)$  holds  $F(i + 1) = F(i) + b$ . Consider  $F_1$  being a function from  $\mathbb{N}$  into Bags  $I$  such that  $\sum p = F_1(\text{len } p)$  and  $F_1(0) = \text{EmptyBag } I$  and for every natural number  $i$  and for every bag  $b$  of  $I$  such that  $i < \text{len } p$  and  $b = p(i + 1)$  holds  $F_1(i + 1) = F_1(i) + b$ . Define  $\mathcal{P}[\text{natural number}] \equiv$  if  $\$1 \leq \text{len } p$ , then  $F(\$1) = F_1(\$1)$ . For every natural number  $i$  such that  $\mathcal{P}[i]$  holds  $\mathcal{P}[i + 1]$  by [5, (16)], [1, (13), (11)], [15, (25)]. For every natural number  $i$ ,  $\mathcal{P}[i]$  from [1, Sch. 2].  $\square$

From now on  $b$  denotes a bag of  $I$ .

Now we state the propositions:

(34)  $\sum \langle b \rangle = b$ . The theorem is a consequence of (33) and (32).

(35) Let us consider (Bags  $I$ )-valued finite sequences  $p, q$ . Then  $\sum(p \wedge q) = \sum p + \sum q$ .

PROOF: Set  $f = p \wedge q$ . Consider  $F$  being a function from  $\mathbb{N}$  into Bags  $I$  such that  $\sum f = F(\text{len } f)$  and  $F(0) = \text{EmptyBag } I$  and for every natural number  $i$  and for every bag  $b$  of  $I$  such that  $i < \text{len } f$  and  $b = f(i + 1)$  holds  $F(i + 1) = F(i) + b$ . Consider  $F_1$  being a function from  $\mathbb{N}$  into Bags  $I$  such that  $\sum p = F_1(\text{len } p)$  and  $F_1(0) = \text{EmptyBag } I$  and for every natural number  $i$  and for every bag  $b$  of  $I$  such that  $i < \text{len } p$  and  $b = p(i + 1)$  holds  $F_1(i + 1) = F_1(i) + b$ . Consider  $F_2$  being a function from  $\mathbb{N}$  into Bags  $I$  such that  $\sum q = F_2(\text{len } q)$  and  $F_2(0) = \text{EmptyBag } I$  and for every natural number  $i$  and for every bag  $b$  of  $I$  such that  $i < \text{len } q$  and  $b = q(i + 1)$  holds  $F_2(i + 1) = F_2(i) + b$ . Define  $\mathcal{P}[\text{natural number}] \equiv$  if  $\$1 \leq \text{len } p$ , then  $F(\$1) = F_1(\$1)$ . For every natural number  $i$  such that  $\mathcal{P}[i]$  holds  $\mathcal{P}[i + 1]$  by [4, (22)], [1, (11), (13)], [15, (25)]. For every natural number  $i$ ,  $\mathcal{P}[i]$  from [1, Sch. 2]. Define  $\mathcal{Q}[\text{natural number}] \equiv$  if  $\$1 \leq \text{len } q$ , then  $F(\text{len } p + \$1) = \sum p + F_2(\$1)$ . For every natural number  $i$  such that  $\mathcal{Q}[i]$  holds  $\mathcal{Q}[i + 1]$  by [4, (22)], [1, (13), (11)], [15, (25)]. For every natural number  $i$ ,  $\mathcal{Q}[i]$  from [1, Sch. 2].  $\square$

Let us consider a (Bags  $I$ )-valued finite sequence  $p$ . Now we state the propositions:

(36)  $\sum(\langle b \rangle \wedge p) = b + \sum p$ . The theorem is a consequence of (35) and (34).

(37) If  $b \in \text{rng } p$ , then  $b \mid \sum p$ . The theorem is a consequence of (8), (7), (33), and (35).

Now we state the proposition:

(38) Let us consider a (Bags  $I$ )-valued finite sequence  $p$ , and an object  $i$ . Suppose  $i \in \text{support } \sum p$ . Then there exists  $b$  such that

- (i)  $b \in \text{rng } p$ , and
- (ii)  $i \in \text{support } b$ .

PROOF: Define  $\mathcal{P}[\text{natural number}] \equiv$  for every (Bags  $I$ )-valued finite sequence  $p$  such that  $\text{len } p = \$1$  for every object  $i$  such that  $i \in \text{support } \sum p$  there exists  $b$  such that  $b \in \text{rng } p$  and  $i \in \text{support } b$ .  $\mathcal{P}[0]$ . For every natural number  $j$  such that  $\mathcal{P}[j]$  holds  $\mathcal{P}[j + 1]$  by [3, (3)], (7), [4, (40)], [15, (25)]. For every natural number  $j$ ,  $\mathcal{P}[j]$  from [1, Sch. 2].  $\square$

Let us consider  $I$  and  $b$ .

A partition of  $b$  is a (Bags  $I$ )-valued finite sequence and is defined by

(Def. 8)  $b = \sum it$ .

Observe that the functor  $\langle b \rangle$  yields a partition of  $b$ . Let  $R$  be a relational structure,  $M$  be a relational extension of (the carrier of  $R$ ) $^\otimes$ ,  $b$  be an element of  $M$ , and  $p$  be a partition of  $b$ . We say that  $p$  is co-ordered if and only if

(Def. 9) for every natural number  $i$  such that  $i, i + 1 \in \text{dom } p$  for every elements  $b_1, b_2$  of  $M$  such that  $b_1 = p(i)$  and  $b_2 = p(i + 1)$  holds  $b_2 \leq b_1$ .

Let  $R$  be a non empty relational structure and  $b$  be a bag of the carrier of  $R$ . We say that  $p$  is ordered if and only if

(Def. 10) for every bag  $m$  of the carrier of  $R$  such that  $m \in \text{rng } p$  for every element  $x$  of  $R$  such that  $m(x) > 0$  holds  $m(x) = b(x)$  and for every bag  $m$  of the carrier of  $R$  such that  $m \in \text{rng } p$  for every elements  $x, y$  of  $R$  such that  $m(x) > 0$  and  $m(y) > 0$  and  $x \neq y$  holds  $x \equiv y$  and for every bag  $m$  of the carrier of  $R$  such that  $m \in \text{rng } p$  holds  $m \neq \text{EmptyBag}(\text{the carrier of } R)$  and for every natural number  $i$  such that  $i, i + 1 \in \text{dom } p$  for every element  $x$  of  $R$  such that  $p_{i+1}(x) > 0$  there exists an element  $y$  of  $R$  such that  $p_i(y) > 0$  and  $x \leq y$ .

In the sequel  $R$  denotes an asymmetric, transitive, non empty relational structure,  $a, b, c$  denote bags of the carrier of  $R$ , and  $x, y, z$  denote elements of  $R$ .

Now we state the propositions:

(39)  $\langle a \rangle$  is ordered if and only if  $a \neq \text{EmptyBag}(\text{the carrier of } R)$  and for every  $x$  and  $y$  such that  $a(x) > 0$  and  $a(y) > 0$  and  $x \neq y$  holds  $x \equiv y$ .

(40) Let us consider a (Bags  $I$ )-valued finite sequence  $p$ , and bags  $a, b$  of  $I$ . Then  $\langle a \rangle \wedge p$  is a partition of  $b$  if and only if  $a \mid b$  and  $p$  is a partition of  $b -' a$ . The theorem is a consequence of (36).

From now on  $p$  denotes a partition of  $b -' a$  and  $q$  denotes a partition of  $b$ .

Now we state the proposition:



(41) If  $q = \langle a \rangle \wedge p$  and  $q$  is ordered, then  $p$  is ordered. The theorem is a consequence of (37) and (25).

Let us consider  $I$ . Let  $m$  be a bag of  $I$  and  $J$  be a set. The functor  $m \upharpoonright J$  yielding a bag of  $I$  is defined by

(Def. 11) for every object  $i$  such that  $i \in I$  holds if  $i \in J$ , then  $it(i) = m(i)$  and if  $i \notin J$ , then  $it(i) = 0$ .

From now on  $J$  denotes a set and  $m$  denotes a bag of  $I$ .

Now we state the propositions:

(42)  $\text{support}(m \upharpoonright J) = J \cap \text{support } m$ .

(43)  $m \upharpoonright J + m \upharpoonright (I \setminus J) = m$ .

(44)  $m \upharpoonright J \mid m$ .

(45) If  $\text{support } m \subseteq J$ , then  $m \upharpoonright J = m$ .

(46)  $\text{support}(m -' m \upharpoonright J) = \text{support } m \setminus J$ .

(47) If  $q$  is ordered and  $q = \langle a \rangle \wedge p$  and  $a(x) > 0$ , then  $a(x) = b(x)$ .

(48) If  $q$  is ordered and  $q = \langle a \rangle \wedge p$  and  $a(x) > 0$  and  $a(y) > 0$  and  $x \neq y$ , then  $x \equiv y$ .

(49) If  $q$  is ordered and  $q = \langle a \rangle \wedge p$ , then  $a \neq \text{EmptyBag}(\text{the carrier of } R)$ .

(50) Let us consider a bag  $c$  of the carrier of  $R$ , and a  $(\text{Bags}(\text{the carrier of } R))$ -valued finite sequence  $r$ . Suppose  $q$  is ordered and  $q = \langle a, c \rangle \wedge r$  and  $c(y) > 0$ . Then there exists  $x$  such that

(i)  $a(x) > 0$ , and

(ii)  $y \leq x$ .

(51) If  $x \in I$  and for every  $y$  such that  $y \in I$  and  $y \neq x$  holds  $x \equiv y$ , then  $x$  is maximal in  $I$ .

(52) If  $q$  is ordered and  $q = \langle a \rangle \wedge p$  and  $c \in \text{rng } p$  and  $c(x) > 0$ , then there exists  $y$  such that  $a(y) > 0$  and  $x \leq y$ .

PROOF: Consider  $i$  being an object such that  $i \in \text{dom } p$  and  $c = p(i)$ . Define  $\mathcal{P}[\text{natural number}] \equiv$  if  $\$1 \in \text{dom } p$ , then for every  $x$  such that  $p_{\$1}(x) > 0$  there exists  $y$  such that  $a(y) > 0$  and  $x \leq y$ .  $\mathcal{P}[1]$  by [4, (28)], [15, (25)], [4, (40)]. For every natural number  $i$  such that  $i \geq 1$  and  $\mathcal{P}[i]$  holds  $\mathcal{P}[i + 1]$  by [1, (13)], [15, (25)], [4, (28)], [16, (3)]. For every natural number  $i$  such that  $i \geq 1$  holds  $\mathcal{P}[i]$  from [1, Sch. 8].  $\square$

Let us assume that  $q$  is ordered and  $q = \langle a \rangle \wedge p$ . Now we state the propositions:

(53)  $x$  is maximal in  $\text{support } b$  if and only if  $a(x) > 0$ .

PROOF:  $a \mid \sum q = b$ . There exists no  $y$  such that  $y \in \text{support } b$  and  $x < y$  by (48), (38), [4, (31), (39)].  $\square$

(54)  $a = b \upharpoonright \{x : x \text{ is maximal in support } b\}$ . The theorem is a consequence of (53) and (47).

Now we state the propositions:

(55) Let us consider a  $(\text{Bags } I)$ -valued finite sequence  $p$ . Suppose  $\sum p = \text{EmptyBag } I$  and for every bag  $a$  of  $I$  such that  $a \in \text{rng } p$  holds  $a \neq \text{EmptyBag } I$ . Then  $p = \emptyset$ . The theorem is a consequence of (37).

(56) Let us consider bags  $a, b$  of  $I$ . If  $a \neq \text{EmptyBag } I$ , then  $a + b \neq \text{EmptyBag } I$ .

(57) Let us consider partitions  $p, q$  of  $b$ . If  $p$  is ordered and  $q$  is ordered, then  $p = q$ .

PROOF: Define  $\mathcal{P}[\text{natural number}] \equiv$  for every  $b$  and  $q$  such that  $\text{len } q = \$_1$  and  $q$  is ordered for every partition  $p$  of  $b$  such that  $p$  is ordered holds  $q = p$ .  $\mathcal{P}[0]$ . For every natural number  $i$  such that  $\mathcal{P}[i]$  holds  $\mathcal{P}[i+1]$  by [5, (130)], (40), (49), (36). For every natural number  $i$ ,  $\mathcal{P}[i]$  from [1, Sch. 2].  $\square$

Let us consider  $I$ . Let  $a, b$  be bags of  $I$ . One can verify that the functor  $\langle a, b \rangle$  yields an element of  $\text{Bags } I \times \text{Bags } I$ . Now we state the proposition:

(58) Suppose  $a \neq \text{EmptyBag}(\text{the carrier of } R)$ . Then  $\{x : x \text{ is maximal in support } a\} \neq \emptyset$ . The theorem is a consequence of (24).

Let us consider  $R$  and  $b$ . The ordered partition of  $b$  yielding a  $(\text{Bags}(\text{the carrier of } R))$ -valued finite sequence is defined by

(Def. 12) there exist functions  $F, G$  from  $\mathbb{N}$  into  $\text{Bags}(\text{the carrier of } R)$  such that  $F(0) = b$  and  $G(0) = \text{EmptyBag}(\text{the carrier of } R)$  and for every natural number  $i$ ,  $G(i+1) = F(i) \upharpoonright \{x : x \text{ is maximal in support}(F(i))\}$  and  $F(i+1) = F(i) -' G(i+1)$  and there exists a natural number  $i$  such that  $F(i) = \text{EmptyBag}(\text{the carrier of } R)$  and  $it = G \upharpoonright \text{Seg } i$  and for every natural number  $j$  such that  $j < i$  holds  $F(j) \neq \text{EmptyBag}(\text{the carrier of } R)$ .

One can verify that the ordered partition of  $b$  yields a partition of  $b$ . Let us note that the ordered partition of  $b$  is ordered as a partition of  $b$ .

Now we state the proposition:

(59)  $b = \text{EmptyBag}(\text{the carrier of } R)$  if and only if the ordered partition of  $b = \emptyset$ . The theorem is a consequence of (32).

Let us consider  $R$ . The functor  $\prec_{\mathcal{M}} R$  yielding a strict relational extension of  $(\text{the carrier of } R)^{\otimes}$  is defined by

(Def. 13) for every elements  $m, n$  of  $it$ ,  $m \leq n$  iff  $m \neq n$  and for every  $x$  such that  $m(x) > 0$  holds  $m(x) < n(x)$  or there exists  $y$  such that  $n(y) > 0$  and  $x \leq y$ .

Let us note that  $\prec_{\mathcal{M}} R$  is asymmetric and transitive.

Let us consider  $I$ . Let  $R$  be a relation between  $I$  and  $I$ .

The functor  $\text{LexOrder}(I, R)$  yielding a binary relation on  $I^*$  is defined by

(Def. 14) for every  $I$ -valued finite sequences  $p, q$ ,  $\langle p, q \rangle \in it$  iff  $p \subset q$  or there exists a natural number  $k$  such that  $k \in \text{dom } p$  and  $k \in \text{dom } q$  and  $\langle p(k), q(k) \rangle \in R$  and for every natural number  $n$  such that  $1 \leq n < k$  holds  $p(n) = q(n)$ .

Let  $R$  be a transitive binary relation on  $I$ . One can verify that  $\text{LexOrder}(I, R)$  is transitive.

Let  $R$  be an asymmetric binary relation on  $I$ . Note that  $\text{LexOrder}(I, R)$  is asymmetric.

Now we state the proposition:

(60) Let us consider an asymmetric binary relation  $R$  on  $I$ , and  $I$ -valued finite sequences  $p, q, r$ . Then  $\langle p, q \rangle \in \text{LexOrder}(I, R)$  if and only if  $\langle r \hat{\ } p, r \hat{\ } q \rangle \in \text{LexOrder}(I, R)$ . The theorem is a consequence of (10).

Let us consider  $R$ . The functor  $\prec\prec_{\mathcal{M}} R$  yielding a strict relational extension of (the carrier of  $R$ )<sup>⊗</sup> is defined by

(Def. 15) for every elements  $m, n$  of  $it$ ,  $m \leq n$  iff  $\langle$ the ordered partition of  $m$ , the ordered partition of  $n \rangle \in \text{LexOrder}(\langle$ the carrier of  $\prec\prec_{\mathcal{M}} R$  $\rangle, \langle$ the internal relation of  $\prec\prec_{\mathcal{M}} R$  $\rangle)$ .

Observe that  $\prec\prec_{\mathcal{M}} R$  is asymmetric and transitive.

Now we state the propositions:

(61) Let us consider elements  $a, b$  of the Dershowitz-Manna order  $R$ . Suppose  $a \leq b$ . Then  $b \neq \text{EmptyBag}(\text{the carrier of } R)$ . The theorem is a consequence of (29).

(62) Let us consider elements  $a, b, c, d$  of the Dershowitz-Manna order  $R$ , and a bag  $e$  of the carrier of  $R$ . Suppose  $a \leq b$  and  $e \mid a$  and  $e \mid b$ . If  $c = a -' e$  and  $d = b -' e$ , then  $c \leq d$ .

(63) Let us consider a (Bags  $I$ )-valued finite sequence  $p$ , and an object  $x$ . Suppose  $x \in I$  and  $(\sum p)(x) > 0$ . Then there exists a natural number  $i$  such that

- (i)  $i \in \text{dom } p$ , and
- (ii)  $p_i(x) > 0$ .

PROOF: Define  $\mathcal{P}[\text{object}] \equiv$  for every (Bags  $I$ )-valued finite sequence  $p$  such that  $p = \$_1$  and  $(\sum p)(x) > 0$  there exists a natural number  $i$  such that  $i \in \text{dom } p$  and  $p_i(x) > 0$ .  $\mathcal{P}[\emptyset]$  by (32), [14, (7)]. For every finite sequence  $p$  and for every object  $a$  such that  $\mathcal{P}[p]$  holds  $\mathcal{P}[p \hat{\ } \langle a \rangle]$  by (7), [4, (40)], [15, (25)], [6, (102)]. For every finite sequence  $p$ ,  $\mathcal{P}[p]$  from [4, Sch. 3].  $\square$

(64) If  $q$  is ordered and  $q_1(x) = 0$  and  $b(x) > 0$ , then there exists  $y$  such that  $q_1(y) > 0$  and  $x \leq y$ .

PROOF: Define  $\mathcal{P}[\text{natural number}] \equiv$  if  $\$1 \in \text{dom } q$ , then for every  $x$  such that  $q_{\$1}(x) > 0$  there exists  $y$  such that  $q_1(y) > 0$  and  $x \leq y$ .  $\mathcal{P}[2]$  by [15, (25)]. For every natural number  $i$  such that  $2 \leq i$  and  $\mathcal{P}[i]$  holds  $\mathcal{P}[i+1]$  by [1, (11)], [15, (25)], [16, (3)]. For every natural number  $i$  such that  $i \geq 2$  holds  $\mathcal{P}[i]$  from [1, Sch. 8]. Consider  $i$  being a natural number such that  $i \in \text{dom } q$  and  $q_i(x) > 0$ .  $\square$

## REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [2] Grzegorz Bancerek. Reduction relations. *Formalized Mathematics*, 5(4):469–478, 1996.
- [3] Grzegorz Bancerek. König’s lemma. *Formalized Mathematics*, 2(3):397–402, 1991.
- [4] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [5] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [6] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [7] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [8] Nachum Dershowitz. Orderings for term-rewriting systems. *Theoretical Computer Science*, 17(3):279–301, 1982. doi:10.1016/0304-3975(82)90026-3.
- [9] Nachum Dershowitz and Zohar Manna. Proving termination with multiset orderings. *Communications of the ACM*, 22(8):465–476, 1979. doi:10.1145/359138.359142.
- [10] Gerard Huet and Derek C. Oppen. Equations and rewrite rules: A survey. Technical report, Stanford, CA, USA, 1980.
- [11] Jean-Pierre Jouannaud and Pierre Lescanne. On multiset ordering. *Information Processing Letters*, 15(2):57–63, 1982. doi:10.1016/0020-0190(82)90107-7.
- [12] Robert Milewski. Natural numbers. *Formalized Mathematics*, 7(1):19–22, 1998.
- [13] Eliza Niewiadomska and Adam Grabowski. Introduction to formal preference spaces. *Formalized Mathematics*, 21(3):223–233, 2013. doi:10.2478/forma-2013-0024.
- [14] Andrzej Trybulec. Binary operations applied to functions. *Formalized Mathematics*, 1(2):329–334, 1990.
- [15] Wojciech A. Trybulec. Non-contiguous substrings and one-to-one finite sequences. *Formalized Mathematics*, 1(3):569–573, 1990.
- [16] Wojciech A. Trybulec and Grzegorz Bancerek. Kuratowski – Zorn lemma. *Formalized Mathematics*, 1(2):387–393, 1990.

*Received December 31, 2015*