

Oceanological and Hydrobiological Studies

International Journal of Oceanography and Hydrobiology

Volume 42, Issue 3

ISSN 1730-413X
eISSN 1897-3191

(347–357)
2013



DOI: 10.2478/s13545-013-0092-9
Review paper

Received: December 14, 2012
Accepted: March 19, 2013

A review of cnidarian epibionts on marine crustacea

Gregorio Fernandez-Leborans*

*Department of Zoology, Faculty of Biology,
Pnta9, Complutense University, 28040 Madrid, Spain*

Key words: epibiosis, cnidaria, crustacea

Abstract

An updated inventory of the cnidarian species living as epibionts on Crustacea was conducted. Cnidarian species that attach themselves to gastropod shells of hermit crabs were also considered. One hundred and forty-eight species of cnidaria were included, with similar numbers of hydrozoans and anthozoans. There were 163 basibiont species that were mostly decapod species. The presence of these epibionts on different crustacean species and their taxonomic positions were indicated. This review is intended to serve as a useful tool for future studies of this type of epibiosis.

INTRODUCTION

Epibiosis is the presence of life forms on the surface of other organisms (Wahl 1989). In many cases the form that colonises, i.e. the epibiont, can be an organism with a sessile phase of its life cycle. The organism on which the epibiont settles is referred to as a basibiont, which provides the attachment surface

and is generally larger than the epibiont (Fernandez-Leborans, Cárdenas 2009). This phenomenon is very common in aquatic environments, where hard substrates available for sessile forms are limited (Wahl 1989, Gili et al. 1993). The existence of epibiosis implies ecological interactions between the partners. In aquatic environments, many crustacean groups, such as cladocerans, copepods, cirripedes, isopods, amphipods and decapods, serve as basibionts for protozoan micro-epibionts and invertebrate macro-epibionts (Ross 1983; Corliss 1979; Fernandez-Leborans, Tato-Porto 2000b). Epibiosis is the evolutionary effect of interactions between environmental factors and benthic life forms (Key et al. 1999). It is a dynamic process, and the benefits and disadvantages for the organisms involved vary depending on the environmental conditions (Bush et al. 2001).

Although, in principle, the epibiosis is not necessarily obligatory, and it does not meet all the requirements for the survival of each species, except for the attachment to a substrate, there may be several effects for both the epibiont and the basibiont. The symbiotic relationship could be either commensalism, mutualism, or parasitism. The epibiosis may produce extensive evolutionary changes in the partners; the effects can include advantages for the epibiont in terms of dispersal and geographical expansion (Gili et al. 1993, Threlkeld et al. 1993), an increase in the nutrient supply, and protection against predation or negative environmental conditions (Abelló et al. 1990). Epibiosis can also be disadvantageous to the epibiont by inducing ontogenetic or behavioural changes (Threlkeld et al. 1993). For example, epibiosis can be beneficial for the basibiont by providing both mimetic protection and cleaning (Ingle 1983; Maldonado, Uriz 1992). In contrast, epibiosis may be negative for the basibiont by restricting its mobility

* e-mail address: greg@bio.ucm.es

and affecting its growth, moulting and functioning of several organs (the eyes, gills, and appendages) (Overstreet 1983, 1987; Reaka 1978). Epibiosis may cause an increased risk of predation (Willey et al. 1990). Epibionts and basibionts may compete for nutrients (Fernandez-Leborans et al. 2006).

Cnidarians are one of the most frequent epibionts on crustaceans. Cnidarian species are frequent on copepods, cirripeds, isopods and decapods. Different aspects of the epibiotic relationships between cnidarians and crustaceans have been studied, i.a. the presence and spatial distribution of epibionts, interactions between the basibiont and the epibiont, patterns of colonisation, effects of abiotic parameters, and the influence on the life cycles and physiology of the organisms involved.

The comparison of presence/occurrence of cnidaria and other epibionts

Cnidaria have frequently been found with other epibionts of diverse crustacean species groups. Protozoan, molluscs, annelids, other crustacean, and bryozoa, etc. commonly share the hard substrates corresponding to the exoskeleton of crustacea (Fernandez-Leborans, Tato-Porto 2000). One of the most studied crustaceans, the hermit crab *Pagurus bernhardus*, may harbour 51 epibiont species belonging to 11 groups (foraminifers, cnidaria, platyhelminthes, nemertines, molluscs, polychaetes, crustacea, nematodes, foronids, bryozoa and tunicata) (Reiss et al. 2003).

Different species from different groups of organisms have been found as epibionts on the shell-crab complex of hermit crabs. The shells and crabs may present different epibionts. Cnidaria, especially hydrozoans, can in many cases dominate a large portion of the surfaces. Hydrozoans are frequently located on areas surrounding the aperture of the shell or define the edges of the apex of the shell's aperture. This location appears to be most effective for the epibiont to use the nutrient particles from crab's activities. Other epibionts, such as ciliate protozoans, cirripeds and polychaetes, are distributed on shell surfaces without hydrozoans. In many cases, the characteristics of the hydrozoan colonies determine an extensive colonisation of the basibiont surface, which is favored by the structure or the base of the colony and the development of zooids. This fact leads several epibiont species to follow an adaptation process for living as hyperepibionts on hydrozoan colonies. Bacteria, microalgae, foraminifera, and

polychaetes have been observed on hydrozoans (Bavestrello et al. 2008) and also some ciliate protozoan species, such as the peritrich ciliate *Vorticella*, the suctorian *Ephelota*, the hypotrich ciliate *Kerona pediculus* (O.F. Müller, 1773), and the lcnophorid *Lcnophora chattoni* Villeneuve-Brachon, 1939 (Bavestrello et al. 2008; Silva-Neto et al. 2012; Warren, Robson 1998). Some of the hydrozoan epibionts, such as *Hydractinia echinata* (Fleming, 1828), are obligate epibionts that account for the highest percentage of organisms on the basibiont surface (Reiss et al. 2003).

The coexistence between hydrozoans and other epibionts can be observed in other crustaceans, such as the lithoded decapod *Lithodes maja* (Linnaeus, 1758) (as *L. maia*). This decapod may have epibionts from diverse groups together with cnidarian epibionts. Two hydroid species of the genus *Obelia* have been found on this crab along with 4 polychaete species, 3 bivalves, 2 amphipods, 2 cirripeds, 3 bryozoans and 1 leech species. In this case, hydrozoan species were also the most abundant species. As in other cases of epibiosis on crustaceans, the structure of the carapace and the size of the basibiont affected the presence and the number of different epibionts (Dvoretzky, Dvoretzky 2008).

In the western areas of Scotland, the crab *Necora puber* (as *Liocarcinus puber*), together with the epibiont hydrozoans *Chytia* sp. and *Leuckartiara* sp., has been shown to present the polychaetes *Pomatoceros triqueter* (Linnaeus, 1767) and *Hydroides norvegica* Gunnerus, 1768, the cirriped *Balanus crenatus* Bruguière, 1789, the entoproct *Barentsia matsushimana* Toriumi, 1951, the ciliate protozoans *Ephelota plana* Annandale, 1911, *Ephelota gemmipara* (Hertwig, 1875), *Chilodochona quennerstedti* Wallengren, 1895 and *Cothurnia longipes* Kahl, 1935 (Fernandez-Leborans, Gabilondo 2008). Furthermore, hydrozoans have been found on the carapace, chelipeds and anterior periopods of another related species of decapod crustaceans, i.e. *Liocarcinus depurator*. These hydrozoans coexist with the suctorian ciliate, which has been found on the same areas as the hydrozoan (Fernandez-Leborans, Gabilondo 2005).

The coexistence of the elllobiopsis *Thalassomyces* sp. and the hydroid *Earleria corachloae* Widmer, Cailliet & Geller, 2010 on the shrimp *Pasiphaea pacifica* Rathbun, 1902 from Monterey Bay (California) was described (Widmer et al. 2010). *Thalassomyces* is a parasitic alveolate protozoan with a nutrient-absorbing "root" and has been documented to exist on diverse crustaceans, including a wide array of

pelagic marine malacostracan crustaceans (shrimp, euphausiids, mysids, amphipods, and copepods) (Silberman et al. 2004). The presence of the ellobiopsid and the hydroid on the shrimp carapace did not seem to affect the swimming behaviour of the shrimp when compared with an uninfected shrimp (Widmer et al. 2010).

The epibiont communities described on two species of the decapods *Cancer* (*C. gracilis* Dana, 1852, *C. productus* J. W. Randall, 1840 and *C. magister* Dana, 1852) included 29 epibiont species from ten phyla. The hydrozoan colonies have been observed on the ventral surfaces and limbs, with the cirripeds as the most abundant ones and the tube-dwelling polychaetes being less common. Other organisms, such as algae, bryozoa, mollusks, porifera and tunicates, have been found on these crabs sampled in Barkley Sound, British Columbia (McGaw 2006).

The spatial distribution of species in these epibiotic communities, in which cnidaria may be present, can be considered in several ways:

- 1) on each basibiont, the colonisation can follow a selection of different habitats on the surface of the basibiont;
- 2) many species of epibionts show a particular pattern of distribution, which may correspond to, as appropriate, the concepts of aggregation, overlap, exclusion, and association (Hayashi, Tsuji 2007);
- 3) epibiont communities appear to behave as a whole, regardless of particular species, which can be verified by the colonisation process;
- 4) the epibiont species tend to occupy the available substrate within the particular requirements of each functional group but also tend to achieve the equilibrium among species and groups, compensating for diversity and density (Fernandez-Leborans et al. 2006);
- 5) as it was evidenced in the study of the epibiont communities of the shrimp *Caridina* from three lakes in Indonesia, the basibiont represents a dynamic environment in which species of the epibiont community develop a colonisation pattern.

The species were located on the host in a particular pattern, which is shown in the results: the species distribution patterns were associated with different lakes. Regardless of the species present, and in all cases, each species was found to follow the same general pattern of distribution (Fernandez-

Leborans et al. 2006). The distribution of epibionts on the surface of basibionts is determined by the following factors:

- 1) biotic and abiotic microlocal parameters;
- 2) water currents generated by feeding and respiratory activities of the basibiont;
- 3) availability of nutrients (Threlkeld et al. 1993);
- 4) behaviour of the basibiont (e.g. burying in sediment, copulation, emersion, nocturnal activity, hiding under rocks);
- 5) the age, sex, body shape, and size of the host (Connell & Keough 1985; Key et al. 1999);
- 6) moult and intermoult periods;
- 7) the presence of particular chemical compounds;
- 8) the presence of protrusions and rugosities on the surface of the basibiont (Viljoen, Van As 1983) and other differences according to particular surfaces of the basibiont;
- 9) mucosity and sediment on the host surface preventing the adhesion of epibionts (Mulisch et al. 1986);
- 10) the presence of a bacterial community and the associated ciliate protozoan fauna on the basibiont surface (Fernández-Leborans, Herrero Córdoba 1997).

In the present revision, 159 crustacean species (including the genera) were recognized as basibionts for 189 species of cnidarian epibionts (Appendix 1), including decapods as the most frequently (92.02%) reported species, while hydrozoans and anthozoans had similar proportions (55.97% and 44.03%, respectively).

There were several reasons for conducting this review, including the following ones:

- 1) Although there are numerous academic contributions regarding the presence of cnidarian epibionts in crustaceans, there are no reviews of different species of cnidarian epibionts and the corresponding species of crustacean basibionts.
- 2) Cnidarians in many cases represent the largest fraction of the epibiont community in crustaceans, both in terms of biomass and the number of individuals.
- 3) Both the diversity of cnidarian epibionts and the crustacean basibiont species in which the former are found are remarkable.
- 4) The cnidarian-crustacean epibiosis has many singular aspects that are related to interactions

between both types of organisms (colonisation, behaviour, the choice of epibiont and basibiont, specificity, defence, predation, feeding, life cycles, synchronisation, physiology, mechanisms antifouling, etc.).

This review provides not only the idea of species diversity of cnidarians and crustaceans involved in the epibiosis but also the relative importance of different taxonomic groups of cnidarians and crustaceans as epibionts. The differences may be related to particular biological characteristics of each party in the community and the characteristics of the whole community in epibiosis and its environment.

REFERENCES

- Abelló, P. & Macpherson E. (1992). Epibiosis and rhizocephalan infestation patterns in relation to the reproductive biology of *Lithodes ferox* Filhol, 1885 (Anomura: Lithodidae). *J. Crust. Biol.* 12, 561–570.
- Abelló, P., Villanueva R. & Gili J.M. (1990). Epibiosis in deep-sea crab populations as indicator of biological and behavioural characteristics of the host. *J. Mar. Biol. Assoc. U. K.* 70, 687–695.
- Acuña, F., Excoffon A, & Scelzo M. (2003). Mutualism between the sea anemone *Antholoba achates* (Drayton, 1846) (Cnidaria: Actiniaria: Actinostolidae) and the spider crab *Libinia spinosa* Milne-Edwards, 1834 (Crustacea: Decapoda, Majidae). *Belgian J. Zool.* 133; 85 - 87.
- Arai, M.N. & Brinckmann-Voss A. (1980). Hydromedusae of British Columbia and Puget Sound. *Can. Bull. Fish. Aquat. Sci.* 204, 1-192.
- Ates, R.M.L. (1995). *Pagurus prideaux* and *Adamsia palliata* are not obligate commensals. *Crustaceana*, 68, 522–524.
- Ates, R.M.L. (2003). A preliminary review of zoanthid-hermit crab symbioses (Cnidaria; Zoantharia/Crustacea; Paguridea). *Zool. Verb. (Leiden)*, 345, 41–48.
- Ayres-Peres, L. & Mantelatto F.L. (2010). Epibiont occurrence on gastropod shells used by the hermit crab *Loxopagurus loxochelis* (Anomura: Diogenidae) on the northern coast of São Paulo, Brazil. *Zoologia*, 27, 222-227.
- Bach, C.E., Hazlett B.A. & Rittschof D. (2006). Sex-specific differences and the role of predation in the interaction between the hermit crab, *Pagurus longicarpus*, and its epibiont, *Hydractinia symbiolongicarpus*. *J. exp. mar. Biol. Ecol.* 333, 181– 189.
- Baeza, J.A. & Stotz W. (2003). Host-use and selection of differently colored sea anemones by the symbiotic crab *Allopetrolisthes spinifrons*. *J. Exp. Mar. Biol. Ecol.* 284, 25– 39.
- Balss, H. (1924). Über Anpassungen und Symbiose der Paguriden. Eine zusammenfassende übersicht. *Z. Morphol. Okol. Tiere*, 1, 752–792.
- Bavestrello, G. (1985). Idroidi simbiotici di paguri e gasteropodi nella Riviera Ligure di Levante. *Oebalia*, 11, 349–362.
- Bavestrello, G., Puce S., Cerrano C., Castellano L. & Arillo A. (2000). Water movement activating fragmentation: a new dispersal strategy for hydractiniid hydroids. *J.Mar. Biol. Assoc. UK.* 80, 361– 362.
- Bavestrello, G., Cerrano C., Camillo C., Puce S., Romagnoli T., Tazioli S. & Totti C. (2008). The ecology of protists epibiontic on marine hydroids. *J. Mar. Biol. Assoc. UK.* 88, 1611–1617.
- Bennet, I. (1962). Hydroids on Marine Snails. *Aus. Nat. Hist.* 14, 93.
- Blochmann, F. & Hilger C. (1888). Über *Gonactinia prolifera* Sars, eine durch Quertheilung sich vermehrende Actinie. *Morpholog. Jahrbuch.* 13, 384-400.
- Boas, J.E.V. (1926). Zur Kenntnis der Einsiedlerkrebse Paguropsis und seiner eigenartigen Behausung. *Selk. Biol. Medd.* 5, 1 – 23.
- Boero, F. & Fresi E. (1986). Zonation and Evolution of a Rocky Bottom Hydroid Community. *P.S.Z.N. I: Marine Ecology* 7, 123-150.
- Bouillon, J. (1994). Classe des Hydrozoaires. In: Grassé P.-P. (ed.), *Traité de Zoologie, Cnidaires* (pp. 29-416). Paris: Masson.
- Bouillon, J., Medel D. & Peña Cantero A.L. (1997). The taxonomic status of the genus *Stylactaria* Stechow, 1921 (Hydroidomedusae, Anthomedusae, Hydractiniidae), with the description of a new species. *Sci. Mar.* 61, 471– 486.
- Bouillon, J., Gravili C., Pagés F., Gili J.-M., Boero F., 2006. An introduction to Hydrozoa. *Mémoires du Muséum national d'Histoire naturelle* 194, 1-591.
- Bouvier, E.L. (1898). Sur la classification, les originès et la distribution des crabes de la famille des Dorrippidés. *Bull. Soc. Philomathique Paris*, 91, 54-70.
- Brooks, W.R. (1989). Hermit crabs alter sea anemone placement patterns for shell balance and reduced predation. *J. Exp. Mar. Biol. Ecol.* 132, 109– 121.
- Brooks, W.R. & Mariscal R.N. (1985). Protection of the hermit crab *Pagurus pollicaris* Say from predators by hydroid-colonized shells. *J. Exp. Mar. Biol. Ecol.* 87, 111 – 118.
- Brooks, W.R. & Mariscal R.N. (1986). Population variation and behavioral changes in two pagurids in association with the sea anemone *Calliactis tricolor* (Lesueur). *J. Exp. Mar. Biol. Ecol.* 103, 275–289.
- Brooks, W.R. & Rittschof D. (1995). Chemical detection and host selection by the symbiotic crab *Porcellana sayana*. *Invertebr. Biol.* 114, 180–185.
- Bruce, J.R., Colman J.S. & Jones N.S. (1963). *Marine Fauna of the Isle of Man and its Surrounding Seas*. Liverpool, 1-307pp.
- Brusca, R.C. & Brusca G.J. (2003). *Invertebrates*. Sunderland, Massachusetts: Sinauer Associates Inc.
- Bush, A.O., Fernández J.C., Esch G.W. & Seed J.R. (2001). *Parasitism: The Diversity and Ecology of Animal Parasites*. Cambridge: Cambridge Univ. Press.
- Buss, L.W. & Yund P.O. (1989). A sibling species group of *Hydractinia* in the north-eastern United States. *J. Mar. Biol. Assoc. U.K.* 69, 857–874.
- Cairns, S.D. & Barnard J.L. (1984). Redescription of *Janaria mirabilis*, a calcified hydroid from the Eastern Pacific. *Bull. South. Calif. Acad. Sci.* 83, 1 – 11.
- Cairns, Stephen D., Dale R. Calder, Anita Brinckmann-Voss, Clovis B. Castro, Daphne G. Fautin,... 2002. *Common and Scientific Names of Aquatic Invertebrates from the United States and Canada: Cnidaria and Ctenophora*, Second Edition. American Fisheries Society Special Publication 28. xi + 115.
- Calder, D.R. (1971). Hydroids and hydromedusae of southern Chesapeake Bay. Virginia Institute of Marine Science, *Special Paper in Marine Science*, 1, 1-125.
- Capart, A. (1948). Le *Lernaeocera branchialis*. *La cellule*, 52, 159-212.

- Carlgren, O. (1928). Zur Symbiose zwischen Aktinien und Paguriden. *Z. Morphol. Okol. Tiere*, 12, 165–173.
- Carlgren, O. (1938). South African Actinaria and Zoantharia. *K. Sven. Vetensk. Akad. Handl.* 3, 1–148.
- Carlgren, O. (1949). A survey of the Ptychodactaria, Corallimorpharia and Actiniaria. *Kungliga Svenska Vetenskaps Akad. Handlingar*, 4, 1–121.
- Carrington, J.T. & Lovett E. (1881). Notes and observations on British stalk-eyed Crustacea. *Zoologist*, 5, 97–101, 137–142, 198–205, 301–307, 358–364, 413–418, 455–461.
- Caruso, T., Falciai L. & Zupo V. (2003). Decapoda Anomura Paguridea: morpho-functional relationships and influence of epibiotic anemones on shell use along a bathymetric cline. *Crustaceana*, 76, 149–166.
- Cazaux, G. (1961). Signification et origine de l'association entre Hydractinie et Pagure: Rôle des tropismes larvaires dans le développement de l'Hydraire. *Bull. Stat. Biol. Arcachon N.S.* 13, 1–5.
- Cerrano, C., Bavestrello G., Puce S. & Sara M. (1998). Biological cycle of *Podocoryna exigua* (Cnidaria: Hydrozoa) from a sandy bottom of the Ligurian Sea. *J. Mar. Biol. Assoc. U.K.* 78, 1101–1111.
- Chintiroglou, Ch. & Doumenc D. (1998). *Isanthus Homolophilus* sp. nov. (Isanthidae: Actiniaria: Anthozoa) from French Polynesia. *J. Mar. Biol. Assoc. U.K.* 78, 829–834.
- Chintiroglou, Ch., Doumenc D. & Koutsoubas D. (1992). Allométrie d'une nouvelle association entre le Décapode Anomure *Pagurus excavatus* (Herbst, 1791) et l'Actinie Acontiaire *Sagartiogeton undatus* (Müller, 1788). *Crustaceana*, 62, 1–12.
- Christensen, H.E. (1967). Ecology of *Hydractinia echinata* (Fleming) (Hydroidea, Athecata): I. Feeding biology. *Ophelia*, 4, 245–275.
- Christidis, J., Chintiroglou Ch. & Culley M.B. (1997). A study of the populations of *Calliactis parasitica* (Couch, 1842) in symbiosis with anomuran decapods in Thermaikos Gulf (N. Aegean Sea). *Crustaceana*, 70, 227–238.
- Conover, M.R. (1976). The influence of some symbionts on the shell-selection behaviour of the hermit crabs, *Pagurus pollicaricus* [sic] and *Pagurus longicarpus*. *Anim. Behav.* 24, 191–194.
- Corliss, J. O. (1979). The ciliated Protozoa: characterization, classification, and guide to the literature, 2nd ed. 1–455. (Pergamon Press, New York, NY).
- Cowles, R.P. (1920). Habits of tropical crustacea: III. Habits and reactions of hermit crabs associated with sea anemones. *Philipp. J. Sci.* 15, 81–90.
- Cuadras, J. & Pereira F. (1977). Invertebrates associated with *Dardanus arrosor* (Anomura, Diogenidae). *Vie Milieu*, 27, 301–310.
- Cunningham, C.W., Buss L.W. & Anderson C. (1991). Molecular and geologic evidence of shared history between hermit crabs and the symbiotic genus *Hydractinia*. *Evolution*, 45, 1301–1316.
- Cutress, C.E. & Ross D.M. (1969). The sea anemone *Calliactis tricolor* and its association with the hermit crab *Dardanus venosus*. *J. Zool. (Lond.)*, 158, 225–241.
- Cutress, C., Ross D.M. & Sutton L. (1970). The association of *Calliactis tricolor* with its pagurid, calappid, and majid partners in the Caribbean. *Can. J. Zool.* 48, 371–376.
- Dales, R.P. (1957). Interrelations of organisms: A. Commensalism. *Mem. Geol. Soc. Amer.* 67, 391–412.
- Daly, M., Ardelean A., Cha H.-R., Campbell A.C. & Fautin D.G. (2004). A new species, *Adamsia obvolva* (Cnidaria: Anthozoa: Actiniaria), from the Gulf of Mexico, and a discussion of the taxonomy of carcinoecium-forming sea anemones. *Bull. Mar. Sci.* 74, 385–399.
- Damiani, C.C. (2003). Reproductive costs of the symbiotic hydroid *Hydractinia symbiolongicarpus* (Buss and Yund) to its hermit crab *Pagurus longicarpus* (Say). *J. Exp. Mar. Biol. Ecol.* 288, 1–20.
- Dechancé, M. & Dufaure J. (1959). Une nouvelle association entre une Actinie et un Pagure. *C. R. Hebd. Séances Acad. Sci.* 249, 1566–1568.
- Di Camillo, C.G., Bo M., Puce S., Tazioli S., Froglic C. & Bavestrello G. (2008). The epibiotic assemblage of *Geryon longipes* (Crustacea: Decapoda: Geryonidae) from the Southern Adriatic Sea. *Ital. J. Zool.* 75, 29–35.
- Doumenc, D. (1975). Actinies bathyales et abyssales de l'océan Atlantique nord. Famille des Hormathiidae (genres *Paracalliactis* et *Phelliactis*) et des Actinostolidae (genres *Actinoscyphia* et *Sicyonis*). *Bull. Mus. nat. Hist. nat. Paris*, 3 ser., 287: Zool. 197, 157–204.
- Doumenc, A.D. & Van Praët M. (1987). Ordre des Actiniaires. Ordre des Ptychodactiniaires. Ordre des Corallimorphaires. In: P-P- Grasse (ed.), *Traité de Zoologie* (pp. 257–401). Paris: Masson.
- Dunn, D.F. & Libermann M.H. (1983). Chitin in sea anemone shells. *Science*, 221, 137–139.
- Dunn, D.F., Devaney D.M. & Roth B. (1980). Stylobates: a shell-forming sea anemone (Coelenterata, Anthozoa, Actiniidae). *Pac. Sci.* 34, 379–388.
- Dvoretzky, A.G. & Dvoretzky V.G. (2008). Epifauna associated with the northern stone crab *Lithodes maia* in the Barents Sea. *Polar Biol.* 31, 1149–1152.
- Edwards, C. (1972). The hydroids and the medusae *Podocoryne areolata*, *P. borealis* and *P. carnea*. *J. Mar. Biol. Assoc. U.K.* 52, 97–144.
- England, K.W. (1971). Actinaria from Mururoa Atoll Tuamotu, Polynesia (Hormathiidae; *Calliactis polyopus* Sagartiidae: *Verrillactis* n. gen. *paguri*). *Cah. Pacifique*, 15, 23–40.
- Evans, F. (1978). The Marine Fauna of the Cullercoats District No. 6. Coelenterata and Ctenophora. *rep. Dove Mar. Lab.* 19, 1–165.
- Faurot, L. (1932). Actinies et pagures etude de psychologie animale. *Arch. Zool. Exp. Gen.* 74, 139–154.
- Fautin, D.G. (1987). *Stylobates loisetteae*, a new species of shell-forming sea anemone (Coelenterata: Actiniidae) from western Australia. *Proc. Calif. Acad. Sci.* 45, 1–7.
- Fernandez-Leborans, G. & Cárdenas C.A. (2009). Epibiotic communities on juvenile king crabs (*Lithodes santolla*) from subantarctic areas. *Polar Biology*, 32, 1693–1703.
- Fernandez-Leborans, G. & Gabilondo R. (2005). Hydrozoan and protozoan epibionts on two decapod species, *Liocarcinus depurator* (Linnaeus, 1758) and *Pilumnus hirtellus* (Linnaeus, 1761), from Scotland. *Zool. Anz.* 244, 59–72.
- Fernandez-Leborans, G. & Gabilondo R. (2006). Taxonomy and distribution of the hydrozoan and protozoan epibionts on *Pagurus bernhardus* (Linnaeus, 1758) (Crustacea, Decapoda) from Scotland. *Acta Zool.* 87, 33–48.
- Fernandez-Leborans, G. & Gabilondo R. (2008). Invertebrate and protozoan epibionts on the velvet swimming crab *Liocarcinus puber* (Linnaeus, 1767) from Scotland. *Acta Zool.* 89, 1–17.
- Fernandez-Leborans, G. & Herrero Córdoba M.J. (1997). Epibiosis of two sessile protozoan ciliate species on decapod crustaceans. *Cytobios*, 91, 143–154.

- Fernandez-Leborans, G. & Tato-Porto M.L. (2000a). A review of the species of protozoan epibionts on crustaceans. I. Peritrich ciliates. *Crustaceana*, 73, 643-684.
- Fernandez-Leborans, G. & Tato-Porto M.L. (2000b). A review of the species of protozoan epibionts on crustaceans. I. Suctorian ciliates. *Crustaceana*, 73, 1205-1237.
- Fernandez-Leborans, G., Zitzler K. & Gabilondo R. (2006). Protozoan ciliate epibionts on the freshwater shrimp *Caridina* (Crustacea, Decapoda, Atyidae) from the Malili lake system on Sulawesi (Indonesia). *J. Nat. Hist.* 40, 1983-2000.
- Folino, N.C. & Yund P.O. (1998). The distribution of hydroid sibling species on hermit crabs in estuaries in the Gulf of Maine. *Estuaries*, 21, 829– 836.
- Frey, R.W. (1987). Hermit crabs: Neglected Factors in Taphonomy and Paleoecology. *Palaios*, 2, 313– 322.
- Genzano, G.N. & Rodriguez, G.M. 1998. Association between hydroid species and their substrates from the intertidal zone of Mar del Plata (Argentina). *Miscellanea Zoológica*, 21, 21-29.
- Gili, J. M., Wervoort W. & Pages F. (1989). Hydroids from the West African coast: Guinea Bissau, Namibia and South Africa. *Scientia Mar.* 53, 67-112.
- Gili, J.M., Abelló P. & Villanueva R. (1993). Epibionts and intermolt duration in the crab *Bathynectes piperitus*. *Mar. Ecol. Progr. Ser.* 98, 107-113.
- Goto, S. (1910). On two species of *Hydractinia* living in symbiosis with a hermit crab. *J. Exp. Zool.* 9, 469– 496.
- Goto, R., Hamamura Y. & Kato M. (2007). Obligate commensalism of *Curvemysella paula* (Bivalvia: Galeommatidae) with hermit crabs. *Mar. Biol.* 151, 1615-1622.
- Grant, W.C. & Pontier P.J. (1973). Fitness in the hermit crab *Pagurus acadianus* with reference to *Hydractinia echinata*. *Bull.-Mt. Desert Isl. Biol. Lab.* 13, 50– 53.
- Haig, J. & Ball E.E. (1988). Hermit crabs from north Australia and eastern Indonesian waters (Crustacea Decapoda: Anomura: Paguroidea) collected during the 1975 *Alpha Helix* Expedition. *Rec. Austral. Mus.* 40, 151-196.
- Hamond, R. (1957). Notes on Hydrozoa of the Norfolk Coast. *J. Linnean Soc. (Zoology)*, 43, 294-324.
- Hand, C. (1975). Behaviour of some New Zealand sea anemones and their molluscan and crustacean hosts. *N.Z. J. Mar. Freshw. Res.* 9, 529– 538.
- Hartnoll, R.G. (1971). The relationship of an amphipod and a spider crab with the snakelocks anemone. *Reports of the Marine Biological Station, Port Erin*, 83, 37-42.
- Hartog, J.C. den (1990). Associated occurrence of *Cyclocoeloma tuberculata* Miers, 1880 (Decapoda: Majidae) and species of Discosomatidae (Anthozoa: Corallimorpharia). *Zool. Med. Leiden*, 64,161-168
- Hayashi, R. & Tsuji K. (2007). Spatial distribution of turtle barnacles on the green sea turtle, *Chelonia mydas*. *Ecol. Res.* 23,121-125.
- Hiro, J. (1939). Notes on the animals found on *Macrocheira keimpeferi* De Haan. III. Hydroids. *Annot. Zool. Jap.* 18, 167-176.
- Holt, E.W. (1890). Foreign substances attached to crabs. *Nature, London*, 41, 463-464.
- Imafuku, M., Yamamoto T. & Ohta M. (2000). Predation on symbiont sea anemones by their host hermit crab *Dardanus pedunculatus*. *Mar. Fresh. Behaviour Physiol.* 33, 221-232.
- Ingle, R.W. (1983). Shallow-water crabs. *Synop. British Fauna. New Series* 25, 1–206.
- Ingle, R.W. (1996). Shallow-water crabs. *Synop. British Fauna. New Series* 25, 2nd ed., 1–243.
- Ingle, R.W. & Christiansen M.E. (2004). *Lobsters, mud shrimps and anomuran crabs. Keys and notes for the identification of species.* Field Studies Council for Linnaean Soc. London. Shrewbury: Estuarine and Coastal Sciences Assoc.
- Jensen, K. & Bender K. (1973). Invertebrates associated with snail shells inhabited by *Pagurus bernhardus* (L.) (Decapoda). *Ophelia*, 10, 185– 192.
- Jones, D.H. (1966). A gymnoblastic hydroid occurring on *Sphyrion lumpi* (Kröyer). *Ann. Magaz. nat. Hist.* 9, 173-181.
- Jungersen, H.F.E. (1911). On a new gymnoblastic Hydroid (*Lebthyocodium sarvotretis* epizoico on a new parasitic Copepod (*Sarvotretes scopeli*) infesting *Scopelus glacialis* Rhdt. *Vidensk. Meddel. fra den naturh. Foren.* 64, 9-33.
- Karlson, R.H. & Cariolou M.A. (1982). Hermit crab shell colonization by *Crepidula convexa* Say. *J. Exp. Mar. Biol. Ecol.* 65, 1–10.
- Karlson, R.H. & Shenk M.A. (1983). Epifaunal abundance, association, and overgrowth patterns on large hermit crab shells. *J. Exp. Mar. Biol. Ecol.* 70. 55–64.
- Karplus, G.C., Fiedler G.C. & Ramcharan P. (1998). The Intraspecific Fighting Behavior of the Hawaiian Boxer Crab, *Lybia edmondsoni* -Fighting with Dangerous Weapons?. *Symbiosis*, 24, 287-302.
- Key, M.M., Winston J.E., Volpe J.W., Jeffries W.B. & Voris H.K. (1999). Bryozoan fouling of the blue crab *Callinectes sapidus* at Beaufort, North Carolina. *Bull. mar. Sci.* 64, 513-533.
- Kramp, P.L. (1935). Polydyr (Coelenterata): I. Ferskvandspolypper og Goplepolypper. *Dan. Fauna*, 41, 1–208.
- Lavaley, M.S.S. & Hartog J.C. den (1995). A case of associated occurrence of the crab *Lauridromia intermedia* (Laurie, 1906) (Crustacea: Decapoda: Dromiidae) and the actinian *Nemanthus annamensis* Carlgren, 1943 (Anthozoa: Actiniaria: Nemanthidae). *Zool. Med. Leiden*, 69, 121-130,
- Leloup, E. (1931). Un cas d'épibiose de l'Hydropolype, *Laomedea geniculata* (Linné). *Bull. Mus. Roy. Hist. Nat. Belg.* 7, 1-3.
- Lemaitre, R. (1996). Hermit crabs of the family Parapaguridae (Crustacea: Decapoda: Anomura) from Australia: species of *Strobopagurus* Lemaitre, 1989, *Sympagurus* Smith, 1883 and two new genera. *Rec. Aust. Mus.* 48, 163–221.
- Little-Ostarello, G. (1973). Natural history of hydrocoral *Allopora californica* Verrill (1866). *Biol. Bull.* 145, 548-564.
- Lopez-Victoria, M., Barrios L.M., Kraus H. & Osorio L.A. (2004). Nuevos aspectos sobre las relaciones simbióticas entre *Dardanus fucosus* (Crustacea: Paguridae), *Calliactis tricolor* (Cnidaria: Hormathiidae) y *Porcellana sayana* (Crustacea: Porcellanidae). *bol. invernmar*, 33, 1-4.
- Maldonado, M. & Uriz M.J. (1992). Relationship between macroalgal functional form groups and substrata stability in a subtropical rocky intertidal system. *J. Exp. Mar. Biol. Ecol.* 74,13–34.
- Marine Biological Association, (1957). *Plymouth Marine Fauna.* Plymouth I-XLIII, 457 pp.
- Matthews, D.C. (1959). Observations on ova fixation in the hermit crab *Eupagurus prideauxii*. *Pubbl. Stn. Zool. Napoli*, 31, 248– 263.
- McGaw, I.J. (2006). Epibionts of sympatric species of *Cancer* crabs in Barkley sound, British Columbia. *J. Crust. Biol.* 26, 85-93.
- McLean, R. (1983). Gastropod shells: a dynamic resource that helps shape benthic community structure. *J. Exp. Mar. Biol. Ecol.* 69, 151– 174.
- Mercado, N.A. & Lytle C.F. (1980). Specificity in the association between *Hydractinia echinata* and sympatric species of hermit crabs. *Biol. Bull. (Woods Hole)*, 159, 337– 348.

- Millard, N.A.H. (1975). Monograph on the Hydroida of southern Africa. *Ann. S. Afr. Mus* 68, 1–513.
- Mills, C.E. (1976). *Podocoryne selena*, a new species of hydroid from the Gulf of Mexico, and a comparison with *Hydractinia echinata*. *Biol. Bull. (Woods Hole)*, 151, 214–224.
- Moser, M. & Taylor S. (1978). Effects of the copepod *Cardiodectes medusaeus* on the lanternfish *Stenobrachius leucopsarus* with notes on hypercastration by the hydroid *Hydrichthys* sp. *Can. J. Zool.* 56, 2372–2376.
- Muirhead, A., Tyler P.A. & Thurston M.H. (1986). Reproductive biology and growth of the genus *Epizoanthus* (Zoantheida) from the north–east Atlantic. *J. Mar. Biol. Assoc. U.K.* 66, 131–143.
- Mulisch, M., Harry O.G., Patterson D.J. & Wyatt C. (1986). Folliculinids (Ciliata: Heterotrichida) from Portaferry, County Down, including a new species of *Metafolliculina* Dons, 1924. *Irish nat. J.* 22, 1–7.
- Nogueira, M.J., De Castro M.R. & Haddad M.A. (2006). *Calliactis tricolor* (Anthozoa, Acontiarina) epibiosis on Brachyura (Crustacea, Decapoda) at the South of Paraná and North of Santa Catarina coast. *Acta Biol. Par., Curitiba*, 35, 233–248.
- Overstreet, R.M. (1983). Metazoan symbionts of crustaceans. In: Bliss DE, editor, *The biology of Crustacea* (pp. 155–250). London: Academic Press.
- Overstreet, R.M. (1987). Solving parasite-related problems in cultured crustacea. *International J. Parasitol.* 17, 309–318.
- Parulekar, A. (1969). *Neoaiphtasia commensali*, gen. et. sp. nov.: an actiniarian commensal of hermit crabs. *J. Bombay Nat. Hist. Soc.* 66, 57–62.
- Paulay, G., Puglisi M.P. & Starmer J.A. (2003). The non-scleractinian Anthozoa (Cnidaria) of the Mariana Islands. *Micronesica*, 35–36, 138–155.
- Puce, S., Cerrano C., Di Camillo C.G. & Bavestrello G. (2008). Hydroidomedusae (Cnidaria: Hydrozoa) symbiotic radiation. *J. Mar. Biol. Assoc. U. K.* 88, 1715–1721.
- Reaka, M.L. (1978). The effects of an ectoparasitic gastropod, *Caledoniella montrouzieri*, upon moulting and reproduction of a stomatopod crustacean, *Gonodactylus viridis*. *Veliger*, 21, 251–254.
- Rees, W.J. (1967). A brief survey of the symbiotic associations of Cnidaria and Mollusca. *Proc. Malacol. Soc. Lond.* 37, 213–231.
- Reiss, H., Knäuper S. & Kröncke I. (2003). Invertebrate associations with gastropod shells inhabited by *Pagurus bernhardus* (Paguridae)—secondary hard substrate increasing biodiversity in North Sea soft-bottom communities. *Sarsia*, 88, 404–415.
- Remane, A. (1980). *Systematische Zoologie*. Stuttgart: Gustav Fischer Verlag.
- Roper, R.E. (1913). The marine Polyzoa of Northumberland. *Reports of the Dove Marine Laboratory*, 2, 36–57.
- Ross, D.M. & Sutton L. (1961). The response of the sea anemone *Calliactis parasitica* to shells of the hermit crab *Pagurus bernhardus*. *Proc. R. Soc. Lond., B Biol. Sci.* 155, 266–281.
- Ross, D.M. & Sutton L. (1961). Detachment of sea anemones by commensal hermit crabs and by mechanical and electrical stimuli. *Nature*, 217, 380–381.
- Ross, D.M. (1979). A behaviour pattern in *Pagurus bernhardus* L. towards its symbiotic actinian *Calliactis parasitica* (Couch). *J. Mar. Biol. Assoc. UK.* 59, 623–630.
- Ross, D.M. & Zamponi M.O. (1982). A symbiosis between *Paracalliactis mediterranea* n. sp. (Anthozoa-Actiniaria) and *Pagurus variabilis* A. Milne-Edwards and Bouvier. *Vie Milieu*, 32, 175–181.
- Ross, D.M. (1967). Behavioural and ecological relationships between sea anemones and other invertebrates. *Oceanogr. Mar. Biol. Annu. Rev.* 5, 291–316.
- Ross, D.M. (1970). The commensal association of *Calliactis polyopus* and the hermit crab *Dardanus gemmatus* in Hawaii. *Can. J. Zool.* 48, 351–357.
- Ross, D.M. (1971). Protection of hermit crabs (*Dardanus* spp.) from octopus by commensal sea anemones (*Calliactis* spp.). *Nature*, 230, 401–402.
- Ross, D.M. (1983). Symbiotic relations. In: Bliss, D. (Ed.), *The Biology of the Crustacea* (pp. 163–212). New York: Academic Press.
- Roule, L. (1900). Description d'une nouvelle espèce Méditerranéenne de zoanthidé, commensale des pagures (Palythoa–Gemmaria–Paguricola). *Bull. Soc. Zool. Fr.* 25, 120–125.
- Ruthensteiner, B., Reinicke G.B. & Straube N. (2008). The Type Material of Hydrozoa described by Eberhard Stechow in the Zoologische Staatssammlung München. *Spixiana*, 31(1), 3–27.
- Samuelsen, T.J. (1970). The biology of six species of Anomura (Crustacea, Decapoda) from Raunefjorden, Western Norway. *Sarsia*, 45, 23–52.
- Sandford, F. (2003). Population dynamics and epibiont associations of hermit crabs (Crustacea: Decapoda: Paguroidea) on Dog Island. *Mem. Mus. Vic.* 60, 45–52.
- Schijfsma, K. (1935). Observations on *Hydractinia echinata* (Flem.) and *Eupagurus bernhardus* (L.). *Arch. Neerl. Zool.* 1, 261–314.
- Schembri, P.J. & McLay C.L. (1983). An annotated key to the hermit crabs (Crustacea : Decapoda : Anomura) of the Otago region (southeastern New Zealand). *New Zealand J. Mar. Fresh. Res.* 17, 27–35.
- Schejter, L. & Mantelatto F.L. (2009). Shelter association between the hermit crab *Sympagurus dimorphus* and the zoanthid *Epizoanthus paguricola* in the southwestern Atlantic Ocean. *Acta Zool.*, DOI: 10.1111/j.1463-6395.2009.00440.x
- Schuchert, P. (1996). The marine fauna of New Zealand: Athecate hydroids and their medusae (Cnidaria, Hydrozoa). *New Zealand Oceanogr. Inst. Mem.* 119, 5–538.
- Schuchert, P. (2007). The European athecate hydroids and their medusae (Hydrozoa, Cnidaria): Filifera part 2. *Rev. suisse Zool.* 114, 195–396.
- Schuchert, P. (2008). The European athecate hydroids and their medusae (Hydrozoa, Cnidaria): Filifera part 3. *Rev. suisse Zool.* 115, 221–302.
- Schuchert, P. (2010). The European athecate hydroids and their medusae (Hydrozoa, Cnidaria): Capitata part 2. *Rev. suisse Zool.* 117, 337–555.
- Silberman, J.D., Collins A.G., Gershwin L.-A., Johnson P.J. & Roger A.J. (2004). Ellobiopsids of the Genus *Thalassomyces* are Alveolates. *J. Eukaryot. Microbiol.* 51, 246–252.
- Silva-Neto, I.D. da, Silva Paiva T. da Pedroso R.J., Alexandre C.J. & Migotto A.E. (2012). Redescription of *Licinophora chattoni* Villeneuve-Brachon, 1939 (Ciliophora, Spirotrichea), associated with *Zyzygus warreni* Calder, 1988 (Cnidaria, Hydrozoa). *Eur. J. Protistol.* 48, 48–62.
- Stachowitsch, M. (1980). The epibiotic and endolithic species associated with the gastropod shells inhabited by the hermit crabs *Paguristes oculatus* and *Pagurus cuanensis*. *P. S. Z. N. Mar. Ecol.* 1, 73–101.
- Števcič, Z. (1966). Z-ivotni kompleks Rakovice *Maja squinado* Herbst (life complex of spinous spider crab *Maja squinado* Herbst). *Ekologija*, 1, 109–119.

- Strenth, N.E. & Chace F.A. Jr (1995). New records of natant decapods (Crustacea, Palaemonidae) from the south Texas Coast. *Texas J. Sci.*, a 0128663722.
- Svoboda, A., Stepanjants S.D. & Ljubenkov J. (2006) The genus *Bouillonia* (Cnidaria: Hydrozoa: Anthoathecata). Three species from the northern and southern hemispheres, with a discussion of bipolar distribution of this genus. *Zool. Med. Leiden*, 80-4, 185-206.
- Teissier, G. (1965). Cnidaire-Cténaire. Inventaire de la Faune Marine de Roscoff. *Trav. Stat. Biolog. Roscoff*, 16, 1-64.
- Templeman, W. (1973). First record of the gymnoblastic hydroid, *Ichthyocodium sarvotretis*, on the copepod *Sphyrion lumpi*, from redfish of the northwest Atlantic. *J. Fish. Res. Board Canada*, 30, 1655-1660.
- Threlkeld, S.T., Chiavelli D.A. & Willey R.L. (1993). The organization of zooplankton epibiont communities. *Trends Ecol. Evol.* 8, 317–321.
- Urzelai, A., Elizalde M., Capellan T., Esteban I., Quiroga A., Zabala Y. & Ibañez M. (1990). Estudio preliminar de las comunidades de *Pagurus alatus* Fabricius, 1775 y *Parapagurus pilosimanus* S. I. Smith, 1879 (Crustacea, Decapoda) y *Epizoanthus paguriphilus* Verrill, 1883 (Anthozoa, Zoantaria) de la fosa de Cap Breton (Golfo de Vizcaya). *Lurralde*, 13, 193– 206.
- Vader, W., Johannessen O.H. & Christiansen B.O. (1981). A pelagic isopod, *Systemus infelix* overgrown with hydroids. *Fauna norv. Ser. A2*, 47-48.
- Van Winkle, D.H., Longnecker K. & Blackstone N.W. (2000). The effects of hermit crabs on hydractiniid hydroids. *Mar. Ecol. Publ. Sta. Zool. Napoli*, 21, 55-67.
- Viljoen, S. & Van AS, J.G. (1983). A taxonomic study of sessile peritrichians of a small impoundment with notes on their substrate preferences. *J. limnol. Soc. Southern Africa*, 9, 33-42.
- Walker, S.E. (1988). Taphonomic significance of hermit crabs (Anomura: Paguridea): epifaunal hermit crab - infaunal gastropod example. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 63, 45– 71.
- Wahl, M. (1989). Marine epibiosis. I. Fouling and antifouling: some basic aspects. *Mar. Ecol. Prog. Ser.* 58, 175-189.
- Warren, A. & Robson E.A. (1998). The identity and occurrence of *Kerona pediculus* (Ciliophora, Hypotrichida), a well-known epi-zoite of *Hydra vulgaris* (Cnidaria, Hydrozoa). *Zool. Verb.* 323, 235–245.
- Widmer, C.L., Cailliet G. & Geller J. (2010). The life cycle of *Earleria corachloae* n. sp. (Cnidaria: Hydrozoa) with epibiotic hydroids on mid-water shrimp. *Mar. Biol.* 157, 49–58.
- Willey, R.L., Cantrell P.A. & Threlkeld S.T. (1990). Epibiotic flagellates increase the susceptibility of some zooplankton to fish predation. *Limnol. Oceanogr.* 35, 952–959.
- Yusa, Y., Yamato S., Marumura M. (2001). Ecology of a parasitic barnacle, *Koleolepas avis*: relationship to the hosts, distribution, left– right asymmetry and reproduction. *J. Mar. Biol. Assoc. U.K.* 81: 781–788.

List of cnidarian species found as epibiont on crustacea. (Classification according to Bouillon et al. 2006, Cairns et al. 2002)

Cnidaria	Crustacea	References	Geographical distribution
	class Hydrozoa; subclass Hydroidina; order Leptothecatae		
<i>Eumudicola validivivae</i> Stechow, 1921	<i>Parapylocheles scorpio</i> (Alcock, 1894)	Ruthensteiner et al. 2008	Western Sumatra; south-west of Pulo Nias
<i>Aglaophenia kirchenpaueri</i> (Heller, 1868)			
<i>Aglaophenia octodonta</i> (Heller, 1868)			
<i>Aglaophenia picardi</i>	Barnacles (<i>Balanus perforatus</i>) (1)	Boero&Fresi 1986	Portofino, Italy
<i>Aglaophenia tubiformis</i> (Heller, 1868)			
<i>Cyrtia hemisphaerica</i> (Linnaeus, 1767)	<i>Inachus dorsetensis</i> (Pennant, 1777); <i>Maja squinado</i> (Herbst, 1788); <i>Hymenosoma orbiculare</i> Desmarest, 1823; <i>Lemaecocera branchialis</i> (Linnaeus, 1767)	Marine Biological Association 1957; Ingle 1983; Millard 1975; Leloup 1931; Capart 1948	(Plymouth, UK); (Northern South West Africa to Mozambique)
<i>Cyrtia hemisphaerica</i> (as <i>Companularia johnstoni</i>)	<i>Pagurus bernhardus</i>	Jensen&Bender 1973	Marine waters of Sweden
<i>Companularia hincsi</i> Alder, 1856	Barnacles (<i>Balanus perforatus</i>) (1)	Boer&Fresi 1986	Portofino, Italy
<i>Companularia integra</i> MacGillivray, 1842			
<i>Obelia geniculata</i> (Linnaeus, 1758) (as <i>Laomedea geniculata</i>)	<i>Lemaecocera branchialis</i> (Linnaeus, 1767)	Leloup 1931, Capart 1948	
<i>Obelia geniculata</i>	<i>Macrocheira kaempferi</i> (Temminck, 1836); <i>Lithodes maja</i> (Linnaeus, 1758) (as <i>L. maja</i>); <i>Paralithodes camtschaticus</i> (Tilesius, 1815); <i>Anilocra physodes</i> (Linnaeus, 1758); <i>Pagurus bernhardus</i> ; <i>Dardanus arrosor</i> ; <i>Jasus lalandi</i> (H. Milne-Edwards, 1837); <i>Lemaecocera branchialis</i>	Dvoretzky&Dvoretzky 2008, Leloup 1931, Reiss et al. 2003, Ross&Zamponi 1982	Barents Sea, North Sea, Mediterranean Sea
<i>Laomedea calceifera</i> (Hincis, 1871) (<i>Laomedea conferta</i>)	<i>Hyas araneus</i> (Linnaeus, 1758)	Hamond 1957; Ingle, 1983	(Monterey Bay, California, USA)
<i>Ealeria corachloae</i> Widmer, Cailliet & Geller, 2010	<i>Paspipha pacifica</i> Rathbun, 1902	Widmer et al. 2010; Hiro 1939	Tonda, Sakai, Wakayama-Ken, Japan
<i>Obelia undotheca</i> Stechow 1923	<i>Oregania gracilis</i> Dana, 1851	Ruthensteiner et al. 2008	Bering Sea, Kamtschatka Peninsula, Awatsch
<i>Obelia longissima</i> (Pallas, 1766)	<i>Lithodes maja</i> ; <i>Paralithodes camtschaticus</i> (Tilesius, 1815); <i>Syscenus infelix</i> Harger, 1880	Dvoretzky&Dvoretzky 2008, Vader et al. 1981	(Barents Sea)
<i>Obelia commissurata</i> McCrady, 1859	<i>Balanus eburneus</i> Gould, 1841	Calder 1971	(Chesapeake Bay, USA)
<i>Obelia dichotoma</i> (Linnaeus, 1758)	<i>Dardanus arrosor</i> , <i>Lepas</i> , <i>Balanus eburneus</i> , <i>Mithraculus forceps</i> A. Milne-Edwards, 1875	Cuadras&Pereira 1977, Calder 1971	(Mediterranean Sea); (Chesapeake Bay, USA)
<i>Obelia</i> sp.	<i>Cancer productus</i> Randall, 1840	Mc Gaw 2006	(Barkley sound, British Columbia)
<i>Orthopsis integra</i> (MacGillivray, 1842)	<i>Dehaanuis dentatus</i> (H. Milne-Edwards, 1834)	Millard 1975	(Southern Africa)
<i>Plumularia obliqua</i>	<i>Balanus perforatus</i>	Boero&Fresi 1986	Portofino, Italy
<i>Plumularia setacea</i> (Linnaeus, 1758)			
<i>Halecium crinis</i> Stechow, 1913	<i>Macrocheira kaempferi</i>	Hiro 1939	(Tonda, Sakai, Wakayama-Ken, Japan)
<i>Halecium beani</i> (Johnston, 1838)	Barnacles (<i>Balanus perforatus</i>) (1)	Boero&Fresi 1986	Portofino, Italy
<i>Halecium mediterraneum</i>			
<i>Halecium labrosum</i> Alder, 1859	<i>Hyas araneus</i> (Linnaeus, 1758); <i>Balanus perforatus</i> (1)	Hamond 1957; Boero&Fresi 1986	(Norfolk Coast, USA); Portofino, Italy
<i>Sertularia gaudichaudi</i>	<i>Balanus perforatus</i>	Boero&Fresi 1986	Portofino, Italy
<i>Sertularia sinensis</i> Jäderholm, 1896	<i>Macrocheira kaempferi</i>	Hiro 1939	(Tonda, Sakai, Wakayama-Ken, Japan)
<i>Sertularia ellisi</i> (Deshayes & Milne Edwards, 1836) (as <i>Sertularia polyzonias</i>)	<i>Maja squinado</i> (Herbst, 1788)	Števíč 1966	
<i>Sertularia turgida</i> (Trask, 1857)	<i>Metacarcinus gracilis</i> (Dana, 1852) (as <i>Cancer gracilis</i>); <i>Metacarcinus magister</i> (Dana, 1852) (as <i>Cancer magister</i>); <i>Cancer productus</i> Randall, 1840	Mc Gaw 2006	(Barkley sound, British Columbia)
<i>Sertularia</i>	<i>Libinia spinosa</i> H. Milne-Edwards, 1834		
<i>Sertularia argentea</i> (Linnaeus, 1758) (as <i>Sertularia cupressina</i> var. <i>argentea</i>)	<i>Pisa armata</i> (Latreille, 1803)	Carrington&Lovett 1881	(Chesapeake Bay, USA)
<i>Sertularia argentea</i> (Linnaeus, 1758)	<i>Libinia</i> , <i>Amphibalanus improvisus</i> (Darwin, 1851) (as <i>Balanus improvisus</i>)	Calder 1971	
<i>Hydrilmania falcata</i> (Linnaeus, 1758)	<i>Pisa armata</i> , <i>Pisa tetraodon</i> (Pennant, 1777)	Carrington&Lovett 1881	
<i>Abietinaria exosporoides</i> Stechow, 1923	<i>Oregania gracilis</i> Dana, 1851	Ruthensteiner et al. 2008	(Pacific Grove, Monterey Bay, California, USA)
<i>Abietinaria pacifica</i> Stechow, 1923	<i>Oregania gracilis</i>	Ruthensteiner et al. 2008	(Pacific Grove, Monterey Bay, California, USA)
<i>Pilaolella</i> sp.	<i>Pagurus bernhardus</i>	Fernandez-Leborans & Gabilondo 2006	Clyde and Argyll, Scotland)
<i>Stegopoma plicatella</i> (Sars, 1863)	<i>Bathynectes piperitus</i> Manning & Holthuis, 1981	Gili et al. 1993	(off southern Namibia, Africa)
	<i>Neolithodes asperimus</i> Barnard, 1947	Gili et al. 1989	(Namibian coast)
<i>Kirchenpaueria pinnata</i> (Linnaeus, 1758)	<i>Hyas araneus</i>	Hamond 1957	(Norfolk coast, USA)
<i>Ventromma halecioides</i> (Alder, 1859)	Barnacles (<i>Balanus perforatus</i>) (1)	Boero&Fresi 1986	Portofino, Italy
<i>Dynamena disticha</i> (Bosc, 1802)			
<i>Nemertesia antennina</i> (Linnaeus, 1758)	<i>Hyas araneus</i> ; <i>Hyas coarctatus</i> Leach, 1816	Hamond 1957	(Norfolk coast, USA)
<i>Holoptesia tenella</i> (Verrill, 1874) (as <i>Schizotracha tenella</i>)	<i>Balanus eburneus</i> Gould, 1841; <i>Libinia</i>	Calder 1971	(Chesapeake Bay, USA)
	order Anthothecatae		
<i>Stylaster californica</i> (Verrill, 1866) (as <i>Allopora californica</i>)	<i>Armato-balanus nefrens</i> (Zullo, 1963) (as <i>Balanus nefrens</i>)	Little-Ostarell 1973	(Carmel river Beach, Bluefish Cove, Carmel, California, USA)
<i>Leucartaria octona</i> (Fleming, 1823)	<i>Macrocheira kaempferi</i> , <i>Geryon longipes</i> (H. Milne-Edwards, 1882)	Hiro 1939, Di Camillo et al. 2008	(Tonda, Sakai, Wakayama-Ken, Japan); (Southern Adriatic Sea)
	<i>Carcinus maenas</i> (Linnaeus, 1758); <i>Hyas</i> , <i>Halicarcinus</i> , <i>Macrocheira kaempferi</i>	Marine Biological Association 1957; Ingle 1983	(Plymouth, UK)
<i>Leucartaria</i> sp.	<i>Liocarcinus depurator</i> , <i>Pagurus bernhardus</i> , <i>Liocarcinus puber</i>	Fernandez-Leborans&Gabilondo 2005, 2006, 2008	(Isle of Cumbrae, Scotland)
<i>Hydrichthys</i> sp.	<i>Cardiodectes medusae</i> (Wilson C.B., 1908) (on the fish <i>Stenobrachius leucosparus</i>)	Moser&Taylor 1978	(Santa Barbara, San Diego, California, USA)
<i>Hydrichthys</i> (lchthyocodium) <i>sarcotretis</i> (Jungersen, 1913)	Eigenmann & Eigenmann, 1890		
	<i>Sarcotretes scopell</i> Jungersen, 1911 (on the fish <i>Benthosema glaciale</i> (Reinhardt, 1837) (as <i>Scopellus glacialis</i>))	Jungersen 1911, Templeman 1973	(SE coast Ireland); (NW Atlantic)
	<i>Sphyrna lumpi</i> (Krøyer, 1845) (on the fish <i>Sebastes mentella</i> Travin, 1951)	Jones 1966 Templeman 1973	(European Atlantic Ocean); (NW Atlantic)
<i>Coryne muscolides</i> (Linnaeus)			
<i>Coryne pusilla</i> Gaertner, 1774	Barnacles (<i>Balanus perforatus</i>) (1); <i>Hyas coarctatus</i> Leach, 1815; <i>Anapagurus hendersoni</i> Barnard, 1947; <i>Anapagurus cf. hendersoni</i> ; <i>Pagurus bernhardus</i>	Boero&Fresi 1986, Roper 1913, Millard 1975, Ingle 1983, Jensen&Bender 1973, Kramp 1935	(Portofino, Italy); (Northumberland, UK); (Plymouth, UK); (False Bay to Mossel Bay, Southern Africa); (Roscoff, France)
<i>Cladocoryne floccosa</i>			
<i>Dicorone conferta</i> (Alder, 1856)			
<i>Dicorone validivivae</i> Stechow, 1923	<i>Parapylocheles scorpio</i> (Alcock, 1894)	Ruthensteiner et al. 2008	Western Sumatra
<i>Eudendrium capillare</i> Alder, 1856	<i>Hyas coarctatus</i> ; <i>Balanus perforatus</i> (1)	Bruce et al. 1963; Ingle, 1983; Boero&Fresi 1986	(Isle of Man, UK); (Portofino, Italy)
<i>Eudendrium racemosum</i>	Barnacles (<i>Balanus perforatus</i>) (1)	Boero&Fresi 1986	Portofino, Italy
<i>Eudendrium ramosum</i> (Linnaeus, 1758)	Barnacles (<i>Balanus perforatus</i>) (1)	Boero&Fresi 1986	Portofino, Italy
<i>Eudendrium album</i> Nutting, 1898	<i>Balanus improvisus</i> Darwin, 1854	Calder 1971	(Chesapeake Bay, USA)
<i>Peratella propagulata</i> Bavestrello, 1987	Hermit crabs	Schuchert, 2007	
<i>Hydractinia</i> sp.	<i>Geryon longipes</i> A. Milne-Edwards, 1882	Di Camillo et al. 2008	(Southern Adriatic Sea)
	<i>Anapagurus chirocanthus</i> (Uljeborg, 1856); <i>Cibbanarius vittatus</i> (Bosc, 1802); <i>Dardanus arrosor</i> ; <i>Paguristes humi</i> Wass, 1955; <i>Pagurus acadianus</i> J. E. Benedict, 1901; <i>P. annulipes</i> (Stimpson, 1860); <i>P. bernhardus</i> ; <i>P. carolinensis</i> McLaughlin, 1975; <i>P. cuanensis</i> Bell, 1845; <i>P. excavatus</i> (Herbst, 1791); <i>P. longicarpus</i> Say, 1817; <i>P. pollicaris</i> Say, 1817; <i>Petrochirus diogenes</i> (Linnaeus, 1758); <i>Balanus eburneus</i> Gould, 1841	Karlson&Caroliou 1982, Ingle&Christiansen 2004; Sandford 2003, Samuelsen, 1970; Mercado&Lytle 1980 (North American Coast Atlantic Ocean); Cuadras&Pereira 1977; Cunningham et al. 1991; Karlson&Shenck 1983 Puce et al. 2008; Reiss et al. 2003 (North Sea); Calder 1971	(Dog Island, St George Sound, Florida, USA); (Raunefjorden, Western Norway); (Mediterranean Sea); (North Atlantic area and Gulf of Mexico); (Chesapeake Bay, USA)
	<i>Pagurus</i> sp.	Schiffma 1935; Cazaux 1961; Bennet 1962; Rees 1967; Christensen 1967; Mills 1976; Mercado&Lytle 1980; Brooks&Mariscal 1986	(Knåhaken Reef in the Øresund, Denmark); (Gulf of Mexico); (North American Coast Atlantic Ocean); (northeastern Gulf of Mexico)
<i>Hydractinia epiconcha</i> Stechow, 1908	<i>Pagurus gracilipes</i> (Stimpson, 1858); <i>Dardanus arrosor</i> (Herbst, 1796) (as <i>Pagurus arrosor</i>)	Balls 1924; Mills 1976	Gulf of Mexico
<i>Hydractinia sodalis</i> Stimpson, 1859	<i>Pagurus constans</i> (Stimpson, 1858)	Goto 1910; Rees 1967	
<i>Hydractinia symbiolongicarpus</i> Buss & Yund, 1989	<i>Pagurus longicarpus</i> Say, 1817; <i>P. pollicaris</i>	Damiani 2003; Bach et al. 2006; Cunningham et al. 1991; Buss&Yund 1989	(Beaufort, NC, USA); (North Atlantic area and Gulf of Mexico); (Long Island Sound, USA)
<i>Hydractinia diogenes</i> Millard, 1959	<i>Diogenes costatus</i> Henderson, 1893	Millard 1975	(Inhaca to Morumbene, South Africa)
<i>Hydractinia epiconcha</i> Stechow, 1908	<i>Dardanus arrosor</i> , <i>Pagurus gracilipes</i>	Dales 1957	
<i>Hydractinia polyclima</i> L. Agassiz, 1862	<i>Pagurus acadianus</i> J. E. Benedict, 1901; <i>P. longicarpus</i> ,	Cunningham et al. 1991; Folino&Yund 1998	(North Atlantic area and Gulf of Mexico)
<i>Hydractinia pugilator</i> (Hesseckel, 1880)	<i>Diogenes pugilator</i> (Roux, 1829)	Puce et al. 2008	
<i>Hydractinia pruvoti</i> (Motz-Kossowska, 1905)	<i>Cibbanarius erythropus</i> (Latreille, 1818)	Puce et al. 2008	
<i>Hydractinia rubricata</i> Schuchert, 1996 (Edwards, 1972) (as <i>Hydractinia selena</i>)	<i>Diacanthurus rubricatus</i> (Henderson, 1888) (as <i>Pagurus rubricatus</i>)	Schuchert, 1996	(Kaikoura Peninsula, New Zealand)
<i>Hydractinia aculeata</i> (Wagner, 1833)	<i>Cestopagurus timidus</i> (Roux, 1830) (as <i>Catopagurides timidus</i>)	Schuchert, 2008	Mediterranean. Type locality: Adriatic Sea, Trieste
<i>Hydractinia fucicola</i> (Sars, 1857)	Hermit crabs	Schuchert 2008	Western Mediterranean, Southern Brittany
<i>Hydractinia carca</i> Bergh, 1887	Hermit crabs	Schuchert 2008	European coasts and Japan Sea
<i>Hydrocolella africana</i> Stechow, 1921	<i>Cibbanarius</i> sp., <i>Dardanus arrosor</i> , <i>Diogenes brevisirostris</i> Stimpson, 1858, <i>D. costatus</i> , <i>Pagurus cuanensis</i> Bell, 1845, <i>Pagurus pilicornis</i> Stebbing, 1924	Millard 1975; Balls 1924	(Orange river mouth to Durban, South Africa)
<i>Janaria mirabilis</i> Stechow, 1921	<i>Manucomplanus cervicornis</i> (Benedict, 1892), <i>M. varians</i> (Benedict, 1892) (as <i>Eupagurus varians</i>)	Cairns&Barnard 1984; Bouvier 1898; Ruthensteiner et al. 2008	(Eastern Pacific) (Gulf of California, USA)
<i>Stylactis fucicola</i>	Barnacles (<i>Balanus perforatus</i>) (1)	Boero&Fresi 1986	Portofino, Italy
<i>Stylactis inermis</i>	gastropod shell inhabited hermit crab	Boero&Fresi 1986	Portofino, Italy
<i>Stylactis aculeata</i>	<i>Macrocheira kaempferi</i>	Hiro 1939	(Tonda, Sakai, Wakayama-Ken, Japan)
<i>Stylactis carinicala</i> Hiro, 1939			
<i>Hydractinia pruvoti</i> (Motz-Kossowska, 1905) (as <i>Stylactaria pruvoti</i>)	<i>Cibbanarius erythropus</i> (Latreille, 1818)	Bavestrello et al. 2000	
<i>Hydractinia spiralis</i> (Hirshita, 1988) (as <i>Stylactaria spiralis</i>)	<i>Pagurus constans</i>	Bouillon et al. 1997	(western Mediterranean)
<i>Stylactaria</i> sp.	<i>Xanthias haswellii</i> Calman, 1911	Bouillon 1994	
<i>Podocoryne selena</i> Mills, 1976	<i>Pagurus longicarpus</i> , <i>Pagurus pollicaris</i> ,	Mills 1976; Brooks&Mariscal 1986	(Gulf of Mexico); (northeastern Gulf of Mexico)
	<i>Cibbanarius vittatus</i> (Bosc, 1802), <i>P. longicarpus</i> , <i>P. pollicaris</i>	Brooks&Mariscal 1985; Mills 1976	(northeastern Gulf of Mexico); (Gulf of Mexico)
<i>Podocoryne areolata</i> (Alder, 1862)	<i>Hyas coarctatus</i> , <i>Anapagurus laevis</i> , <i>Pagurus bernhardus</i>	Evans 1978; Edwards 1972	(north east England, UK)

<i>Podocoryne borealis</i> (Mayer, 1900)	<i>Anapagurus laevis</i> (Bell, 1945), <i>Pagurus bernhardus</i>	Edwards 1972	(Clyde Sea Area, south west Scotland)
<i>Hydractinia carnea</i> (M. Sars, 1846) (as <i>Podocoryne carnea</i>)	<i>Anapagurus laevis</i> , <i>Dardanus arrosor</i> , <i>Paguristes eremita</i> , <i>Pagurus bernhardus</i> , <i>P. longicarpus</i> , <i>P. palliaris</i> , <i>P. cuanensis</i> Bell, 1845, <i>P. prideaux</i> Leach, 1815, <i>P. pubescens</i> Krøyer, 1838	Edwards 1972; Cudras&Pereira 1977; Balis 1924; Stachowitsch 1980; Cunningham et al. 1991; McLean 1983	(Clyde Sea Area, south west Scotland); (Mediterranean Sea); (North Adriatic Sea, Gulf of Trieste); (North Atlantic areas and Gulf of Mexico)
	<i>Pagurus</i> sp.	Schijsma 1935; Cazaux 1961; Bennet 1962; Rees 1967; Christensen 1967; Mills 1976; Mercanod&Lytle 1980	(Krhåken Reef in the Øresund, Denmark); (Gulf of Mexico); (North American Coast Atlantic Ocean)
<i>Podocoryne exigua</i> (Haeckel, 1879)	<i>Diogenes pugilator</i> (Roux, 1829); <i>Paguristes eremita</i> (Linnaeus, 1767)	Cerrano et al. 1998; Savestrello 1985	Ligurian Sea
<i>Podocoryne</i> sp.	<i>Pagurus bernhardus</i>	Matthews 1959	(Côte d'Ivoire, Addah)
<i>Polyhydra spinifera</i> Stechow, 1962	<i>Diogenes pugilator</i>	Ruthensteiner et al. 2008	
<i>Leuckartaria octona</i> (Fleming, 1823) (as <i>Perigonimus repens</i>)	<i>Pagurus bernhardus</i>	Kramp 1935 Jensen&Bender 1973	
<i>Perigonimus</i> sp.	<i>Paguristes eremita</i> , <i>Pagurus cuanensis</i>	Stachowitsch 1980	(North Adriatic Sea, Gulf of Trieste)
<i>Bougainvillia ramosa</i> (van Beneden, 1844)	Barnacles (<i>Balanus perforatus</i>) (1)	Boero&Fresi 1986	(Portofino, Italy)
<i>Bougainvillia meinertiae</i> Jäderholm, 1923	<i>Ceratothoa imbricata</i> (Fabricius, 1775)	Millard 1975	(Cape Agulhas, Southern Africa)
<i>Bougainvillia rugosa</i> Clarke, 1882	Libinia	Calder 1971	(Chesapeake Bay, USA)
<i>Bougainvillia supercilialis</i> (L. Agassiz, 1849)	various crustaceans	Aral&Brinckmann-Voss 1980	(British Columbia and Puget Sound)
<i>Bougainvillia</i> sp.	<i>Paguristes eremita</i> , <i>Pagurus cuanensis</i>	Stachowitsch 1980	(North Adriatic Sea, Gulf of Trieste)
<i>Turritopsis nutricula</i> (McCrary, 1857)	Barnacles (<i>Balanus perforatus</i>) (1)	Boero&Fresi 1986	Portofino, Italy
<i>Turritopsis nutricula</i> McCrary, 1857	<i>Balanus improvisus</i>	Calder 1971	(Chesapeake Bay, USA)
<i>Tabularia larynx</i> Ellis and Solander, 1786	Barnacles (<i>Balanus perforatus</i>) (1)	Boero&Fresi 1986	Portofino, Ligurian Sea, Italy
<i>Tabularia mesembryanthemum</i> Allman, 1871	<i>Macrocheira kaempferi</i>	Hiro 1939	(Tonda, Sakai, Wakayama-Ken, Japan)
<i>Boullonia denhartogi</i> Svoboda, Stepanjants & Ljubenkov, 2006	<i>Paralomis formosa</i> Henderson, 1888	Svoboda et al. 2006	Portofino, Ligurian Sea, Italy
<i>Ectopleura larynx</i> (Ellis and Solander, 1786)	Barnacles (<i>Balanus perforatus</i>) (1)	Boero&Fresi 1986	Portofino, Ligurian Sea, Italy
<i>Ectopleura dumortieri</i> (van Beneden, 1844)	<i>Balanus improvisus</i>	Calder 1971	(Chesapeake Bay, USA)
<i>Clovecinia multitecuculata</i> Millard, 1975	<i>Dardanus arrosor</i>	Millard 1975	(False Bay and Natal, Southern Africa)
<i>Corymorpha</i> (<i>Euphyssa</i>) <i>balsi</i> Stechow, 1932	<i>Schiophrys dama</i> (Herbst 1804)	Ruthensteiner et al. 2008	(Heirsson Prong, Shark Bay, Western Australia)
<i>Sarsia phlogaster</i>	Barnacles (<i>Balanus perforatus</i>) (1)	Boero&Fresi 1986	Portofino, Italy
<i>Sarsia tubulosa</i> M. Sars, 1835 (as <i>Sarsia sarsii</i>)	<i>Balanus</i>	Genzano&Rodriguez 1998	(Mar del Plata, Argentina)
<i>Antennella africana</i> Broch, 1914	<i>Macropodia falcifera</i> (Stimpson, 1857)	Millard 1975	(Luderitz Bay to Natal, South Africa)
<i>Gattya humilis</i> Allman, 1885	<i>Dehaanuis dentatus</i> (H. Milne Edwards, 1834)	Millard 1975	(Natal, Southern Africa)
<i>Hydracthis sarcotretis</i> (Jungersen, 1913) (as <i>Ichthyocodium sarcotretis</i>)	<i>Sarcotretes scopeli</i> Jungersen, 1911	Bouillon 1994	Northern Atlantic coasts
	<i>Sphyrian lumpi</i> (Krøyer, 1845)	Bouillon 1994	
<i>Rosalinda incrustans</i> (Kramp, 1947)	<i>Anamothia rissanoa</i> (Roux, 1828)	Schuchert 2010	(Atlantic Ocean, west of Gibraltar; western Mediterranean: Costa Brava and Corsica)
class Anthozoa, subclass Octocorallia, order Alcyonacea			
Octocorallia			
Xenidae			
<i>Xenia</i>	<i>Cryptodromiopsis unidentata</i> (Ruppell, 1830)	Lavaley&Hartog 1995	(Indo-West-Pacific, Indonesia)
<i>Alcyonium digitatum</i> Linnaeus, 1758	<i>Cyclocloma tuberculata</i> Miers, 1880	Hartog 1990	
	<i>Pisa armata</i> , <i>Pisa tetraodon</i>	Carrington&Lovett 1881	
<i>Nemanthus annamensis</i> Carlgren, 1943	<i>Lauridromia intermedia</i> (Laurie, 1906)	Lavaley&Hartog 1995	(Kenyan coast, Africa)
<i>Anemoma sulcata</i> (Pennant, 1777)	<i>Inachus phalangium</i> (Fabricius, 1775), <i>Macropodia rastrata</i> (Linnaeus, 1761), <i>Dromia marmorata</i> Forst., 1814, <i>Stenocypus furcata</i> (Olivier, 1793)	Hartnoll 1971 Lavaley&Hartog 1995; Hartog 1990	(Canary Islands)
<i>Phymactis clematis</i> (Drayton in Dana, 1846)	<i>Allotopetrolithes spinifrons</i> (H. Milne Edwards, 1837)	Baeza&Stotz 2003	(La Pampilla, Peninsula of Coquimbo, north-central Chile)
<i>Sagartia expansa</i> (Stimpson, 1856) (as <i>Cancrisia expansa</i>)	<i>Daripoides fackhina</i> (Herbst, 1785)	Lavaley&Hartog 1995	(SE Asia)
<i>Phlyctenanthus australis</i> Carlgren, 1949	<i>Libinia spinosa</i> H. Milne-Edwards, 1834	Acuña et al. 2003	(Mar del Plata, Buenos Aires province, Argentina)
<i>Isanthus homalophilus</i> Chintiroglou & Doumenq, 1998	<i>Lamoha inflata</i> (Guinot & Richer de Forges, 1981) (as <i>Hypsophrys inflata</i>)	Chintiroglou&Doumenq 1998	(French Polynesia)
<i>Diadumene cincta</i> Stephenson, 1925	<i>Hyas araneus</i>	Ingle 1996	
<i>Urticina felina</i> (Linnaeus, 1761)	<i>Hyas araneus</i>	Ingle 1996	
<i>Urticina consors</i> Verrill, 1882	<i>Parapagurus pilosimanus</i> S. I. Smith, 1879; <i>Sympagurus pictus</i> S. I. Smith, 1883	Balis 1924	
<i>Aiptasia</i> sp.	<i>Paguristes eremita</i> , <i>Pagurus cuanensis</i>	Stachowitsch 1980	(North Adriatic Sea, Gulf of Trieste)
<i>Neoaipasia commensalis</i> Parulekar, 1969	<i>Cibanarius padavensis</i> De Man, 1888, <i>Diogenes custos</i> (Fabricius, 1798)	Parulekar 1969	(Andaman and Nicobar Islands, western India)
<i>Actinia equina</i> (Linnaeus, 1758)	<i>Cancer pagurus</i>	Ingle 1987	(Plymouth, UK)
	<i>Paguristes eremita</i>	Balis 1924	
	<i>Cancer pagurus</i>	Holt 1890	
<i>Triactis producta</i> Klunzinger, 1877 (as <i>Actinia prehensia</i>)	<i>Lybia tessellata</i> (Latreille, in Milbert, 1812)	Karpulis et al. 1998	(Island of Oahu, Hawaii, USA)
<i>Stylobates aeneus</i> Dall, 1903	<i>Sympagurus dofeigni</i> (Balis, 1912)	Dunn et al. 1980	(Hawaii, Guam, USA)
<i>Stylobates cancrisocia</i> (Carlgren, 1928)	<i>Sympagurus trispinosus</i> (Balis, 1911)	Dunn et al. 1980	(off east Africa)
<i>Stylobates loisetoeae</i> Fautin, 1987	<i>Parapagurus</i> sp.	Fautin 1987	(western Australia)
<i>Bunodactis chrysobothrys</i> Parry, 1951	<i>Leptomithrax longipes</i> (Thomson, 1902)	Hand 1975	(New Zealand)
<i>Hormathia coronata</i> (Gosse, 1858)	<i>Dardanus arrosor</i>	Balis 1924	
	<i>Dardanus pedunculatus</i> (Herbst, 1804), <i>D. arrosor</i> , <i>D. deformis</i> (H. Milne-Edwards, 1836), <i>D. impressus</i> (De Haan, 1849), <i>D. isopoides</i> (Forsk., 1775), <i>D. tricolor</i> (Forsk., 1775), <i>D. sinistripes</i> (Stimpson, 1859), <i>D. gemmatum</i> (H. Milne-Edwards, 1848), <i>D. megistos</i> (Herbst, 1804), <i>Anapagurus polynesiensis</i> (Nobili, 1906), <i>Catapagurides fragilis</i> (Melin, 1939), <i>Parapagurus pilosimanus</i> , <i>Cryptodromiopsis saraburei</i> (M. J. Rathbun, 1910)	Imafuku et al. 2000; Carlgren 1949; England 1971; Dales 1957; Yusa et al. 2001; Lavaley&Hartog 1995; Ross 1970; Pauly et al. 2003	(Shirahama, Wakayama Prefecture, Japan) (South Africa) (Muruoa Atoll Tuamotu, Polynesia) (western Japan) (Hawaii Islands) (Mariana Islands)
<i>Callioctis polytypus</i> (Forsk., 1775)	<i>Pagurus bernhardus</i>	Balis 1924	
<i>Callioctis parasitica</i> (Couch, 1844) (as <i>Callioctis effoeta</i>)	<i>Dardanus arrosor</i>	Carlgren 1949; Yusa et al. 2001	(Central Kuroshio Current, East and southern China, east sea Japan) (western Japan)
<i>Callioctis japonica</i> Carlgren, 1928	<i>Dardanus arrosor</i>	Dales 1957	
<i>Callioctis algaensis</i> Carlgren, 1938	<i>Dardanus arrosor</i>	Dales 1957	
<i>Callioctis conchicola</i> Parry, 1952 (as <i>Callioctis conchicola</i>)	<i>Diacanthus rubricatus</i> (Henderson, 1888), <i>Paguristes subpilosus</i> Henderson, 1888, <i>Leptomithrax longipes</i>	Hand 1975	(New Zealand)
	<i>Paguristes barbatus</i> (Heller, 1862), <i>Areopaguristes pilosus</i> (H. Milne Edwards, 1836) (as <i>Paguristes pilosus</i>), <i>Diacanthus spinulimanus</i> (Miers, 1876) (as <i>Pagurus spinulimanus</i>)	Schembri&McLay 1983	(Otago region, southeastern New Zealand)
<i>Callioctis parasitica</i>	<i>Pagurus bernhardus</i>	Reiss et al. 2003; Ingle&Christiansen 2004	(North Sea)
	<i>Cibanarius erythropus</i> , <i>Dardanus arrosor</i> , <i>Paguristes eremita</i> , <i>Pagurus cuanensis</i> , <i>Pagurus alatus</i> (Fabricius, 1775), <i>Pagurus excavatus</i> , <i>Pagurus cuanensis</i> , <i>Pagurus</i> sp.	Ross&Sutton 1961; Cudras&Pereira 1977; Ross 1971; Christidis et al. 1997; Faurot 1932 Chintiroglou&Doumenq 1991	(Adriatic Sea, Aegean Sea, Celtic Seas, North Sea, South European Atlantic Shelf, South India and Sri Lanka) (Mediterranean Sea)
<i>Callioctis tricolor</i>	<i>Pagurus palliaris</i> (shell), <i>Cibanarius vittatus</i> (Bosc, 1802), <i>Dardanus venosus</i> , <i>Pagurus impressus</i> , <i>Pagurus longicarpus</i> , <i>Petrochirus diogenes</i> , <i>P. granulatus</i> Olivier, 1811	Brooks&Mariscal 1986; Cutress&Ross 1969 Brooks&Rittschof 1995; Sandford 2003; Frey 1987; Conover 1976 McLean 1983; Carlgren 1949	(northeastern Gulf of Mexico); (Dog Island, Florida, USA); (Tampa Bay, Florida, USA); (Atlantic ocean: Eastern Atlantic, from all coasts of western Europe and Britain)
	<i>Callinectes danae</i> S. I. Smith, 1869, <i>Hepatus pudibundus</i> (J. F. W. Herbst, 1785), <i>Libinia ferreirei</i> Brito Capello, 1871, <i>L. spinosa</i> , <i>Stenocypus furcata</i> (Olivier, 1791), <i>Hepatus epheliticus</i>	Nogueira et al. 2006; Cutress et al. 1970	(south of Paraná and North of Santa Catarina coast) (Caribbean waters)
	<i>Dardanus jocosus</i> Biffar & Provenzano, 1972	López-Victoria et al. 2004	(Concha Bay, Tayrona Natural Park, Colombian Caribbean)
<i>Callioctis</i> sp.	<i>Paguristes eremita</i> (shell), <i>Dardanus</i> sp., <i>Loxopagurus loxochelis</i> (Moreira, 1901)	Caruso et al. 2003; Cowles 1920, Ayres-Peres&Mantelatto 2010	(Sicilian Channel, Italy) (northern coast of São Paulo, Brazil)
<i>Paracalliactis lacazei</i> Dechance & Dufauré, 1959	<i>Anapagurus laevis</i> (Bell, 1945)	Dechance&Dufauré 1959	
<i>Paracalliactis mediterranea</i> Ross & Zamponi, 1982	<i>Dardanus arrosor</i> , <i>Pagurus variabilis</i> Milne-Edwards & Bouvier, 1892	Ross&Zamponi 1982	(Mediterranean Sea)
<i>Paracalliactis japonica</i> Carlgren, 1928	<i>Dardanus arrosor</i>	Carlgren 1928	
<i>Paracalliactis rosea</i>	<i>Diacanthus rubricatus</i> , <i>Lophopagurus incertus</i> (Henderson, 1888), <i>Paguristes subpilosus</i> , <i>Parapagurus dimorphus</i> (Studer, 1883), <i>Polypagurus crenatus</i> (Borradaile, 1916), <i>Leptomithrax longipes</i>	Hand 1975	(New Zealand)
<i>Paracalliactis rosea</i> Hand, 1975	<i>Paguristes barbatus</i> , <i>Parapagurus dimorphus</i> , <i>Lophopagurus</i>	Schembri&McLay 1983	(Otago region, southeastern New Zealand)
<i>Paracalliactis stephensoni</i> Carlgren, 1928	<i>Parapagurus pilosimanus</i>	Carlgren 1928	
<i>Paracalliactis validiove</i>	<i>Oncopagurus bicristatus</i> A. Milne-Edwards, 1880; <i>Sympagurus andersoni</i> (Henderson, 1896)	Carlgren 1928	
<i>Sagartiogeton undatus</i> (Müller, 1778)	<i>Pagurus excavatus</i>	Chintiroglou et al. 1992	(French Polynesia)
Anthozoa	<i>Dardanus gemmatum</i> , <i>D. deformis</i> , <i>D. megistos</i>	Pauly et al. 2003	(Mariana Islands)
<i>Verrillactis paguri</i> (Verrill, 1869)	<i>Diogenes edwardsii</i> (De Haan, 1849)	Goto et al. 2007	(Seto Inland Sea, Japan)
<i>Sagartiormorpha carlgreni</i> Kwietniewski, 1898 (as <i>Sagartiormorpha paguri</i>)	<i>Dardanus</i> sp., <i>Dardanus deformis</i> , <i>D. gemmatum</i> , <i>Diogenes edwardsii</i>	Cutress&Ross 1969; Haig&Ball 1988; Goto et al. 2007	(Australia and eastern Indonesian waters) (Seto Inland Sea, Japan)
<i>Sagartiormorpha</i> sp.	<i>Diogenes edwardsii</i>	Ross 1983	
<i>Phymanthea pluvia</i> (Drayton in Dana, 1846)	<i>Allotopetrolithes spinifrons</i> (H. Milne-Edwards, 1837)	Baeza&Stotz 2003	(La Pampilla, Peninsula of Coquimbo, north-central Chile)
<i>Bolocerades</i>	<i>Lilia</i>	Doumenq&Van Praët 1987	
<i>Antholoba achates</i> (Drayton in Dana, 1846)	<i>Libinia spinosa</i> , <i>Hepatus chilensis</i> H. Milne-Edwards, 1837	Acuña et al. 2003	(Mar del Plata, Buenos Aires province, Argentina)
<i>Paranthus rapiformis</i> (Le Sueur, 1817)	<i>Pontania domestica</i> Gibbs, 1850	Strent&Chace 1995	(south Texas Coast, USA)
subclass Hexacorallia, order Actiniaria			
<i>Gonactinia prolifera</i> (Sars, 1835)	<i>Anapagurus chiroacanthus</i> , <i>Pagurus bernhardus</i> , <i>P. cuanensis</i>	Samuelsen 1970 ; Jensen&Bender 1973	(Raunefjorden, Western Norway)
subclass Hexacorallia, order Zoantheida			
<i>Epizoanthus abyssorum</i> Verrill, 1885	<i>Parapagurus pilosimanus</i>	Ates 2003	(northern coast, USA)
<i>Epizoanthus oreanensis</i> Delle Chiaje, 1823	<i>Paguristes eremita</i> , <i>Pagurus cuanensis</i>	Ates 2003	(Adriatic Sea, Aegean Sea, Celtic Seas, Western Mediterranean)
<i>Epizoanthus carolinianus</i> Carlgren, 1923	<i>Parapagurus bowyeri</i> Stebbing, 1910, <i>P. pilosimanus</i>	Carlgren 1928; Balis 1924; Ates 2003	(South Africa); (Agulhas Bank Area); (South Africa)
<i>Epizoanthus chuni</i> Carlgren, 1923	<i>Parapagurus arcuatus</i>	Balis 1924; Ates 2003	(East African Coral Coast)
<i>Epizoanthus frenzelii</i> Pax, 1932	<i>Paguristes eremita</i>	Ates 2003	(Adriatic Sea)
<i>Epizoanthus incrustatus</i> (Dubé & Koren, 1847)	<i>Anapagurus laevis</i> , <i>Catapagurus sharreri</i> H. Milne-Edwards, 1880, <i>Pagurus bernhardus</i> , <i>P. excavatus</i> , <i>P. pallius</i> (S. I. Smith, 1882), <i>P. pubescens</i> , <i>Sympagurus pictus</i>	Muirhead et al. 1986; Balis 1924; Ates 2003	(north-east Atlantic); (Celtic Seas, North Sea, Scotian Shelf, South and West Ireland, Southern Norway, Virginian coast)
<i>Epizoanthus indicus</i> Lwowsky, 1913	<i>Parapagurus</i> sp.	Ates 2003	(Japan)

<i>Epizoanthus paguricola</i> (Roule, 1900)	<i>Anapagurus chiroacanthus</i> , <i>A. laevis</i> , <i>Pagurus cuanensis</i> , <i>Simpagurus dimorphus</i>	Roule 1900; Ates 2003 Schjeller&Mantelatto 2009 (southwestern Atlantic Ocean)	(Mediterranean Sea) (Western Mediterranean, Western and Northern Madagascar)
<i>Epizoanthus paguriphilus</i> Verrill, 1882	<i>Pagurus alatus</i> , <i>Parapagurus pilosimanus</i>	Urzelai et al. 1990; Muirhead et al. 1986; Bals 1924	(Cap Breton, Biscay Bay) (north-east Atlantic)
<i>Epizoanthus pagurapsidis</i> Ates, 2003	<i>Paguropsis typica</i> Henderson, 1888	Boas 1926; Ates 2003	
<i>Epizoanthus parasiticus</i> Hertwig, 1882	<i>Pagurus pubescens</i>	Ates 2003	
<i>Epizoanthus senegambiensis</i> Pax & Muller, 1956	<i>Diogenes ovatus</i> Miers, 1879	Ates 2003	(senegambian coast)
<i>Epizoanthus steueri</i> Carlgren, 1923	<i>Paguristes eremita</i>	Ates 2003	(Adriatic Sea)
<i>Epizoanthus studeri</i>	<i>Parapagurus dimorphus</i>	Carlgren 1938; Ates 2003	
<i>Epizoanthus vatovai</i> Pax & Lochter, 1935	<i>Paguristes eremita</i>	Ates 2003	(Adriatic Sea)
<i>Epizoanthus</i> sp.	<i>Anapagurus pusillus</i> Henderson, 1888, <i>Nematopagurus muricatus</i> (Henderson, 1896), <i>Oncopagurus minutus</i> (Henderson, 1896), <i>Paguristes balanophilus</i> Alcock, 1905, <i>P. puniceus</i> Henderson, 1896, <i>Parapagurus abyssorum</i> , <i>P. andreui</i> Macpherson, 1984, <i>P. bouvieri</i> , <i>P. latimanus</i> Henderson, 1888, <i>P. pilosimanus</i> , accepted as <i>Sympagurus burkenroadi</i> Thompson, 1943 (as <i>Sympagurus papposus</i>), <i>S. villosus</i> Lemaitre, 1996	Bals 1924; Ates 2003; Lemaitre 1996	(Indian coasts, Indonesia, French Polynesia, South Africa) (Australia)
<i>Palythoa nelliae</i> Pax, 1935	<i>Cryptodramiopsis unidentata</i> (Ruppell, 1830)	Lavaleye&Hartog 1995	(South Africa)
<i>Palythoa eupaguri</i> Marion, 1882	<i>Parapagurus pilosimanus</i> Smith, 1879 (ex <i>Eupagurus jacobii</i>)	Ates 2003	(Atlantic Ocean)
	subclass Hexacorallia, order Corallimorpharia		
<i>Discosoma</i>	<i>Cyclocoeloma tuberculata</i> Miers, 1880	Lavaleye&Hartog 1995 (Indo-West-Pacific, Indonesia)	(Indo-West-Pacific, Indonesia)