Peter Bugiel, Stanisław Wędrychowicz, and Beata Rzepka*

A few problems connected with invariant measures of Markov maps - verification of some claims and opinions that circulate in the literature

<https://doi.org/10.1515/anona-2020-0221>

Received November 9, 2019; accepted February 4, 2020.

Abstract: It is well known that C^2 -transformation φ of the unit interval into itself with a Markov partition (2.1) $\pi = \{I_k : k \in K\}$ admits φ -invariant density g ($g \geq 0$, $\|g\| = 1$) if: (2.2) $|(\varphi^n)'| \geq C_1 > 1$ for some n (*expanding condition*); (2.3) $|\varphi''(x)/(\varphi'(y))^2| \leq C_2 < \infty$ (*second derivative condition*); and (2.4) $\#\pi < \infty$ or $\varphi(I_k) = [0, 1]$, for each $I_k \in \pi$. If (2.4) is deleted, then the situation dramatically changes. The cause of this fact was elucidated in connection with so-called *Adler's Theorem* ([1] and [2]).

However after that time in the literature occur claims and opinions concerning the existence of invariant densities and their properties for Markov Maps, which satisfy (2.2), (2.3) and do not satisfy (2.4), revealing unacquaintance with that question.

In this note we discuss the problems arising from the mentioned claims and opinions. Some solutions of that problems are given, in a systematic way, on the base of the already published results and by providing appropriate examples.

Keywords: Markov maps; invariant measure; expanding transformation; recurrence; aperiodicity

MSC 2010: 28D05, 37A99, 26A18

1 Introduction

We begin with the celebrated result of [3]. The authors were well aware that their result cannot be extended to expanding transformations with countably many one-to-one pieces in a simple way (see Th. 2, and the comment below on Cond. (17) there). The real task in that period of time was to find reasonable additional conditions which would guarantee the existence of density invariant under the action of expanding map with countably many one-to-one pieces. Several attempts was made to accomplish that task (for more details see e.g. a review article [4], and also [5], or [6], Sect. 6).

One of the mentioned attempts was published in [7], as *Adler's Theorem*. Since no proof was given there, the question arose whether it is true [8]. A solution was published in [1], and [2].

After the above two notes and a few other ones, related with them, were published, some further claims and opinions concerning the existence of invariant densities and their lower and upper bounds for Markov Maps appear in the literature.

Peter Bugiel, Faculty of Mathematics and Computer Science, Jagiellonian University Cracow (Kraków), Poland,

E-mail: uibugiel@cyf-kr.edu.pl

Stanisław Wędrychowicz, Department of Nonlinear Analysis, Rzeszów University of Technology, al. Powstańców Warszawy 8, 35-959 Rzeszów, Poland, E-mail: swedrych@prz.edu.pl

***Corresponding Author: Beata Rzepka**, Department of Nonlinear Analysis, Rzeszów University of Technology, al. Powstańców Warszawy 8, 35-959 Rzeszów, Poland, E-mail: brzepka@prz.edu.pl

Those claims and opinions reveal that their authors were unacquainted with the essence of the problem. That problem is rather of delicate nature. It involves, among other things, the so-called *measure-theoretic recurrence property*.

In this note we clear up, in a systematic way, the essence of the problems with the aid of examples, comments and some published results.

2 The problems and examples

2.1 Existence of invariant densities

We begin with the following

Definition 2.1. Let $\varphi : I \rightarrow I$ be countably piecewise one-to-one and C^2 , here $I \subset \mathbb{R}$ denotes an interval. It is called of *Markov type* if there exists a partition (mod 0) $\pi = \{I_k : k \in K\}$ such that: each I_k is an interval, $\varphi_k := \varphi|_{I_k}$ is one-to-one and C^2 from I_k onto $J_k = \varphi(I_k)$ and the following condition holds:

$$\text{for each } j, k \in K, \text{ if } \varphi(I_j) \cap I_k \neq \emptyset, \text{ then } I_k \subseteq \varphi(I_j). \quad (2.1)$$

The following result follows from Ths. 1, and 2 in [3]:

Corollary 2.2. Assume that φ is of Markov type and satisfies:

$$\text{for some } n : \quad (\varphi^n)'(x) \geq C_1 > 1 \text{ whenever } (\varphi^n)' \text{ is defined}; \quad (2.2)$$

$$|\varphi''(x)/(\varphi'(y))^2| \leq C_2 < \infty \text{ whenever } \varphi''(x) \text{ and } \varphi'(y) \text{ are defined}; \quad (2.3)$$

$$\text{and } \#\pi < \infty \text{ or } \varphi(I_k) = [0, 1], \text{ for each } I_k \in \pi. \quad (2.4)$$

Then:

$$\text{there exists } \varphi\text{-invariant density } g \text{ } (g \geq 0, \|g\| = 1). \quad (2.5)$$

Now we delete (2.4) and pose, at the beginning, the following question:

Problem 2.3. Assume that φ is of Markov type, in the sense of Def. 2.1, and satisfies: Conditions (2.2), and (2.3). Does (2.5) holds?

The following simple example gives negative answer to the question in Problem 2.3:

Example 2.4. Let $I_k = [1 - 2^{-(k-1)}, 1 - 2^{-k}]$ ($k = 1, 2, \dots$) and $\varphi : [0, 1] \rightarrow [0, 1]$ such that every I_k is mapped linearly on $I_k \cup I_{k+1}$. Then $|\varphi'| \geq 3/2$, φ is of Markov type and fulfils (2.3). Nevertheless (2.5) does not hold because each point moves to the right under the action of φ .

Notice that φ in this example has the following defective property:

$$\bigcup_{j=1}^{\infty} \varphi^j(I_k) = \bigcup_{j=k}^{\infty} I_j \supsetneq \emptyset \text{ as } k \rightarrow \infty, \text{ for } k \geq 2.$$

The transformation given in [9] is another example which gives negative answer to the question in Problem 2.3.

Let us now eliminate that defect by imposing the so-called indecomposability condition (see [10], Definitions 2.2, 2.3, and Cond. 2.[M₁4]):

$$\bigcup_{j=1}^{\infty} \varphi^j(I_k) = [0, 1], \quad \text{for each } I_k \in \pi. \quad (2.6)$$

Note that indecomposability condition (2.6) is equivalent to the following so-called *transitivity condition*:

$$\text{for every } I_i, I_j \in \pi \text{ one has: } I_i \subset \varphi^n(I_j) \text{ for some } n \geq 1. \quad (2.7)$$

Definition 2.5. A transformation φ of Markov type which satisfies Cond. (2.6) is called a *Markov Map*.

Now we pose, analogously as before, the following question:

Problem 2.6. Assume that φ is of Markov type, in the sense of Def. 2.1, and satisfies Conditions: (2.2), (2.3) and (2.6).

Does (2.5) holds?

In this case the following example gives negative answer to the question in Problem 2.6 [11]:

Example 2.7. Let $A = \{0\} \cup \{1/k : k = 1, 2, \dots\}$ and $I_k = (1/(k+1), 1/k]$ for $k = 1, 2, \dots$. Then we define φ as follows:

$$\begin{aligned} \varphi|_{I_1}(x) &= 2x - 1; \\ \text{and for any } k = 2, 3, \dots, \varphi|_{I_k} &\text{ is the increasing linear such that} \\ \varphi(I_k) &= (0, 1/(k-1)]; \\ \varphi(0) &= 0. \end{aligned}$$

Proof. Clearly, φ is of Markov type and satisfies all the assumptions of Problem 2.6. Nevertheless there is no φ -invariant density. This is so because it is ergodic and has σ -finite absolutely invariant measure concentrated on the whole I (there exists piecewise constant, not integrable, and φ -invariant function). $\square$

Notice that φ in this example fulfils even more restrictive condition than Cond. (2.6). Namely, it fulfils

$$\varphi^k(I_k) = (0, 1] \quad \text{for each } I_k \in \pi. \quad (2.8)$$

Finally, let us consider instead of Cond. (2.6) its more restrictive version; the following condition:

$$\text{there exists } j \geq 2 \text{ such that } I_i \subset \bigcup_{r=1}^j \varphi^r(I_k) \text{ for every } I_i, I_k \in \pi. \quad (2.9)$$

Once more we pose the following question:

Problem 2.8. Assume that φ is of Markov type in the sense of Def. 2.1, and satisfies Conditions: (2.2), (2.3) and (2.9).

Does (2.5) holds?

The so-called *Adler's Theorem* asserts that the answer to the question in Problem 2.8 is positive [7]. But no proof is given there. Further, the comments on that theorem in [12] are restricted to a history of the theorem. However, it was noted that *Adler's Theorem* may not hold in [8].

The counterexamples, published in [1] and [2], disprove *Adler's Theorem*, i.e. they give negative answer to the question in Problem 2.8.

In the former paper was also proposed a correction of *Adler's Theorem*. Namely, in the case of bounded interval I the following additional condition was proposed (see Cond. (1.H3) there):

$$\inf\{|\varphi(I_k)| : k \in K\} > 0; \quad \text{where } |\cdot| \text{ denotes Lebesgue measure.} \quad (2.10)$$

While in the case of unbounded interval I , it was proposed the following (see Cond. (1.H4) there):

$$\lim_{n \rightarrow \infty} R(n) = 0 \text{ where } R(n) = \sup_{k \in K} \int_{I \setminus V_n} \tilde{\sigma}_k(x) dx, \quad \tilde{\sigma}_k(x) = \frac{\sigma_k(x)}{|I_k|}, \quad (2.11)$$

$$\sigma_k(x) = |(\varphi_k^{-1})'(x)| \mathbf{1}_{\varphi(I_k)}(x), \quad \{V_n\}_{n=1}^{\infty} \text{ is a sequence of subsets of } I,$$

and each V_n is the union of a finite number of I_k 's such that $V_n \subset V_{n+1}$,

$$\bigcup_{n=1}^{\infty} V_n = I \pmod{|\cdot|}.$$

Note that under the assumptions of Problem 2.8 the two Conditions (2.10) and (2.11) are equivalent.

A more efficient than the last two above conditions is the following one:

$$\int h_1 dx > 0 \text{ where } h_1 = \inf_s \sum_{k \in K} \tilde{\sigma}_k \int_{I_k} \tilde{\sigma}_s(x) dx, \quad (2.12)$$

and $\tilde{\sigma}_k$ is defined in (2.11).

Note that Condition (2.12) is an analogue of the widely known condition from the theory of Markov Chain, the analogue is explained in [13]. Its efficiency is illustrated by examples in ([13], Ex. 2.1) and in ([6], Exs. 4.3.1, and 4.3.2). One has also to underline, that

Remark 2.9. Condition (2.12) additionally assures aperiodicity but Condition (2.10) does not (see Example 2.13, below).

The role which each of the last three conditions plays in the problem of the existence of invariant density consists in guaranteeing that the needed *measure-theoretic recurrence property* holds.

Since the transformations given in Example 2.4 and in [9] have global attractors (single point and the Cantor set, respectively), they are without that property. Note also that they do not satisfy the simple Condition (2.10).

On the other hand, it is not easy to decide, without Condition (2.12), whether or not the above mentioned transformations of Examples 4.3.1, or 4.3.2 in [6] have the needed *measure-theoretic recurrence property*.

Theorems stated in [14] as Theorem 1.2 and, in more abstract setting, as Theorem 1.3 contain the theorem questioned in [8].

There is also given a proof of Th. 1.3 which is incorrect (the thesis of the Lemma 1.5 does not hold, in general). That fact is not noted in [15].

Theorem 2.2 in [16] is a version of Th. 1.3 from [14]. It is stated under the indispensable Condition (2.10). This condition is incorporated, as Condition c), in the definition of Markov Map (Definition p. 353).

However, in connection with the Assertion c) of that theorem and the opinion on transitivity Assumption contained in Remark 4c) p. 354, here Condition (2.7), one has to raise two questions. The first question reads:

Problem 2.10. Assume that φ is of Markov type in the sense of Def. 2.1. What is the essential role played by Condition (2.7) in the theory of Markov Maps?

We begin with example of a Markov type map without property (2.7) (see also [10], Example 2.1, *W-transformation*):

Example 2.11. Let $0 < a < 1$, and then let $\psi : I = [0, 1] \rightarrow I$ be defined by

$$\psi(x) = \begin{cases} x/a & \text{if } 0 \leq x < a, \\ -(1+a)x + (a^2 + a + 1) & \text{if } a \leq x < 1. \end{cases}$$

Proof. Clearly, ψ is a transformation of Markov-type with respect to the following intervals: $I_1 = [0, a^2)$, $I_2 = [a^2, a)$, $I_3 = [a, 1)$.

The interval I_1 is the so-called inessential interval; the remainder two intervals I_2, I_3 are essential [10]. Transformation ψ restricted to the last two intervals satisfies already condition (2.7). Consequently, the invariant density is supported by $I_2 \cup I_3$. $\square$

In general, any transformations of M-type can be decomposed into transformations with property (2.7) and an inessential part (see for details [10]).

The second question:

Problem 2.12. Assume that φ is of Markov type in the sense of Def. 2.1 which satisfies: (2.5) and the so-called transitivity condition (2.7).

Does the invariant density is necessarily exact in the sense of Rokhlin ([17])?

As it shows the transformation of the below Example 2.13, *transitivity condition* (2.7) does not assure, in general, exactness. It has to be complemented to *exclude periodicity of Markov Map*. However, as it is noted in Remark 2.9, Condition (2.12) involves already aperiodicity.

The role of such complementary condition plays Condition (1.H5) in ([18], Remark (1.1)) (see also Cond. (3.H₁4) in [13]), or Cond. (4.1.H₁3) in [6]). That condition reads:

$$\text{There exist an integer } \tilde{n} \geq 1 \text{ and } I_{\tilde{k}} \text{ such that } \varphi^{\tilde{n}}(I_{\tilde{k}}) = [0, 1]. \quad (2.13)$$

Note that it is a weak version of Cond. (2.16) below.

Example 2.13. (of a Markov map with properties (2.7) and not exact)

Put $I_k = [k, k+1)$ for $k = 0, 1, 2, 3$. Let $\chi_k : I_k \rightarrow I_2 \cup I_3$ for $k = 0, 1$ be linear, increasing, and onto. Analogously, let $\chi_k : I_k \rightarrow I_0 \cup I_1$ for $k = 2, 3$ be linear, increasing, and onto.

Finally, define $\chi : I = [0, 4) \rightarrow I$ by $\chi(x) = \chi_k(x)$ iff $x \in I_k$.

Then χ trivially fulfils the Conds. (2.2), and (2.3) of Corollary 2.2 and is a Markov Map which satisfies Condition (2.7) (actually Condition (2.9), for $j = 3$) and Condition (2.10), but it is not exact. Therefore it is a counterexample to the Assertions c) and d) of Theorem 2.2 in [16].

The proof of the last fact is based on the following criterion of exactness [17]:

Let $(I, \mathfrak{F}, \varphi : I \rightarrow I; d\mu)$ where I is a space, $\mathfrak{F}$ is a σ -algebra of its subsets, φ transformation with μ invariant measure. Then

$$\varphi \text{ is exact} \Leftrightarrow \text{for every } A \in \mathfrak{F}, \lim_{n \rightarrow \infty} \mu(\varphi^n(A)) = 1. \quad (2.14)$$

Proof. Now we are going to show that χ is not exact. Note first that $d\mu = 1/4 dx$ is the unique invariant density.

Further $\chi(I_k) = I_2 \cup I_3$ for $k = 0, 1$ and $\chi(I_k) = I_0 \cup I_1$ for $k = 2, 3$. From these relations it follows that

$$\mu(\varphi^n(I_k)) = 1/2 \quad \text{for every } I_k, k = 0, 1, 2, 3, \text{ and } n = 1, 2, 3, \dots, \quad (2.15)$$

and therefore the criterion (2.14) is not fulfilled. $\square$

Finally, one needs to complete the opinion on Cond. (2.10) expressed in the Remark 4c), p. 354 of the cited book [16]. The authors claim that it can be somewhat weakened but it is certainly not possible to dispense with it altogether if we want to have that Markov Maps necessarily have invariant density.

The essential role played by Cond. (2.10) for the existence of invariant density has been already explained above.

As for the weakening of that condition, it is in general less efficient than Condition (2.12) (see: Convergence Theorem; Coroll. 1.1 in [18] – 1-dimensional case; or 3.1. Theorem; 3.1. Coroll. in [13] – multi-dimensional case). The efficiency of Condition (2.12) is also shown below by Examples 2.19 and 2.20.

In the introduction of [19] is noted that in Chapter 7, Section 4 of the book [20], in English, the proof appears to have an error.

Actually the theorem contained in Chapter 7, Section 4 of that book does not hold. This is so because the theorem in question is stated under somewhat less restrictive conditions than that of the so-called *Adler's Theorem*. Therefore the above mentioned counterexamples in [1], and [2] disprove that theorem as well.

In a review [21] of the book [22], in Polish, the reviewer claims that the proof of the Theorem 1, § 4, Section 7 on p. 164 is not correct.

This problem is clear up in [23]. It turns out that this is the very same problem as that raised in the introduction of [19].

One has to return to the already mentioned above note [19]. At the end of that note is questioned the double inequality of Remark 1 in [24]. That remark states:

*If countably piecewise C^1 -Markov Map satisfies conditions:
Cond. (2.2, for $n = 1$), Cond. (2.3), and*

$$\text{there exists one integer } \tilde{n} \text{ such that } \varphi^{\tilde{n}}(I_k) = [0, 1] \text{ for each } I_k \in \pi, \quad (2.16)$$

then the invariant measure is unique and its density is bounded away from 0.

The author also claims in ([24], p. 38) that the density g of the unique invariant measure satisfies the following double inequality:

$$M_1 \leq g(x) \leq M_2 \quad \text{for some constants } M_1, M_2 > 0. \quad (2.17)$$

On the other hand, the authors in ([19], p. 868) question the above double inequality (2.17).

Remark 2.14. Additionally the authors suggest that the fault is connected with the use of the idea of regularity functional. This is not the case. The regularity functional has been used to get bounds (see e.g. [6], [13], or [18]).

However, the bounds of the double inequality are, in general, not constants as in (2.17), but functions (see: the above cited papers or Coroll. 2.21, below).

Next if $\tilde{n} = 1$ in (2.16), then the remark in question is obviously correct.

Finally, in connection with the discussed Remark 1 in ([24], pp. 37-38), one has to raise two further questions. The first question is still connected with the problem of the existence of invariant densities for Markov Maps:

Problem 2.15. *Assume that φ is of Markov type in the sense of Def. 2.1 and satisfies: Cond. (2.2) for $n = 1$, Cond. (2.3) and Cond. (2.16) for $\tilde{n} \geq 2$. Does (2.5) holds?*

The second question is connected with the problem of the existence of the lower and upper bounds of invariant density. It is delayed until the second subsection.

As for the question in Problem 2.15, first note that Cond. (2.16) is essentially more restrictive than Cond. (2.9) thus it is a restrictive version of *Adler's Theorem*. Nevertheless, the answer to that question is negative too.

It follows from the repeatedly cited counterexamples published in [1] and [2]. More exactly, the defined in [2] Markov Map $\tilde{\tau} : I = [0, 1] \rightarrow I$ satisfies

$$\tilde{\tau}^2(\tilde{I}_k) = [0, 1] \quad \text{for each } \tilde{I}_k \text{ (see there Final remark (b)).}$$

2.2 Bounds of invariant densities

We begin this subsection with the question announced at the end of the previous one. It can be formulated as follows:

Problem 2.16. Assume that φ is of Markov type in the sense of Def. 2.1 which satisfies Cond. (2.5). Does the invariant density satisfies the double inequality (2.17)?

Regarding the Problem 2.16. As was above noted, the authors in ([19], p. 868) question the above double inequality (2.17) in [24] and suggest that the fault is connected with the use of the idea of regularity functional (see Remark 2.14).

On the other hand, in ([25], Ex. 4) is given an example of Markov Map S in the sense of Definition 2.5 and it is shown that the double inequality (2.17) does not hold.

Remark 2.17.

- (a) More specifically, the Markov Map S satisfies Conditions (2.2), (2.3), and (2.10), above and therefore, as it is proved in ([25], Proposition 2), it belongs to a class considered in [26].

Then the author shows that the invariant density h of the Markov Map S satisfies:

$$\lim_{x \rightarrow 1} h(x) = 0. \quad (2.18)$$

Therefore it does not satisfy Cond. (2.17) (it is not bounded away from 0).

- (b) Note that the relation (2.18) is an immediate consequence of the double inequality of the below Coroll. 2.21.

The author claims that the property (2.18) of S is associated with Cond. (2.10). There is however no argumentation given that this is the case.

Actually, that property of invariant density is neither caused by Cond. (2.10) nor by any other condition that assures existence of invariant density.

Example 4 in [25] illustrates in the reality quite another fact. Namely, it shows that assumption (2.22) in Coroll. 2.22 cannot be omitted.

Indeed, it follows from the following

Proposition 2.18.

- (a) There exist two Markov Maps ψ and $\tilde{\psi}$ which do not belong to the class considered in [26]. Precisely, they do not satisfy Cond. (*) of Proposition 2 in ([25], p. 1274) and therefore, a fortiori, they do not satisfy Cond. (2.10).
- (b) There is ψ -invariant density g_ψ such that:

$$\liminf_{x \rightarrow 1} g_\psi(x) = 0, \quad (2.19)$$

and therefore g_ψ does not satisfy Cond. (2.17) (it is not bounded away from 0).

- (c) There is $\tilde{\psi}$ -invariant density $g_{\tilde{\psi}}$ which satisfies Cond. (2.17).

Proof. The two transformations are given in the following two examples:

Example 2.19. Let $p_k = 1 - 2^{-k}$, and put $I_k = [p_k, p_{k+1})$, for $k = 0, 1, 2, \dots$

Then define linear mappings $\psi_{2k} : I_{2k} \rightarrow [0, p_{2k+3})$ for $k = 0, 1, 2, \dots$;

$\psi_1 : I_1 \rightarrow [0, p_4)$; and $\psi_{2k+1} : I_{2k+1} \rightarrow I_{2k}$ for $k = 1, 2, 3, \dots$

Finally, define $\psi : [0, 1) \rightarrow [0, 1)$ by $\psi(x) = \psi_k(x)$ iff $x \in I_k$.

Example 2.20. Markov Map $\tilde{\psi}$ is a result of simple modification of ψ in such a way that its first linear mappings ψ_0 is replaced with the linear mapping $\tilde{\psi}_0$ from I_0 onto the whole $[0, 1]$.

Now we show that the two Markov Maps ψ and $\tilde{\psi}$ have the properties listed in the proposition. Part (a) of Prop. 2.18 follows directly from definitions of the two Markov Maps.

The proof of the remainder two Parts (b) and (c) is based on the following two corollaries:

Corollary 2.21. Let a Markov map $\varphi : [0, 1] \rightarrow [0, 1]$ satisfy: Conditions (2.2) and (2.3) of Corollary 2.2 and additionally Cond. (2.10) or, the more efficient, Cond. (2.12). Then:

there is a unique φ -invariant density g_0 such that

$$C_1^{-1}\tilde{g} \leq g_0 \leq C_1\tilde{g}, \quad (2.20)$$

where $C_1 > 0$ is a constant and $\tilde{g} > 0$ on $\{g_0 > 0\}$ is given by

$$\tilde{g} = \sum_{k \in K} \tilde{\sigma}_k \int_{I_k} g_0 dx, \quad (2.21)$$

and $\tilde{\sigma}_k$ is defined in (2.11).

Proof. (see: Convergence Theorem; Coroll. 1.1 in [18], 1-dimensional case, or 3.1. Theorem, 3.1. Coroll. in [13], multi-dimensional case). $\square$

The second corollary reads:

Corollary 2.22. *If, in particular, φ satisfies:*

there is $\pi_1 \subset \pi$, $\#\pi_1 < \infty$ such that:

$$\inf_{x \in \varphi(I_k)} \tilde{\sigma}_k(x) > 0 \text{ for } I_k \in \pi_1, \text{ and } \bigcup_{I_k \in \pi_1} \varphi(I_k) = \{g_0 > 0\} \quad (2.22)$$

then there is a constant $C_0 > 0$ such that $g_0 \geq C_0$.

Proof. This fact is a simple consequence of the assumptions of the previous Corollary 2.21 and the double inequality (2.20) together with (2.21). $\square$

The proof of (2.19) consists of two parts. In the first part it is proved that there exists a unique ψ -invariant density g_ψ ; in the second part it is proved that it satisfies (2.19).

To prove the existence of g_ψ we show that ψ satisfies Cond. (2.12). To this end observe that from the inequalities

$$\tilde{\sigma}_{2k}(x) \geq \mathbf{1}_{[0, p_5)}(x), \text{ for } i = 0, 1, 2, \dots, \text{ and } \tilde{\sigma}_1(x) \geq \mathbf{1}_{[0, p_4)}(x) \quad (2.23)$$

where $\tilde{\sigma}_k$ is defined in (2.11), it follows

$$\begin{aligned} \sum_{k=0}^{\infty} \tilde{\sigma}_k \mathbf{1}_{\psi(I_k)} \int_{I_k} \tilde{\sigma}_i \mathbf{1}_{\psi(I_i)} dx &\geq \tilde{\sigma}_1 \mathbf{1}_{\psi(I_1)} \int_{I_1} \tilde{\sigma}_i \mathbf{1}_{\psi(I_i)} dx + \sum_{k=0}^{\infty} \tilde{\sigma}_{2k} \mathbf{1}_{\psi(I_{2k})} \int_{I_{2k}} \tilde{\sigma}_i \mathbf{1}_{\psi(I_i)} dx \\ &\geq \mathbf{1}_{[0, p_4)} \int_{I_1 \cup (\bigcup_{k=0}^{\infty} I_{2k})} \tilde{\sigma}_i \mathbf{1}_{\psi(I_i)} dx > 0, \end{aligned} \quad (2.24)$$

for any $\tilde{\sigma}_i$. It implies Cond. (2.12).

As for (2.19), note first that the density $\tilde{g}$ given by (2.21) of Coroll. 2.21 and associated with g_ψ can be written as a sum

$$\tilde{g} = g_1 + g_2, \quad (2.25)$$

where

$$g_1 = \tilde{\sigma}_1 \mathbf{1}_{\psi(I_1)} \int_{I_1} g_\psi dx + \sum_{i=1}^{\infty} \tilde{\sigma}_{2i+1} \mathbf{1}_{\psi(I_{2i+1})} \int_{I_{2i+1}} g_\psi dx,$$

and

$$g_2 = \sum_{i=0}^{\infty} \tilde{\sigma}_{2i} \mathbf{1}_{\psi(I_{2i})} \int_{I_{2i}} g_\psi dx.$$

Further for g_1 one has

$$g_1 = \frac{1}{p_4} \mathbf{1}_{I_0 \cup I_1 \cup I_2 \cup I_3} \int_{I_1} g_\psi dx + \frac{1}{|I_2|} \mathbf{1}_{I_2} \int_{I_3} g_\psi dx + \sum_{i=2}^{\infty} \frac{1}{|I_{2i}|} \mathbf{1}_{I_{2i}} \int_{I_{2i+1}} g_\psi dx, \quad (2.26)$$

and for g_2 one has

$$\begin{aligned} g_2 &= \sum_{i=0}^{\infty} \frac{1}{|\psi(I_{2i})|} \mathbf{1}_{\psi(I_{2i})} \int_{I_{2i}} g_\psi dx \\ &\leq \frac{1}{p_3} \left(\mathbf{1}_{I_0 \cup I_1 \cup I_2} \int_{A_0} g_\psi dx + \sum_{i=1}^{\infty} \mathbf{1}_{I_{2i+1} \cup I_{2i+2}} \int_{A_{2i}} g_\psi dx \right), \end{aligned} \quad (2.27)$$

where

$$A_{2i} = \bigcup_{k=0}^{\infty} I_{2(i+k)}, \quad \text{for } i = 0, 1, 2, \dots$$

Thus the relation (2.19) follows from (2.20) and (2.21) of Coroll. 2.21, and the relations (2.25), (2.26), and (2.27).

As for the Part (c), note that the simple modification of ψ described in Example 2.20 leads to a Markov Map $\tilde{\psi}$ which satisfies Assumption (2.22) of Coroll. 2.22. Therefore its $\tilde{\psi}$ -invariant density $g_{\tilde{\psi}}$ satisfies Cond. (2.17). $\square$

References

- [1] P. Bugiel, A note on invariant measures for Markov maps of an interval, *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete* **70** (1985), 345–349; Zbl. Math. 606.28013; MR **87a**:28019.
- [2] P. Bugiel, Correction and addendum to: A note on invariant measures for Markov maps of an interval, *Probability Theory Rel. Fields* **76** (1987), 255–256; (formerly: *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*); Zbl. Math. 651.28010; MR **88i**:28028.
- [3] A. Lasota and J.A. Yorke, On the existence of invariant measures for piecewise monotonic transformations, *Transactions of the American Mathematical Society* Vol. **186** (1973), 481–488.
- [4] M. Iosifescu, Mixing properties for f -expansions, in: *Probab. Theory and Math. Stat.*, Prohorov et al. (eds.), VNU Science Press 1987, Vol. **2** (1987), 1–8.
- [5] M. Iosifescu and S. Grigorescu, *Dependence with Complete Connections and its Applications*, Cambridge University Press, Cambridge Tracts in Mathematics **96**, Cambridge, 1991.
- [6] P. Bugiel, Distortion inequality for the Frobenius-Perron operator and some of its consequences in ergodic theory of Markov maps in $\mathbb{R}^d$, *Annales Polonici Mathematici* **LXVIII.2** (1998), 125–157 (pdf copy is available in net).
- [7] R. Bowen, Invariant measures for Markov Maps of the interval, *Commun. Math. Phys.* **69** (1979), 1–14.
- [8] C. Series, Additional comments to Bowen (1979), *Commun. Math. Phys.* **69** (1979), pp. 17.
- [9] R. Rudnicki, On a one-dimensional analogue of the Smale horseshoe, *Annal. Polon. Math.* Vol. **54.2** (1991), 147–153.
- [10] P. Bugiel, On the number of all absolutely continuous, ergodic measures of Markov type transformations defined on an interval, *Math. Nachr.* **129** (1986), 261–268; Zbl. Math. 622.28014; MR **88a**:58111.
- [11] P. Góra, Countably piecewise expanding transformations without absolutely continuous invariant measure, *Dynamical Systems and Ergodic Theory*, Banach Center Publications, PWN - Polish Scientific Publishers, Warszawa Vol. **23** (1989), 113–117.
- [12] R.L. Adler, Afterword to Bowen, *Commun. Math. Phys.* **69** (1979), 15–17.
- [13] P. Bugiel, Ergodic properties of Markov maps in $\mathbb{R}^d$, *Probab. Theory Relat. Fields* **88** (1991), 483–496.
- [14] R. Mañé, *Ergodic Theory and Differentiable Dynamics*, Ergebnisse der Mathematik und ihrer Grenzgebiete, 3.Folge, Bd.8, Springer-Verlag, Berlin, 1987.
- [15] J. Franks, A review of the book: [14], *Bull. Amer. Math. Soc.* **20** (1989), 190–193.
- [16] W. Melo and S. Strien, *One-dimensional Dynamics*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Springer-Verlag, 1993.
- [17] V.A. Rohlin, Exact endomorphisms of Lebesgue spaces, *Izv. Akad. Nauk SSSR Ser. Mat.* **25** (1961), 490–530; *Amer. Math. Soc. Transl. (2)* **39** (1964), 1–36.

- [18] P. Bugiel, On the convergence of the iterates of the Frobenius-Perron operator associated with a Markov map defined on an interval. The lower-function approach, *Annales Polonici Mathematici* **LIII.2** (1991), 131–137 (pdf copy is available in net).
- [19] P. Góra and A. Boyarsky, Compactness of invariant densities for families of expanding, piecewise monotonic transformations, *Can. J. Math.* Vol. **XLI** (1989), no. 5, 855–869.
- [20] I.P. Cornfeld, S.V. Fomin and Ya.G. Sinai, *Ergodic Theory*, Springer-Verlag, New York, 1980 (Grundlehren Math. Wiss. 245).
- [21] H. Żołądek, A review of the book: [22], *Mathematical News (Wiadomości Matematyczne)* **28.2** (1990), 278–280 (in Polish).
- [22] S.W. Fomin, I.P. Kornfeld and J.G. Sinaj, *Ergodic Theory (Teoria Ergodyczna)*, PWN, Warszawa 1987, translated (from the Russian to Polish) by Jacek Jakubowski, pp. 402, ISBN 83-01-07118-4.
- [23] P. Bugiel and S. Wędrychowicz, Remark concerning the review on the book: [22], *Roczniki Polskiego Towarzystwa Matematycznego, Seria II: Wiadomości Matematyczne (Annals of Polish Mathematical Society, Series II: Mathematical News)* **40** (2004) (in Polish) (pdf copy is available in net).
- [24] G. Pianigiani, First return map and invariant measure, *Israel Jour. Math.* **35** (1980), nos 1-2, 32–47.
- [25] R. Zweimüller, Ergodic structure and invariant densities of non-Markovian interval maps with indifferent fixed points, *Nonlinearity* **11** (1998), 1263–1276.
- [26] M. Rychlik, Bounded variation and invariant measures, *Studia Math.* **76** (1983), 69–80.