Semi-free subgroups of a profinite surface group

Matan Ginzburg and Mark Shusterman

Communicated by Pavel A. Zalesskii

Abstract. We show that every closed normal subgroup of infinite index in a profinite surface group Γ is contained in a semi-free profinite normal subgroup of Γ . This answers a question of Bary-Soroker, Stevenson, and Zalesskii

1 Introduction

The classical theorem of Nielsen and Schreier states that every subgroup of a free group is free. Trying to extend this result to profinite groups fails, for example $\mathbb{Z}_2 \leq \hat{\mathbb{Z}}$. This naturally gives rise to the question of finding conditions upon which a subgroup of a free profinite group is free.

Some results in this direction are known, for instance, Melnikov's characterization of normal subgroups of free profinite groups, and Haran's diamond theorem. Results of a slightly different flavor have been obtained by Shusterman in [7], where, for example, the following is proven.

Theorem 1.1. *Let F be a nonabelian finitely generated free profinite group, and let $H \leq_c F$ be a closed subgroup of infinite index in F . Then there exists a free profinite subgroup $H \leq L \leq_c F$ of rank $\aleph_0$.*

In particular, weakly maximal subgroups (see Definition 2.2) are free.

In this work we consider an analog of the above for profinite surface groups. These groups show up as étale fundamental groups of curves over an algebraically closed field of characteristic 0.

We will be interested in semi-free profinite subgroups (of profinite surface groups), a notion introduced in [1], where it is shown that a group is free profinite if and only if it is projective and semi-free. As shown in [8], projectivity of a subgroup of a profinite surface group is equivalent to a simple condition on its index (as a supernatural number). Henceforth, we will focus on semi-freeness.

Our main result is the following.

Theorem 1.2. *Let $N \triangleleft_c \Gamma_g$ be a normal subgroup of infinite index in a profinite surface group of genus $g \geq 2$. Then there exists a semi-free profinite subgroup $M \triangleleft_c \Gamma_g$ that contains N .*

This answers a question raised by Bary-Soroker, Stevenson, and Zalesskii (in [2, Remark 4.1]), who used their diamond theorem to establish the special case where Γ_g/N is not hereditarily just infinite.

Our method also gives the following analog of the aforementioned results of Shusterman.

Theorem 1.3. *Weakly maximal subgroups of profinite surface groups are semi-free profinite.*

Weakly maximal subgroups (see Definition 2.2) were also studied in the context of branch groups, for instance in [4].

2 Preliminaries

In this section we give the basic definitions and claims that will be used in the rest of this paper. We will work in the category of profinite groups, namely we assume that every subgroup is closed, every homomorphism is continuous, and so on. For a finitely generated profinite group G we denote by $d(G)$ the minimal size of a generating set of G .

Definition 2.1. An infinite profinite group G is called *just infinite* if for every $\{1\} \neq N \triangleleft_c G$ the quotient G/N is finite. Equivalently, every nontrivial normal subgroup of G is open. We say that G is *hereditarily just infinite* if every open normal subgroup of it is just infinite.

Definition 2.2. Let H be a closed subgroup of infinite index in a profinite group G . We say that H is *weakly maximal* in G if every $H \leq K \leq_c G$ is open.

Definition 2.3. Given groups G, A, B and surjective homomorphisms

$$\alpha : A \rightarrow B, \quad \beta : G \rightarrow B,$$

we define the *embedding problem* $\mathcal{E}(G, A, B, \alpha, \beta)$ as the problem of finding a homomorphism $\varphi : G \rightarrow A$ such that $\beta = \alpha \circ \varphi$. Such a homomorphism φ is called a *solution* to the problem. If moreover φ is surjective, then it is called a *proper solution*. An embedding problem $\mathcal{E}(G, A, B, \alpha, \beta)$ is called *finite* if A is finite. Note that since α is surjective, B is also finite.

Definition 2.4. An embedding problem $\mathcal{E}(G, A, B, \alpha, \beta)$ is called *split* if there is a homomorphism $\gamma : B \rightarrow A$ such that $\alpha \circ \gamma = \text{id}_B$. In such a case we have $A \cong \text{Ker}(\alpha) \rtimes B$.

Definition 2.5. A profinite group G of rank $\aleph_0$ is called *semi-free* if every finite split embedding problem $\mathcal{E}(G, A, B, \alpha, \beta)$ has a proper solution.

Definition 2.6. The *profinite surface group of genus g* is the group given by the profinite presentation

$$\Gamma_g = \left\langle x_1, \dots, x_g, y_1, \dots, y_g \mid \prod_{i=1}^g [x_i, y_i] = 1 \right\rangle.$$

The set $x_1, \dots, x_g, y_1, \dots, y_g$ from such a presentation will be called a *surface basis*.

Fact 2.7. An open subgroup H of a genus g profinite surface group Γ is a profinite surface group of genus $[\Gamma : H](g - 1) + 1$.

Claim 2.8. Let $\mathcal{E}(G, A, B, \alpha, \beta)$ be an embedding problem and let $\varphi : G \rightarrow A$ be a solution of $\mathcal{E}$. If $\text{Ker}(\alpha) \subseteq \text{Im}(\varphi)$, then φ is a proper solution.

Proof. Let $a \in A$. Denote $b = \alpha(a)$ and let $g \in G$ be such that $\beta(g) = b$. Then

$$\alpha(a) = \beta(g) = \alpha(\varphi(g)),$$

therefore

$$a\varphi(g)^{-1} \in \text{Ker}(\alpha) \subseteq \text{Im}(\varphi).$$

As $\text{Im}(\varphi)$ is a subgroup, we conclude that $a \in \text{Im}(\varphi)$ so $\text{Im}(\varphi) = A$. □

3 Semi-free subgroup

We need the following variant of [5, Lemma 6.1].

Lemma 3.1. *Let*

$$\Gamma = \left\langle x_i, y_i \mid \prod_{i=1}^g [x_i, y_i] = 1 \right\rangle$$

be a profinite surface group, and let $N \leq_c \Gamma$. Consider the diagram

$$\begin{array}{ccc} & \Gamma & \\ & \downarrow \beta & \searrow \bar{\beta} \\ N & & \\ \downarrow \beta & & \\ A & \xrightarrow{\alpha} & B, \end{array}$$

where A, B are finite groups, α is a surjection, and $\bar{\beta}|_N = \beta$ is a surjection as well. Denote by

$$\mathcal{E} = \mathcal{E}(N, A, B, \alpha, \beta), \quad \bar{\mathcal{E}} = \mathcal{E}(\Gamma, A, B, \alpha, \bar{\beta})$$

the two finite embedding problems in the above diagram. Let $\varphi: \Gamma \rightarrow A$ be a solution to $\bar{\mathcal{E}}$, and set

$$K := \text{Ker}(\alpha), \quad n := |K\varphi(\Gamma)|, \quad s := d(K).$$

Suppose that $g \geq sn + s$ and that for all $1 \leq i, j \leq sn + s$,

$$\varphi(x_i) = \varphi(x_j), \quad \varphi(y_i) = \varphi(y_j), \quad x_i x_1^{-1} \in N.$$

Then $\mathcal{E}$ admits a proper solution.

Proof. Choose a set of generators $B := \{k_1, \dots, k_s\}$ of K . Define $\eta(x_1) = \varphi(x_1)$ and

$$\eta(x_{in+j+1}) = \varphi(x_{sn+i+1})k_{i+1} = \varphi(x_1)k_{i+1}, \quad 0 \leq i \leq s-1, \quad 1 \leq j \leq n.$$

Let η coincide with φ for all other generators of Γ , that is,

$$\eta(x_i) = \varphi(x_i), \quad \eta(y_j) = \varphi(y_j), \quad sn+1 < i \leq g, \quad 1 \leq j \leq g.$$

Since

$$[\varphi(x_{sn+i+1})k_{i+1}, \varphi(y_{sn+i+1})] \in K\varphi(\Gamma), \quad 0 \leq i \leq s-1,$$

we get

$$[\eta(x_{in+2}), \eta(y_{in+2})]^n = 1 = [\varphi(x_{in+2}), \varphi(y_{in+2})]^n, \quad 0 \leq i \leq s-1.$$

Therefore

$$\begin{aligned}
 & \prod_{i=1}^g [\eta(x_i), \eta(y_i)] \\
 &= [\eta(x_1), \eta(y_1)] \cdot \prod_{i=0}^{s-1} [\eta(x_{in+2}), \eta(y_{in+2})]^n \cdot \prod_{i=sn+2}^g [\eta(x_i), \eta(y_i)] \\
 &= [\varphi(x_1), \varphi(y_1)] \cdot \prod_{i=0}^{s-1} [\varphi(x_{in+2}), \varphi(y_{in+2})]^n \cdot \prod_{i=sn+2}^g [\varphi(x_i), \varphi(y_i)] \\
 &= \prod_{i=1}^g [\varphi(x_i), \varphi(y_i)] = 1.
 \end{aligned}$$

Thus η extends to a homomorphism. Since $x_1^{-1}x_i \in N$ for $1 \leq i \leq sn + s$, we conclude that $k_{j+1} = \eta(x_1^{-1}x_{jn+2}) \in \eta(N)$ for $0 \leq j \leq s-1$, hence $\eta(N)$ contains K , so the result follows by invoking Claim 2.8. $\square$

We also need the following generalization of [2, Lemma 2.2].

Lemma 3.2. *Let Γ_g be a profinite surface group of genus g , let*

$$\mathcal{E} = \mathcal{E}(\Gamma_g, A, B, \alpha, \beta)$$

be a finite split embedding problem, and suppose that $g \geq 2|A|^2|B|$. Then $\mathcal{E}$ has a proper solution, and Γ_g has a surface basis $x_1, \dots, x_g, y_1, \dots, y_g$ such that

$$x_i x_1^{-1}, y_i y_1^{-1} \in \text{Ker}(\beta), \quad 1 \leq i \leq m, \quad m = \frac{2|A|^2}{|B|}.$$

Furthermore, if $N \leq_c \Gamma_g$ is such that $\beta(N) = B$ and

$$x_i x_1^{-1} \in N, \quad 1 \leq i \leq m,$$

then the embedding problem $\bar{\mathcal{E}} = \mathcal{E}(N, A, B, \alpha, \beta|_N)$ is properly solvable.

Proof. Let $z_1, \dots, z_g, w_1, \dots, w_g$ be a surface basis of Γ_g . Let γ be a section of α and set $\varphi = \gamma \circ \beta$. As before, put $K = \text{Ker}(\alpha)$, $s = d(K)$, and $n = |K\varphi(\Gamma_g)| = |A|$. Note that

$$s \leq |K| = \frac{|A|}{|B|}$$

and thus

$$sn + s \leq \frac{|A|^2 + |A|}{|B|} \leq \frac{2|A|^2}{|B|} = m.$$

Each of the pairs $(\varphi(z_i), \varphi(w_i))$ can attain at most $|B|^2$ values, whence by the pigeonhole principle (since $g \geq m|B|^2$) there are

$$1 \leq j_1 < j_2 < \cdots < j_m \leq g$$

such that

$$\varphi(z_{j_1}) = \cdots = \varphi(z_{j_m}), \quad \varphi(w_{j_1}) = \cdots = \varphi(w_{j_m}).$$

Suppose $j_1 \neq 1$. Then

$$\begin{aligned} 1 &= \prod_{i=1}^g [z_i, w_i] \\ &= [z_{j_1}, w_{j_1}] \left[\prod_{i=1}^{j_1-1} [z_i, w_i] \right]^{[z_{j_1}, w_{j_1}]} \prod_{i=j_1+1}^g [z_i, w_i] \\ &= [z_{j_1}, w_{j_1}] \cdot \prod_{i=1}^{j_1-1} [z_i^{[z_{j_1}, w_{j_1}]}, w_i^{[z_{j_1}, w_{j_1}]}] \cdot \prod_{i=j_1+1}^g [z_i, w_i] \end{aligned}$$

and so we can replace $\{z_i, w_i\}_{i=1}^g$ with a new surface basis

$$\begin{aligned} &z_{j_1}, z_1^{[z_{j_1}, w_{j_1}]}, \dots, z_{j_1-1}^{[z_{j_1}, w_{j_1}]}, z_{j_1+1}, \dots, z_g, \\ &w_{j_1}, w_1^{[z_{j_1}, w_{j_1}]}, \dots, w_{j_1-1}^{[z_{j_1}, w_{j_1}]}, w_{j_1+1}, \dots, w_g. \end{aligned}$$

By repeating this process with $j_2, \dots, j_m$, we obtain a surface basis $x_1, \dots, x_g, y_1, \dots, y_g$ of Γ_g such that $x_i = z_{j_i}, y_i = w_{j_i}$ for $1 \leq i \leq m$, and so

$$\varphi(x_1) = \cdots = \varphi(x_m), \quad \varphi(y_1) = \cdots = \varphi(y_m).$$

Since $sn + s \leq m$, we can apply Lemma 3.1 (with $N = \Gamma_g$ if necessary) and the result follows. $\square$

Corollary 3.3. *The finite split embedding problem*

$$\begin{array}{ccccccc} & & & & \Gamma_g & \amalg & G \\ & & & & \downarrow \beta & & \\ 1 & \longrightarrow & K & \longrightarrow & K \rtimes H & \longrightarrow & H \longrightarrow 1 \end{array}$$

has a proper solution once $g \geq 2|K|^2|H|^3$, Γ_g is a profinite surface group of genus g , and G is any profinite group.

Proof. Write H_0 for $\beta(\Gamma_g)$. According to Lemma 3.2, the finite split embedding problem

$$\begin{array}{ccccccc} & & & & \Gamma_g & & \\ & & & & \downarrow \beta|_{\Gamma_g} & & \\ 1 & \longrightarrow & K & \longrightarrow & K \rtimes H_0 & \longrightarrow & H_0 \longrightarrow 1 \end{array}$$

has a proper solution φ_1 . Let $\varphi_2 : G \rightarrow K \rtimes H$ be the map defined by

$$\varphi_2(f) = (1, \beta|_G(f))$$

for every $f \in G$. There exists a unique homomorphism

$$\varphi : \Gamma_g \amalg G \rightarrow K \rtimes H$$

such that

$$\varphi(\gamma) = \varphi_1(\gamma), \quad \varphi(f) = \varphi_2(f) \quad \text{for all } \gamma \in \Gamma_g \text{ and all } f \in G.$$

By the universal property of free products, φ is a solution to the original embedding problem. Since K is contained in $\text{Im}(\varphi_1)$, it is also contained in $\text{Im}(\varphi)$ so by Claim 2.8, φ is a proper solution. $\square$

Theorem 3.4. *Let $N \triangleleft_c \Gamma_g$ be a normal subgroup of the profinite surface group of genus $g \geq 2$ such that Γ_g/N is hereditarily just infinite. Then N is semi-free.*

Proof. Let

$$\begin{array}{ccccccc} & & & & N & & \\ & & & & \downarrow \beta & & \\ 1 & \longrightarrow & K & \longrightarrow & K \rtimes H & \longrightarrow & H \longrightarrow 1 \end{array}$$

be a finite split embedding problem for N . We shall prove it has a proper solution.

Using [3, Lemma 1.2.5 (c)], we can extend our embedding problem to a subgroup

$$N \leq F \triangleleft_o \Gamma_g$$

such that

$$[\Gamma_g : F] \geq \frac{2|K|^2|H|^3 + m - 1}{g - 1},$$

where $m = 2|K|^2|H|$. By Fact 2.7, F is a profinite surface group of genus

$$h = [\Gamma_g : F](g - 1) + 1.$$

Note that

$$h - m \geq 2|K|^2|H|^3.$$

Applying Lemma 3.2 to F , the extended embedding problem and m , we obtain a proper solution φ , and a surface basis $x_1, \dots, x_h, y_1, \dots, y_h$ of F such that

$$x_j x_1^{-1}, y_j y_1^{-1} \in \text{Ker}(\beta), \quad 1 \leq j \leq m.$$

If $x_i x_1^{-1} \in N$ for all $1 \leq i \leq m$, then by Lemma 3.2, our original embedding problem has a proper solution. Assume henceforth that $x_i x_1^{-1} \notin N$ for some i with $1 \leq i \leq m$.

The homomorphism β factors modulo

$$L = \langle x_2 x_1^{-1}, \dots, x_m x_1^{-1}, y_2 y_1^{-1}, \dots, y_m y_1^{-1} \rangle^F.$$

Note that

$$F/L \cong \left\langle x_1, x_{m+1}, \dots, x_h, y_1, y_{m+1}, \dots, y_h \mid [x_1, y_1]^m \cdot \prod_{i=m+1}^h [x_i, y_i] = 1 \right\rangle$$

and that

$$\beta([x_1, y_1]^m) = \beta([x_1, y_1])^m = 1,$$

so β even factors through

$$\begin{aligned} F/\langle L, [x_1, y_1]^m \rangle^F &\cong \left\langle x_i, y_i \mid \prod_{i=m+1}^h [x_i, y_i] = 1 \right\rangle \amalg \langle x_1, y_1 \mid [x_1, y_1]^m = 1 \rangle \\ &\cong \Gamma_{h-m} \amalg G, \end{aligned}$$

where $G = \langle x_1, y_1 \mid [x_1, y_1]^m = 1 \rangle$.

Let

$$\psi: F \rightarrow F/\langle L, [x_1, y_1]^m \rangle^F$$

be the quotient map, and set $M = \psi(N)$. Since Γ_g/N is hereditarily just infinite, F/N is just infinite, so $\psi^{-1}(M)$ either equals N or is open in F . The former is impossible as there exists $1 \leq i \leq m$ such that

$$x_i x_1^{-1} \in \text{Ker}(\psi) \subseteq \psi^{-1}(M), \quad x_i x_1^{-1} \notin N.$$

Hence $\psi^{-1}(M)$ is an open subgroup of F , so M can be seen as an open subgroup of $\Gamma_{h-m} \amalg G$.

We can now write the original embedding problem as

$$\begin{array}{c}
 N \\
 \downarrow \psi|_N \\
 M \\
 \downarrow \bar{\beta} \\
 1 \longrightarrow K \longrightarrow K \rtimes H \longrightarrow H \longrightarrow 1,
 \end{array}$$

so it is sufficient to properly solve it for M . Applying the Kurosh theorem (see [6, Theorem D.3.1]), we find that $M \cong \Gamma_t \sqcup \overline{G}$, where

$$t \geq h - m \geq 2|K|^2|H|^3$$

and so by Corollary 3.3 the desired proper solution exists. $\square$

Repeating the above proof verbatim, one obtains the following.

Theorem 3.5. *Let Γ_g be a surface group and $N \leq_c \Gamma_g$ weakly maximal. Then N is semi-free profinite.*

Acknowledgments. The authors would like to thank Lior Bary-Soroker and Pavel Zalesskii for useful remarks and discussions. Mark Shusterman is grateful to the Azrieli Foundation for the award of an Azrieli Fellowship.

Bibliography

- [1] L. Bary-Soroker, D. Haran and D. Harbater, Permanence criteria for semi-free profinite groups, *Math. Ann.* **348** (2010), no. 3, 539–563.
- [2] L. Bary-Soroker, K. F. Stevenson and P. A. Zalesskii, Subgroups of profinite surface groups, *Math. Res. Lett.* **18** (2011), no. 3, 459–471.
- [3] M. D. Fried and M. Jarden, *Field Arithmetic*, 3rd ed., *Ergeb. Math. Grenzgeb.* (3) 11, Springer, Berlin, 2008.
- [4] R. I. Grigorchuk, Branch groups, *Mat. Zametki* **67** (2000), no. 6, 852–858.
- [5] A. Pacheco, K. F. Stevenson and P. Zalesskii, Normal subgroups of the algebraic fundamental group of affine curves in positive characteristic, *Math. Ann.* **343** (2009), no. 2, 463–486.
- [6] L. Ribes and P. Zalesskii, *Profinite Groups*, 2nd ed., *Ergeb. Math. Grenzgeb.* (3) 40, Springer, Berlin, 2010.

- [7] M. Shusterman, Free subgroups of finitely generated free profinite groups, *J. Lond. Math. Soc. (2)* **93** (2016), no. 2, 361–378.
- [8] P. A. Zalesskii, Profinite surface groups and the congruence kernel of arithmetic lattices in $\mathrm{SL}_2(\mathbb{R})$, *Israel J. Math.* **146** (2005), 111–123.

Received April 15, 2018; revised May 7, 2018.

Author information

Matan Ginzburg, Raymond and Beverly Sackler School of Mathematical Sciences,
Tel-Aviv University, Tel-Aviv, Israel.
E-mail: matan.ginzburg@gmail.com

Mark Shusterman, Raymond and Beverly Sackler School of Mathematical Sciences,
Tel-Aviv University, Tel-Aviv, Israel.
E-mail: markshus@mail.tau.ac.il