Mohammed Baashen*, Bakr F. Abdel-Wahab, Amany S. Hegazy, Benson M. Kariuki and Gamal A. El-Hiti

Crystal structure of 1-phenyl-*N'*-(1-phenyl-5-(thiophen-2-yl)-1*H*-pyrazole-3-carbonyl)-5-(thiophen-2-yl)-1*H*-pyrazole-3-carbohydrazide,

$C_{28}H_{20}N_6O_2S_2$

<https://doi.org/10.1515/ncrs-2017-0375>

Received February 26, 2018; accepted May 14, 2018; available online May 30, 2018

Abstract

$C_{28}H_{20}N_6O_2S_2$, triclinic, $P\bar{1}$ (no. 2), $a = 10.6738(6)$ Å, $b = 11.7869(7)$ Å, $c = 12.5381(7)$ Å, $\alpha = 112.842(6)^\circ$, $\beta = 91.963(4)^\circ$, $\gamma = 116.129(6)^\circ$, $V = 1264.38(15)$ Å³, $Z = 2$, $R_{gt}(F) = 0.0523$, $wR_{ref}(F^2) = 0.1390$, $T = 296(2)$ K.

CCDC no.: 1843111

Tables 1 and 2 contain details on crystal structure and measurement conditions and a list of the atoms including atomic coordinates and displacement parameters.

Source of material

The title compound was synthesized from reaction of an equimolar mixture of 1-phenyl-5-(thiophen-2-yl)-1*H*-pyrazole-3-carbohydrazide and 2-(methoxymethylene)malononitrile in ethanol under reflux for 1.5 h. The solid obtained on cooling was collected by filtration, dried and recrystallized

*Corresponding author: Mohammed Baashen, Department of Chemistry, College of Science and Humanities, Shaqra University, Duwadimi, Saudi Arabia, e-mail: mbaashen@su.edu.sa

Bakr F. Abdel-Wahab: Department of Chemistry, College of Science and Humanities, Shaqra University, Duwadimi, Saudi Arabia and Applied Organic Chemistry Department, National Research Centre, Dokki, Giza, Egypt

Amany S. Hegazy and Benson M. Kariuki: School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, UK

Gamal A. El-Hiti: Cornea Research Chair, Department of Optometry, College of Applied Medical Sciences, King Saud University, P.O. Box 10219, Riyadh 11433, Saudi Arabia

Table 1: Data collection and handling.

Crystal:	Colourless needle
Size:	0.30 × 0.19 × 0.15 mm
Wavelength:	Mo K α radiation (0.71073 Å)
μ :	0.25 mm ^{−1}
Diffractometer, scan mode:	SuperNova, ω
θ_{max} , completeness:	29.7°, >99%
$N(hkl)_{measured}$, $N(hkl)_{unique}$, R_{int} :	10719, 6002, 0.019
Criterion for I_{obs} , $N(hkl)_{gt}$:	$I_{obs} > 2 \sigma(I_{obs})$, 4380
$N(param)_{refined}$:	417
Programs:	CrysAlis ^{PRO} [9], SHELX [10], WinGX [11]

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	x	y	z	U_{iso}^*/U_{eq}
S1 ^a	0.53465(11)	0.34985(13)	0.71518(12)	0.0664(4)
C1 ^a	0.4846(5)	0.4560(5)	0.8228(4)	0.0627(12)
H1 ^a	0.549379	0.537074	0.891002	0.075*
C2 ^a	0.3408(5)	0.4085(6)	0.7972(5)	0.0651(11)
H2 ^a	0.292964	0.450287	0.843816	0.078*
C3 ^a	0.2758(8)	0.2832(11)	0.6862(9)	0.0656(16)
H3 ^a	0.177161	0.234682	0.652859	0.079*
C4 ^a	0.3610(2)	0.2385(2)	0.63239(17)	0.0466(5)
S1A ^b	0.2424(7)	0.2796(10)	0.6921(9)	0.0688(14)
C1A ^b	0.3838(13)	0.4266(18)	0.8129(15)	0.057(3)
H1A ^b	0.369843	0.491069	0.875773	0.069*
C2A ^b	0.5156(14)	0.4382(16)	0.8067(13)	0.059(3)
H2A ^b	0.602666	0.507921	0.862585	0.070*
C3A ^b	0.4940(12)	0.3221(17)	0.6974(14)	0.067(3)
H3A ^b	0.569287	0.307032	0.674116	0.081*
C4A ^b	0.3610(2)	0.2385(2)	0.63239(17)	0.0466(5)
C5	0.3135(2)	0.10572(19)	0.52627(17)	0.0421(4)
C6	0.2007(2)	−0.0253(2)	0.49954(18)	0.0473(5)
H6	0.134030	−0.048065	0.544076	0.057*
C7	0.2068(2)	−0.11664(19)	0.39219(18)	0.0447(4)
C8	0.5003(2)	0.19227(19)	0.41823(16)	0.0425(4)
C9	0.6187(3)	0.1766(3)	0.4017(2)	0.0677(7)
H9	0.621872	0.099208	0.402195	0.081*
C10	0.7305(3)	0.2750(4)	0.3846(3)	0.0898(9)
H10	0.811686	0.265428	0.373942	0.108*
C11	0.7275(3)	0.3892(3)	0.3826(2)	0.0809(9)
H11	0.806201	0.456831	0.371479	0.097*
C12	0.6079(3)	0.4026(2)	0.3970(2)	0.0694(7)

Table 2 (continued)

Atom	x	y	z	U_{iso}^*/U_{eq}
H12	0.604777	0.479025	0.394240	0.083*
C13	0.4937(2)	0.3062(2)	0.41518(18)	0.0508(5)
H13	0.412489	0.315899	0.425448	0.061*
C14	0.1114(2)	−0.2710(2)	0.32795(19)	0.0482(5)
C15	0.0942(2)	−0.5481(2)	0.06679(19)	0.0493(5)
C16	0.0018(2)	−0.70304(19)	0.00660(18)	0.0452(5)
C17	0.0174(2)	−0.7993(2)	−0.09108(18)	0.0478(5)
H17	0.085527	−0.779631	−0.135102	0.057*
C18	−0.0892(2)	−0.93014(19)	−0.10910(16)	0.0421(4)
S2 ^c	−0.1851(3)	−1.21667(18)	−0.1788(2)	0.0564(5)
C19 ^c	−0.1175(2)	−1.0688(2)	−0.19663(17)	0.0440(4)
C20 ^c	−0.0800(17)	−1.0914(14)	−0.3047(9)	0.057(2)
H20 ^c	−0.042354	−1.022008	−0.330743	0.068*
C2 ^c	−0.1048(12)	−1.2311(7)	−0.3716(8)	0.0511(13)
H21 ^c	−0.087427	−1.264754	−0.446696	0.061*
C22 ^c	−0.1571(10)	−1.3091(8)	−0.3113(5)	0.0500(13)
H22 ^c	−0.175656	−1.401789	−0.338654	0.060*
S2A ^d	−0.0696(7)	−1.0881(6)	−0.3229(4)	0.0560(8)
C19A ^d	−0.1175(2)	−1.0688(2)	−0.19663(17)	0.0440(4)
C20A ^d	−0.1586(18)	−1.1886(11)	−0.1788(14)	0.059(3)
H20A ^d	−0.179414	−1.193369	−0.108754	0.071*
C21A ^d	−0.1649(17)	−1.3030(13)	−0.2809(9)	0.056(2)
H21A ^d	−0.200408	−1.394590	−0.289816	0.067*
C22A ^d	−0.1120(19)	−1.2596(10)	−0.3630(12)	0.054(2)
H22A ^d	−0.100014	−1.316450	−0.433204	0.064*
C23	−0.2811(2)	−1.00608(19)	−0.00114(17)	0.0431(4)
C24	−0.4113(2)	−1.0925(2)	−0.0840(2)	0.0565(6)
H24	−0.423716	−1.086218	−0.154914	0.068*
C25	−0.5238(3)	−1.1892(2)	−0.0603(2)	0.0696(7)
H25	−0.613023	−1.247945	−0.115284	0.083*
C26	−0.5048(3)	−1.1988(3)	0.0434(3)	0.0768(8)
H26	−0.580538	−1.264818	0.058451	0.092*
C27	−0.3737(3)	−1.1112(3)	0.1256(3)	0.0794(8)
H27	−0.361020	−1.118217	0.196119	0.095*
C28	−0.2615(3)	−1.0134(3)	0.1042(2)	0.0625(6)
H28	−0.173220	−0.952913	0.160305	0.075*
N1	0.38018(17)	0.08720(15)	0.43565(14)	0.0425(4)
N2	0.31608(18)	−0.05003(16)	0.35200(15)	0.0463(4)
N3	0.14238(19)	−0.33627(16)	0.22880(16)	0.0553(5)
H3B	0.211634	−0.288734	0.204268	0.066*
N4	0.0618(2)	−0.48181(17)	0.16540(17)	0.0599(5)
H4	−0.008465	−0.529077	0.189308	0.072*
N5	−0.10724(18)	−0.76601(16)	0.04906(15)	0.0467(4)
N6	−0.16261(17)	−0.90569(15)	−0.02352(14)	0.0435(4)
O1	0.01181(19)	−0.33486(16)	0.36309(17)	0.0776(6)
O2	0.1941(2)	−0.48422(16)	0.03243(17)	0.0810(6)

Occupancies: ^a = 0.768(3); ^b = 0.232(3); ^c = 0.609(3); ^d = 0.391(3).

from dimethylformamide to colourless crystals (Mp. > 300 °C; 67%).

Experimental details

The two thiophenyl groups are disordered and the occupancies refined to 0.768(3)/0.232(3) and 0.609(3)/0.391(3). Both components of either disordered group were refined with

similar geometry and displacement parameters (SAME, SIMU and ISOR instructions in SHELXL [10]). The hydrogen atoms were placed in calculated positions (AFIX 43 instruction in SHELXL [10]) with the U_{iso} values set to 1.2 U_{eq} (C, N).

Comment

N,N'-Diacylhydrazines have been synthesized efficiently using various synthetic approaches as biologically active compounds [1–4]. Also, they show fungicidal and herbicidal activities [5–7]. The X-ray crystal structure for a similar compound has been published [8].

The asymmetric unit comprises one molecule in which both thiophenyl groups are disordered. The two thiophenyl groups are twisted from the least-squares plane of the pyrazole-carbohydrazone group by 45.3(3)° and 31.1(2)° whereas the two phenyl groups are twisted by 53.1(8)° and 59.8(1)°. The pyrazole-carbohydrazone groups of the molecules form planes parallel to (111) in the crystal. The molecules are arranged in pairs which are linked by edge-to-face contacts between phenyl groups with centroid-to-centroid distances of 4.88 Å. The separation between the planes of the pyrazole-carbohydrazone groups of adjacent molecules is 3.46 Å.

Acknowledgements: The project was supported by King Saud University, Deanship of Scientific Research, Research Chairs.

References

1. Zhen, M.; Peng, Y.: *N,N*-Dicarboxymethyl hydrazine: an old but neglected reagent for chemoselective derivatization of carbonyl compounds. *Org. Biomol. Chem.* **14** (2016) 3443–3449.
2. Sun, G.-X.; Sun, Z.-H.; Yang, M.-Y.; Liu, X.-H.; Ma, Y.; Wei, Y.-Y.: Design, synthesis, biological activities and 3D-QSAR of new *N,N'*-diacylhydrazines containing 2,4-dichlorophenoxy moieties. *Molecules* **18** (2013) 14876–14891.
3. Liu, X. H.; Shi, Y. X.; Ma, Y.; He, G. R.; Dong, W. L.; Zhang, C. Y.; Wang, B. L.; Wang, S. Cui, Z.; Zhang, L.; Huang, J.; Yang, X.; Ling, Y.: Synthesis and bioactivity of novel *N,N'*-diacylhydrazine derivatives containing furan (III). *Chin. J. Chem.* **28** (2010) 1257–1266.
4. Kantelehner, W.; Mezger, J.; Stoyanov, E. V.; Leonhardt, S.: A new synthesis of *N,N'*-bis(dimethylaminomethylene) hydrazine. *Synthesis* **21** (2008) 3380–3382.
5. Zhao, Q.; Shang, J.; Liu, Y.; Wang, K.; Bi, F.; Hauang, R.; Wang, Q.: Synthesis and insecticidal activities of novel *N*-sulfenyl-*N'*-tert-butyl-*N,N'*-diacylhydrazines. *J. Agric. Food Chem.* **55** (2007) 9614–9619.
6. Zhang, X. Y.; Li, Y. S.; Weng, J. Q.; Tan, C. X. Synthesis and herbicidal activity of some new diacylhydrazine derivatives fluoride-containing pyrazolyl moiety. *Chin. J. Org. Chem.* **31** (2011) 1295–1299.
7. Liu, X. H.; Li, B. J.; Li, Z. M.: Synthesis of some *N,N'*-diacylhydrazine derivatives with radical-scavenging

- and antifungal activity. *Chem. Bio. Drug. Des.* **73** (2009) 320–327.
8. Klapötke, T. M.; Krumm, B.; Scharf, R.: *N*-Nitrosarcosine: an economic precursor for the synthesis of new energetic materials. *Chem. Asian J.* **11** (2016) 3134–3144.
9. CrysAlis^{PRO}. Agilent Technologies, Yarnton, England (2014).
10. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr. A* **64** (2008) 112–122.
11. Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Crystallogr.* **45** (2012) 849–854.