Musa A. Said*, Mohamed R. Aouad, Saud M. Almutairi, David L. Hughes*
and Mouslim Messali*

Crystal structure of 1-heptylpyridazin-1-ium iodide, $C_{11}H_{19}N_2I$

Table 1: Data collection and handling.

Crystal:	Light yellow prism
Size:	$0.35 \times 0.11 \times 0.10$ mm
Wavelength:	Mo $K\alpha$ radiation ($0.71073 \text{ \AA}$)
μ :	2.21 mm^{-1}
Diffractometer, scan mode:	Xcalibur 3/Sapphire3, Thin slice φ and ω -scans
$\theta_{\max}$, completeness:	27.5° , $>99\%$
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	11282, 3285, 0.041
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 2794
$N(\text{param})_{\text{refined}}$:	127
Programs:	CrysAlis [1], SHELX [2, 3], WinGX [4]

<https://doi.org/10.1515/ncrs-2018-0090>

Received March 1, 2018; accepted April 12, 2018; available online June 9, 2018

Abstract

$C_{11}H_{19}N_2I$, triclinic, $P\bar{1}$ (no. 2), $a = 5.7074(3) \text{ \AA}$, $b = 8.9811(5) \text{ \AA}$, $c = 14.4312(8) \text{ \AA}$, $\alpha = 100.989(4)^\circ$, $\beta = 94.768(4)^\circ$, $\gamma = 97.327(4)^\circ$, $V = 715.83(7) \text{ \AA}^3$, $Z = 2$, $R_{\text{gt}}(F) = 0.0386$, $wR_{\text{ref}}(F^2) = 0.0913$, $T = 295(2) \text{ K}$.

CCDC no.: 1836644

The asymmetric unit of the title crystal structure is shown in the figure. Tables 1 and 2 contain details on crystal structure and measurement conditions and a list of the atoms including atomic coordinates and displacement parameters.

Source of materials

The title ionic liquid was prepared according to a method reported earlier [5, 6]. To a solution of pyridazine (1 g, 12.5 mmol in 10 mL of toluene) was added dropwise

1-iodoheptane (2.825 g, 12.5 mmol) and the mixture was placed in a closed container and exposed to irradiation for 5 hours at room temperature using a sonication bath. Completion of the reaction was marked by the precipitation of a solid from the initially obtained clear and homogenous mixture in toluene. The pyridazin-1-ium-based ionic liquid is isolated by filtration and washed three times with ethyl acetate to remove any unreacted starting materials and solvent. Finally the 1-heptylpyridazin-1-ium iodide was dried at a reduced pressure to remove all volatile organic compounds to produce a yellow powder. (Yield 77%, m.p. $89\text{--}92^\circ\text{C}$). Crystals were obtained from a mixture of dichloromethane and *n*-hexane (1:2). Elemental analysis: Anal. Calc. for $C_{11}H_{19}IN_2$: C, 43.15%; H, 6.25%; N, 9.15%; Found: C, 43.10%; H, 6.19%; N, 9.11%. $^1\text{H NMR}$ (DMSO, 400 MHz): $\delta = 0.81$ (t, 3H), 1.23–1.30 (m, 8H), 1.99 (quintet, 2H), 4.83 (t, 2H), 7.65 (t, 1H), 8.77 (t, 1H), 9.66 (d, 1H), 10.02 (d, 1H); $^{13}\text{C NMR}$ (DMSO, 100 MHz): $\delta = 14.4$ (CH_3), 22.6 (CH_2), 25.8 (CH_2), 28.8 (CH_2), 29.9 (CH_2), 31.7 (CH_2), 65.0 (CH_2), 136.5 (CH), 137.0 (CH), 150.3 (CH), 155.0 (CH).

Experimental details

The diffraction data were processed using the CrysAlisPro-CCD and -RED [1] programs. The structure was determined by the intrinsic phasing routines in the SHELXT program [2] and refined by full-matrix least-squares methods, on F^2 's, in SHELXL [3]. Hydrogen atoms were included in idealized positions and their U_{iso} values were set to ride on the U_{eq} values of the parent carbon atoms.

*Corresponding authors: Musa A. Said and Mouslim Messali, Chemistry Department, Taibah University, P.O. Box 30002, Code 14177, Al-Madinah Al-Munawarah, Kingdom of Saudi Arabia, e-mail: musa_said04@yahoo.co.uk (M.A. Said); mouslim@mail.be (M. Messali); and David L. Hughes, School of Chemistry, University of East Anglia, Norwich, NR4 7TJ, United Kingdom, e-mail: d.l.hughes@uea.ac.uk

Mohamed R. Aouad: Chemistry Department, Taibah University, P.O. Box 30002, Code 14177, Al-Madinah Al-Munawarah, Kingdom of Saudi Arabia

Saud M. Almutairi: King Abdulaziz City for Science and Technology, Riyadh 11442, P.O. Box 6086, Saudi Arabia

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	x	y	z	<i>U</i> _{iso} */ <i>U</i> _{eq}
I	0.90826(4)	0.75400(3)	0.58391(2)	0.06733(12)
N1	0.5799(5)	0.6420(3)	0.34437(19)	0.0579(7)
N2	0.7257(5)	0.7389(4)	0.3100(2)	0.0685(8)
C3	0.6800(7)	0.8800(4)	0.3215(3)	0.0741(10)
H3	0.7812	0.9504	0.2984	0.089*
C4	0.4921(8)	0.9308(5)	0.3657(3)	0.0754(11)
H4	0.4659	1.0322	0.3718	0.090*
C5	0.3474(7)	0.8281(6)	0.3997(3)	0.0766(11)
H5	0.2169	0.8566	0.4294	0.092*
C6	0.3969(6)	0.6810(5)	0.3895(2)	0.0676(9)
H6	0.3026	0.6089	0.4141	0.081*
C11	0.6434(8)	0.4853(5)	0.3325(3)	0.0798(11)
H11A	0.8012	0.4903	0.3648	0.096*
H11B	0.5328	0.4234	0.3624	0.096*
C12	0.6393(10)	0.4109(6)	0.2323(4)	0.0920(14)
H12A	0.6896	0.3110	0.2289	0.110*
H12B	0.7520	0.4720	0.2027	0.110*
C13	0.3966(9)	0.3914(6)	0.1775(3)	0.1012(15)
H13A	0.3547	0.4920	0.1750	0.121*
H13B	0.2812	0.3412	0.2116	0.121*
C14	0.3793(10)	0.2984(8)	0.0763(4)	0.1124(17)
H14A	0.4929	0.3495	0.0419	0.135*
H14B	0.4239	0.1984	0.0787	0.135*
C15	0.1357(11)	0.2765(9)	0.0219(4)	0.126(2)
H15A	0.0218	0.2274	0.0570	0.151*
H15B	0.0927	0.3765	0.0185	0.151*
C16	0.1157(14)	0.1844(13)	−0.0753(5)	0.160(3)
H16A	0.1625	0.0851	−0.0725	0.192*
H16B	0.2248	0.2350	−0.1116	0.192*
C17	−0.1308(14)	0.1613(12)	−0.1255(5)	0.184(4)
H17A	−0.1341	0.1008	−0.1883	0.275*
H17B	−0.2392	0.1092	−0.0907	0.275*
H17C	−0.1769	0.2591	−0.1298	0.275*

Comment

Ionic liquids (ILs) have received increased attention in recent years due to their outstanding and unique properties, such as negligible vapor pressure, non-volatility, non-flammability, excellent thermal stability, and high electrical conductivity [7]. Based on these characteristics, a broad range of applications of ILs has been reported in a variety of different fields, including the electrodeposition of metals and the development of corrosion inhibitors [8–10]. Furthermore, a number of biological activities of pyridazine derivatives have been reported including anticancer, antituberculosis, antihypertensive and antimicrobial activities [11–13].

The asymmetric unit of the title crystal structure contains one cation and the iodide anion. Bond lengths and angles in the organic cation are in the expected ranges. The heptyl group has an all-*trans* arrangement and is aligned about C(11)

Table 3: ‘Weak’ hydrogen bonds, in Ångstroms and degrees.

D–H...A	d(D–H)	d(H...A)	d(D...A)	<(DHA)
C(3)–H(3)...I#1	0.93	3.11	3.739(4)	126.8
C(4)–H(4)...I#2	0.93	3.08	3.861(4)	143.2
C(5)–H(5)...I#3	0.93	3.15	3.892(4)	137.8
C(11)–H(11A)...I#4	0.97	3.08	3.808(4)	133.4
C(11)–H(11B)...I#5	0.97	3.04	3.985(5)	164.8

Symmetry transformations used to generate equivalent atoms:

#1 : 2–x, 2–y, 1–z #2 : 1–x, 2–y, 1–z #3 : x–1, y, z
#4 : 2–x, 1–y, 1–z #5 : 1–x, 1–y, 1–z.

with a *cis* N(2)–N(1)–C(11)–C(12) torsion angle of −61.6(5)°; the N(2)–N(1)–C(11)–H(11b) angle is *trans* at 177.2°.

The iodide ion lies over the pyridinium ring at 3.686 Å from N(1). There are also five short H...I contacts in the range 3.04–3.15 Å, to neighboring cations, forming ‘weak’ C–H...I hydrogen bonds, Table 3, which link ions in planes parallel to the *ab* plane, at *z* ∼ 1/2. All the short inter-ion distances involve the iodide ion; the heptyl chains lie parallel but do not show any close contacts between chains. In a recent work from our group, we found a bromide anion linked to a pyridinium cation by a C–H...Br hydrogen bond with the H...Br distance 2.89 Å; there were four further short C–H...Br contacts, to three separate cations, at distances ranging from *ca* 3.07 to 3.11 Å [14].

References

1. Program CrysAlis PRO. Oxford Diffraction Ltd., Yarnton, Oxford, UK (2011).
2. Sheldrick, G. M.: SHELXT – Integrated space-group and crystal-structure determination. *Acta Crystallogr.* **A71** (2015) 3–8.
3. Sheldrick, G. M.: Crystal structure refinement with SHELXL. *Acta Crystallogr.* **C71** (2015) 3–8.
4. Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Cryst.* **45** (2012) 849–854.
5. Messali, M.: An efficient and green sonochemical synthesis of some new eco-friendly functionalized ionic liquids. *Arab. J. Chem.* **7** (2014) 63–70.
6. Messali, M.: Eco-friendly synthesis of a new class of pyridinium-based ionic liquids with attractive antimicrobial activity. *Molecules* **20** (2015) 14936–14949.
7. Earle, M. J.; Esperança, J. M. S. S.; Gilea, M. A.; Lopes, J. N. C.; Rebelo, L. P. N.; Magee, J. W.; Seddon, K. R.; Widegren, J. A.: The distillation and volatility of ionic liquids. *Nature* **439** (2006) 831–834.
8. Ibrahim, M. A. M.; Messali, M.: Ionic liquid [BMPy]Br as an effective additives during zinc electrodeposition from aqueous sulphate bath. *Prod. Finish.* **76** (2011) 14.
9. Lgaz, H.; Benali, O.; Salghi, R.; Jodeh, S.; Larouj, M.; Hamed, O.; Messali, M.; Samhan, S.; Zougagh, M.; Oudda, H.: Pyridinium derivatives as corrosion inhibitors for mild steel in 1M HCl: electrochemical, surface and quantum chemical studies. *Der Pharma Chem.* **8** (2016) 172–190.

10. Al-Ghamdi, A. F.; Messali, M.; Ahmed, S. A.: Electrochemical studies of new pyridazinium-based ionic liquid and its determination in different detergents. *J. Mater. Environ. Sci.* **2** (2011) 215–224.
11. Rodríguez-Ciría, M.; Sanz, A. M.; Yunta, M. J. R.; Gomez-Contreras, F.; Navarro, P.; Fernandez, I.; Pardo, M.; Cano, C.: Synthesis and cytotoxic activity of *N,N*-bis-{3-[*N*-(4-chlorobenzo[*g*]-phthalazin-1-yl)]aminopropyl}-*N*-methylamine: a new potential DNA bisintercalator. *Bioorg. Med. Chem.* **11** (2003) 2143–2148.
12. Demirayak, S.; Karaburun, A. C.; Beis, R.: Some pyrrole substituted aryl pyridazinone and phthalazinone derivatives and their antihypertensive activities. *Eur. J. Med. Chem.* **39** (2004) 1089–1095.
13. Caprosu, M. D.; Butnariu, R. M.; Mangalagiu, I. I.: Synthesis and antimicrobial activity of some new pyridazine derivatives. *Heterocycles* **65** (2005) 1871–1879.
14. Said, M. A.; Aouad, M. R.; Hughes, D. L.; Almeahadi, M. A.; Messali, M.: Synthesis and crystal structure of a new pyridinium bromide salt: 4-methyl-1-(3-phenoxypropyl) pyridinium bromide. *Acta Crystallogr.* **E73** (2017) 1831–1834.