Yong Hong*, Wei Liu and Gaozhang Gou

Crystal structure of [4-chloro-2-(((2-((3-ethoxy-2-oxidobenzylidene)amino)phenyl)imino)(phenyl)methyl)phenolato- $\kappa^4 N, N', O, O'$]nickel(II) - ethyl acetate (1/1), $C_{32}H_{29}ClN_2NiO_5$

<https://doi.org/10.1515/ncrs-2018-0071>

Received May 27, 2018; accepted July 26, 2018; available online August 8, 2018

Abstract

$C_{32}H_{29}ClN_2NiO_5$, triclinic, $P\bar{1}$ (no. 2), $a = 9.7657(5)$ Å, $b = 10.1262(6)$ Å, $c = 16.0745(9)$ Å, $\alpha = 83.025(2)^\circ$, $\beta = 77.437(2)^\circ$, $\gamma = 66.240(2)^\circ$, $V = 1419.02(14)$ Å³, $Z = 2$, $R_{gt}(F) = 0.0403$, $wR_{ref}(F^2) = 0.0908$, $T = 293(2)$ K.

CCDC no.: 1822661

The crystal structure is shown in the figure. Tables 1 and 2 contain details on crystal structure and measurement conditions and a list of the atoms including atomic coordinates and displacement parameters.

Source of material

(*E*)-2-(((2-aminophenyl)imino)(phenyl)methyl)-4-chlorophenol (mono-imine) was synthesized by the method described

Table 1: Data collection and handling.

Crystal:	Colorless block
Size:	0.32 × 0.21 × 0.11 mm
Wavelength:	Mo $K\alpha$ radiation (0.71073 Å)
μ :	0.82 mm ⁻¹
Diffractometer, scan mode:	Bruker APEX-II, φ and ω -scans
θ_{max} , completeness:	25°, >99%
$N(hkl)_{measured}$, $N(hkl)_{unique}$, R_{int} :	32913, 4987, 0.053
Criterion for I_{obs} , $N(hkl)_{gt}$:	$I_{obs} > 2 \sigma(I_{obs})$, 4029
$N(param)_{refined}$:	372
Programs:	Bruker programs [1], SHELX [2]

[4]. 5-Chloro-2-hydroxybenzophenone (0.233 g, 1 mmol), 1,2-diaminobenzene (0.108 g, 1 mmol), piperidine (0.085 g, 1 mmol), and triethylorthoformate (0.148 g, 1 mmol) were refluxed for 2 h in ethanol (10 mL), then the solution was cooled and the mono-imine was obtained by filtration. Unsymmetrical tetradentate Schiff base ligand (*UTSBL*) was prepared by the obtained mono-imine and 3-ethoxy-2-hydroxybenzaldehyde. The mono-imine (0.323 g, 1 mmol) and 3-methoxy-2-hydroxybenzaldehyde (0.168 g, 1 mmol) were added to the solution of ethanol (10 mL) and refluxed for 2 h. The precipitate thus formed was collected by filtration. *UTSBL* (0.471 g, 1 mmol) and nickel acetate tetrahydrate (0.249 g, 1 mmol) were refluxed in methanol and ethyl acetate for about 1 h. After filtration, the solution was evaporated slowly at room temperature to obtain green prismatic crystals.

Experimental details

The H atoms were positioned geometrically with $d(C-H) = 0.93-0.98$ Å and refined as riding with $U_{iso}(H) = 1.2 U_{eq}(\text{carrier})$ or $1.5 U_{eq}(\text{methyl})$.

Discussion

The complex of symmetrical or unsymmetrical sal(salphen)tetradentate (N_2O_2) Schiff bases raise certain concerns for their intriguing applications in various fields [4–8]. There have been reported similar symmetrical crystal structures of Ni(II) [9, 10]. Herein, we report the crystal structure of a similar Ni(II) complex.

The dihedral angles between the aryl rings (C3–C4–C5–C6–C7–C8 and C10–C11–C12–C13–C14–C15), (C3–C4–C5–C6–C7–C8 and C23–C24–C25–C26–C27–C28),

*Corresponding author: Yong Hong, College of Science, Honghe University, Mengzi 661199, Yunnan Province, P.R. China; and Key Laboratory of Natural Pharmaceutical and Chemical Biology of Yunnan, Province, Mengzi 661199, Yunnan, P.R. China, e-mail: hongyong1018@163.com, hongyong1018@gmail.com

Wei Liu and Gaozhang Gou: College of Science, Honghe University, Mengzi 661199, Yunnan Province, P.R. China; and Key Laboratory of Natural Pharmaceutical and Chemical Biology of Yunnan, Province, Mengzi 661199, Yunnan, P.R. China

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	x	y	z	<i>U</i> _{iso} */ <i>U</i> _{eq}
Ni1	0.30792(4)	0.37454(4)	0.64762(2)	0.02772(11)
Cl1	−0.23263(13)	0.32020(12)	1.01227(6)	0.0815(3)
O2	0.42601(19)	0.47224(19)	0.65928(11)	0.0341(4)
O3	0.1748(2)	0.4804(2)	0.73702(12)	0.0391(5)
O1	0.5733(2)	0.5936(2)	0.71920(12)	0.0437(5)
N1	0.4329(2)	0.2829(2)	0.55049(13)	0.0279(5)
N2	0.2014(2)	0.2561(2)	0.64774(13)	0.0275(5)
O5	0.3524(3)	0.1789(3)	0.93543(15)	0.0694(7)
C23	0.0529(3)	0.3147(3)	0.78929(16)	0.0302(6)
C8	0.5644(3)	0.4484(3)	0.61874(16)	0.0302(6)
C16	0.1132(3)	0.2266(3)	0.71513(16)	0.0278(6)
C24	0.0796(3)	0.4414(3)	0.79322(16)	0.0316(6)
C15	0.2452(3)	0.1860(3)	0.56815(15)	0.0276(6)
C10	0.3713(3)	0.2028(3)	0.51498(16)	0.0274(6)
C12	0.3538(3)	0.0722(3)	0.40630(17)	0.0392(7)
H12	0.390948	0.032008	0.352885	0.047*
C17	0.0767(3)	0.0955(3)	0.71997(16)	0.0306(6)
C3	0.6490(3)	0.5137(3)	0.64882(17)	0.0333(6)
C11	0.4249(3)	0.1459(3)	0.43476(16)	0.0349(6)
H11	0.509197	0.157475	0.400047	0.042*
C25	−0.0006(3)	0.5311(3)	0.86344(17)	0.0378(7)
H25	0.013076	0.616352	0.865176	0.045*
C28	−0.0430(3)	0.2795(3)	0.85994(17)	0.0414(7)
H28	−0.056965	0.193763	0.860712	0.050*
C13	0.2263(3)	0.0582(3)	0.45769(17)	0.0397(7)
H13	0.176948	0.009949	0.437961	0.048*
C7	0.6362(3)	0.3630(3)	0.54644(17)	0.0325(6)
C14	0.1717(3)	0.1146(3)	0.53733(17)	0.0349(6)
H14	0.085347	0.105167	0.570836	0.042*
C22	−0.0700(3)	0.1067(3)	0.72112(17)	0.0393(7)
H22	−0.148540	0.197215	0.719599	0.047*
C27	−0.1152(3)	0.3679(3)	0.92664(18)	0.0446(8)
C9	0.5633(3)	0.2896(3)	0.51506(17)	0.0327(6)
H9	0.612731	0.241759	0.464511	0.039*
C2	0.6604(4)	0.6333(4)	0.7645(2)	0.0498(8)
H2A	0.698921	0.700806	0.729961	0.060*
H2B	0.746346	0.548553	0.777926	0.060*
C6	0.7849(3)	0.3484(3)	0.50412(19)	0.0418(7)
H6	0.830256	0.294362	0.455475	0.050*
C18	0.1913(3)	−0.0394(3)	0.72479(17)	0.0396(7)
H18	0.289069	−0.047430	0.726238	0.047*
C26	−0.0971(3)	0.4965(3)	0.92850(17)	0.0415(7)
H26	−0.150030	0.558086	0.973448	0.050*
C21	−0.0994(4)	−0.0165(4)	0.72453(19)	0.0514(9)
H21	−0.197897	−0.009101	0.725783	0.062*
C4	0.7939(3)	0.4944(3)	0.60712(19)	0.0405(7)
H4	0.848343	0.536289	0.627721	0.049*
C31	0.3058(5)	0.0584(4)	0.9518(3)	0.0772(12)
H31A	0.394328	−0.031728	0.949911	0.093*
H31B	0.249589	0.057986	0.909079	0.093*
C5	0.8609(3)	0.4126(3)	0.5342(2)	0.0443(8)
H5	0.958390	0.402279	0.506106	0.053*
C19	0.1621(4)	−0.1623(3)	0.72748(19)	0.0541(9)
H19	0.239839	−0.253042	0.730215	0.065*
C20	0.0167(5)	−0.1497(4)	0.7261(2)	0.0611(10)

Table 2 (continued)

Atom	x	y	z	<i>U</i> _{iso} */ <i>U</i> _{eq}
H20	−0.002722	−0.232395	0.726165	0.073*
C1	0.5594(4)	0.7013(4)	0.8442(2)	0.0716(11)
H1A	0.472789	0.782797	0.830338	0.107*
H1B	0.614432	0.732633	0.874790	0.107*
H1C	0.525632	0.632228	0.879123	0.107*
O4	0.4719(4)	0.1022(4)	0.8062(2)	0.1100(11)
C29	0.4823(6)	0.3156(5)	0.8531(3)	0.0885(13)
C32	0.2080(6)	0.0744(5)	1.0378(3)	0.0997(15)
H32A	0.121950	0.164885	1.039380	0.150*
H32B	0.265538	0.071984	1.079774	0.150*
H32C	0.173125	−0.003231	1.049850	0.150*
C30	0.4392(5)	0.1886(5)	0.8603(3)	0.0750(12)

(C3—C4—C5—C6—C7—C8 and C17—C18—C19—C20—C21—C22) are 15.19°, 37.47°, 56.80°, respectively. The dihedral angle between the two chelating ring system (Ni—N1—O2—C7—C8—C9 and Ni—N2—O3—C16—C23—C24) is 23.53° which indicated some distortion around the metal center. The packing of crystal structure is *via* van der Waals forces.

Acknowledgements: This work was financially supported by National Natural Science Foundation of China (Grant No. 21461007) and Open Fund of Master Construction Discipline in Yunnan Province (Grant No. HXB1401).

References

1. Bruker, *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, WI, USA (2009).
2. Sheldrick, G. M.: A short history of *SHELX*. *Acta Crystallogr. A* **64** (2008). 112–122.
3. Dolomanov, O. V.; Bourhis, L. J.; Gildea, R. J.; Howard, J. A. K.; Puschmann, H.: OLEX2: a complete structure solution, refinement and analysis program. *J. Appl. Crystallogr.* **42** (2009) 339–341.
4. Atkins, R.; Brewer, G.; Kokot, E.; Mockler, G. M.; Sinn, E.: Copper(II) and nickel(II) complexes of unsymmetrical tetradentate Schiff base ligands. *Inorg. Chem.* **24** (1985) 127–134.
5. Kleij, A. W.; Tooke, D. M.; Spek, A. L.; Reek, J. N. H.: A convenient synthetic route for the preparation of nonsymmetric metallo-salphen complexes. *Eur. J. Inorg. Chem.* **22** (2005) 4626–4634.
6. Kleij, A. W.: Nonsymmetrical salen ligands and their complexes: synthesis and applications. *Eur. J. Inorg. Chem.* **2** (2009) 193–205.
7. Escudero-Adán, E. C.; Belmonte, M. M.; Benet-Buchholz, J.; Kleij, A. W.: Practical approach to structurally diverse monimine salts and nonsymmetrical metallosalphen complexes. *Org. Lett.* **12** (2010) 4592–4595.
8. Whiteoak, C. J.; Salassa, G.; Kleij, A. W.: Recent advances with π -conjugated salen systems. *Chem. Soc. Rev.* **41** (2012) 622–631.

9. Bhattacharya, S.; Mondal, S.; Sasmal, S.; Sparkes, H. A.; Howard, J. A. K.; Nayak, M.; Mohanta, S.: Bis(nitrate)diaquauranyl(vi) synthon to generate $[1 \times 2 + 1 \times 1]$ and $[1 \times 1 + 1 \times 1]$ co-crystalized $3d \cdots 5f$ self-assemblies. *CrystEngComm* **13** (2011) 1029–1036.
10. San Felices, L.; Escudero-Adán, E. C.; Benet-Buchholz, J.; Kleij, A. W.: Isolation and structural characterization of a binuclear intermediate species pertinent to transmetalation of Zn(salphen) complexes and the formation of polynuclear salen structures. *Inorg. Chem.* **48** (2008) 846–853.