Mohammad Hayal Alotaibi, Hanan A. Mohamed, Bakr F. Abdel-Wahab, Amany S. Hegazy, Benson M. Kariuki* and Gamal A. El-Hiti*

Crystal structure of *N'*-(1-(2-hydroxyphenyl)ethylidene)-5-methyl-1-phenyl-1*H*-1,2,3-triazole-4-carbohydrazide, C₁₈H₁₇N₅O₂

<https://doi.org/10.1515/ncrs-2018-0417>

Received October 8, 2018; accepted October 26, 2018; available online November 13, 2018

Abstract

C₁₈H₁₇N₅O₂, monoclinic, *P*₂₁/*n* (no. 14), *a* = 20.3702(7) Å, *b* = 7.3482(2) Å, *c* = 23.2504(10) Å, β = 106.507(4)°, *V* = 3336.8(2) Å³, *Z* = 8, *R*_{gt}(*F*) = 0.0571, *wR*_{ref}(*F*²) = 0.1507, *T* = 296(2) K.

CCDC no.: 1875460

Table 1: Data collection and handling.

Crystal:	Colourless block
Size:	0.37 × 0.20 × 0.15 mm
Wavelength:	Mo <i>K</i> α radiation (0.71073 Å)
μ:	0.09 mm ^{−1}
Diffractometer, scan mode:	SuperNova, ω
θ _{max} , completeness:	29.7°, >99%
<i>N</i> (<i>hkl</i>) _{measured} , <i>N</i> (<i>hkl</i>) _{unique} , <i>R</i> _{int} :	32673, 8345, 0.033
Criterion for <i>I</i> _{obs} , <i>N</i> (<i>hkl</i>) _{gt} :	<i>I</i> _{obs} > 2 σ(<i>I</i> _{obs}), 5134
<i>N</i> (<i>param</i>) _{refined} :	458
Programs:	CrysAlis ^{PRO} [1], SHELX [2, 3], WinGX, ORTEP [4]

*Corresponding authors: Benson M. Kariuki, School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, UK, e-mail: kariukib@cardiff.ac.uk; and Gamal A. El-Hiti, Department of Optometry, College of Applied Medical Sciences, King Saud University, P.O. Box 10219, Riyadh 11433, Saudi Arabia, e-mail: gelhiti@ksu.edu.sa

Mohammad Hayal Alotaibi: National Center for Petrochemicals Technology, King Abdulaziz City for Science and Technology, P.O. Box 6086, Riyadh 11442, Saudi Arabia

Hanan A. Mohamed and Bakr F. Abdel-Wahab: Department of Chemistry, College of Science and Humanities, Shaqra University, Duwadimi, Saudi Arabia; and Applied Organic Chemistry Department, National Research Centre, Dokki, Giza, Egypt

Amany S. Hegazy: School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, UK

The asymmetric unit, containing two crystallographically independent molecules, of the title crystal structure is shown in the figure. Tables 1 and 2 contain details on crystal structure and measurement conditions and a list of the atoms including atomic coordinates and displacement parameters.

Source of material

The title compound was synthesized based on a literature procedure [5] and recrystallized from dimethylformamide to give colourless block crystals (86%).

Experimental details

All hydrogen atoms were placed in calculated positions and refined using a riding model. N–H bonds were fixed at

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} [*] / <i>U</i> _{eq}
C1	0.12057(9)	0.8874(2)	0.47514(9)	0.0489(4)
C2	0.08609(10)	0.8835(3)	0.41350(9)	0.0565(5)
C3	0.12032(12)	0.9311(3)	0.37181(10)	0.0689(6)
H3	0.097213	0.927515	0.331097	0.083*
C4	0.18760(12)	0.9833(3)	0.38990(11)	0.0691(6)
H4	0.210018	1.013592	0.361514	0.083*
C5	0.22207(10)	0.9909(3)	0.44999(11)	0.0644(6)
H5	0.267610	1.027692	0.462396	0.077*
C6	0.18894(10)	0.9440(3)	0.49135(10)	0.0582(5)
H6	0.212782	0.950030	0.531862	0.070*
C7	0.08730(9)	0.8324(3)	0.52096(9)	0.0509(4)
C8	0.12478(12)	0.8414(4)	0.58642(10)	0.0779(7)
H8A	0.102238	0.926203	0.605873	0.117*
H8B	0.171001	0.880668	0.591325	0.117*
H8C	0.125129	0.723117	0.604032	0.117*
C9	−0.07306(10)	0.6544(3)	0.52256(9)	0.0519(5)
C10	−0.10264(9)	0.6101(3)	0.57162(8)	0.0480(4)
C11	−0.16850(9)	0.5589(2)	0.56824(8)	0.0466(4)
C12	−0.22810(10)	0.5189(3)	0.51571(9)	0.0580(5)
H12A	−0.252997	0.417098	0.524717	0.087*
H12B	−0.212234	0.490555	0.481621	0.087*
H12C	−0.257561	0.623413	0.506853	0.087*
C13	−0.22141(9)	0.5069(2)	0.65369(8)	0.0466(4)
C14	−0.28706(9)	0.5689(3)	0.62847(9)	0.0563(5)
H14	−0.297984	0.632300	0.592378	0.068*
C15	−0.33675(10)	0.5359(3)	0.65739(11)	0.0658(6)
H15	−0.381453	0.575041	0.640188	0.079*
C16	−0.32030(11)	0.4458(3)	0.71119(11)	0.0650(6)
H16	−0.353582	0.426215	0.730834	0.078*
C17	−0.25458(11)	0.3844(3)	0.73618(10)	0.0611(5)
H17	−0.243494	0.323191	0.772692	0.073*
C18	−0.20502(10)	0.4134(3)	0.70715(9)	0.0554(5)
H18	−0.160779	0.369865	0.723658	0.066*
C19	0.49738(9)	0.2062(3)	0.61744(10)	0.0540(5)
C20	0.42966(10)	0.2652(3)	0.59025(10)	0.0621(5)
C21	0.38566(11)	0.2931(3)	0.62507(12)	0.0749(7)
H21	0.340885	0.329834	0.606804	0.090*
C22	0.40661(12)	0.2679(3)	0.68561(12)	0.0724(6)
H22	0.376246	0.288004	0.708201	0.087*
C23	0.47271(11)	0.2125(3)	0.71350(11)	0.0678(6)
H23	0.487254	0.195850	0.754838	0.081*
C24	0.51682(10)	0.1823(3)	0.67939(10)	0.0607(5)
H24	0.561273	0.144453	0.698396	0.073*
C25	0.54578(10)	0.1731(3)	0.58201(10)	0.0573(5)
C26	0.61401(10)	0.0825(3)	0.60817(11)	0.0732(6)
H26A	0.617815	−0.021000	0.584101	0.110*
H26B	0.617676	0.043331	0.648354	0.110*
H26C	0.650059	0.167180	0.608726	0.110*
C27	0.53785(10)	0.2418(3)	0.43069(10)	0.0608(5)
C28	0.58545(10)	0.2273(3)	0.39434(9)	0.0567(5)
C29	0.57247(9)	0.2507(3)	0.33380(9)	0.0540(5)
C30	0.50873(10)	0.3001(3)	0.28653(10)	0.0689(6)
H30A	0.520297	0.367360	0.255371	0.103*
H30B	0.480210	0.373509	0.303775	0.103*
H30C	0.484577	0.191345	0.269959	0.103*
C31	0.65374(10)	0.2345(3)	0.26991(10)	0.0561(5)

Table 2 (continued)

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} [*] / <i>U</i> _{eq}
C32	0.61490(11)	0.1478(3)	0.21909(10)	0.0640(5)
H32	0.575629	0.084315	0.219738	0.077*
C33	0.63476(13)	0.1562(3)	0.16747(11)	0.0741(6)
H33	0.608371	0.099877	0.132765	0.089*
C34	0.69322(15)	0.2471(4)	0.16680(13)	0.0838(8)
H34	0.706720	0.250872	0.131821	0.101*
C35	0.73196(14)	0.3326(4)	0.21768(15)	0.0876(8)
H35	0.771589	0.394283	0.216972	0.105*
C36	0.71239(11)	0.3275(3)	0.27001(12)	0.0734(6)
H36	0.738326	0.385813	0.304504	0.088*
N1	0.02520(8)	0.7749(2)	0.50145(7)	0.0536(4)
N2	−0.00829(8)	0.7204(2)	0.54201(7)	0.0564(4)
H2	0.011791	0.728044	0.579840	0.068*
N3	−0.06542(8)	0.6244(3)	0.63013(7)	0.0591(4)
N4	−0.10436(8)	0.5859(3)	0.66399(7)	0.0599(4)
N5	−0.16782(7)	0.5458(2)	0.62651(7)	0.0476(4)
N6	0.52485(9)	0.2239(2)	0.52699(9)	0.0629(5)
N7	0.56550(9)	0.2020(3)	0.48976(8)	0.0663(5)
H7	0.607043	0.164449	0.503194	0.080*
N8	0.65296(8)	0.1891(3)	0.41904(8)	0.0667(5)
N9	0.68330(8)	0.1874(3)	0.37689(8)	0.0685(5)
N10	0.63434(8)	0.2248(2)	0.32432(8)	0.0570(4)
O1	0.01960(8)	0.8355(3)	0.39170(7)	0.0796(5)
H1	0.005693	0.795392	0.419006	0.119*
O2	−0.10394(7)	0.6361(2)	0.46992(6)	0.0719(4)
O3	0.40414(8)	0.2961(3)	0.53053(8)	0.0874(5)
H3A	0.435081	0.288670	0.514508	0.131*
O4	0.47790(8)	0.2840(3)	0.41023(8)	0.0838(5)

0.86 Å (AFIX 43 instruction in SHELXL [4]), with displacement parameters 1.2 times *U*_{eq}(N). Aromatic C—H distances were set to 0.93 Å (AFIX 43) and their *U*_{iso}(H) set to 1.2 times the *U*_{eq}(C). Methyl C—H distances were set to 0.96 Å and their *U*_{iso}(H) to 1.5 times the *U*_{eq}(C) with the groups allowed to rotate about the C—C bonds (AFIX 137). O—H bonds were fixed at 0.82 Å (AFIX 147), with displacement parameters 1.2 times *U*_{eq}(O).

Comment

Various 1*H*-1,2,3-triazoles show miscellaneous biological activities [6–10]. Recently, we have reported the crystal structures for two 1*H*-1,2,3-triazole-4-carbohydrazides [11, 12].

The asymmetric unit consists of two crystallographically independent molecules (*cf.* the figure). Bond lengths and angles excellently fit with those of our previous experiments [11, 12]. The twist angles between the planes through the hydroxyphenyl and the triazolyl groups are 3.37(6)° for molecule (C1–C18, **A**) and 6.02(6)° for the molecule (C19–C36, **B**). The angles between the planes through the triazolyl and phenyl groups are 37.05(6)° for **A** and 46.14(6)° for **B**.

The molecules are stacked along [010] in the crystal structure with **A** and **B** forming separate columns.

Interactions of π – π type occur between hydroxyphenyl and triazolyl groups of neighbouring molecules related by inversion symmetry within the columns. The associated centroid-to-centroid distances are 3.97 Å and 3.66 Å for molecules **A** and **B** respectively. Intramolecular O–H $\cdots$ N hydrogen bonds are observed with O $\cdots$ N distances of 2.561(2) Å and 2.540(2) Å and O–H $\cdots$ N angles of 144.3° and 143.8° for **A** and **B** respectively.

Acknowledgements: Mohammad Hayal Alotaibi thanks King Abdulaziz City for Science and Technology (KACST), Saudi Arabia for financial support (Award No. 020-0180).

References

1. Rigaku Oxford Diffraction: CrysAlis^{PRO}. Rigaku Oxford Diffraction, Yarnton, England (2015).
2. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr. A* **64** (2008) 112–122.
3. Sheldrick, G. M.: Crystal structure refinement with SHELXL. *Acta Crystallogr. C* **71** (2015) 3–8.
4. Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Crystallogr.* **45** (2012) 849–854.
5. Youssef, N. S.; El-Zahany, E. A. M.; Drweesh, S. A.; Abdel-Wahab, B. F.; Ali, M. M.: Synthesis, characterization and anticancer activity of some transition metal complexes of new Schiff base triazole derivatives. *Smart Nanocomposites* **5** (2014) 89–109.
6. Ferreira, S. B.; Soderro, A. C.; Cardoso, M. F.; Lima, E. S.; Kaiser, C. R.; Silva, F. P.; Ferreira, V. F.: Synthesis, biological activity, and molecular modeling studies of 1*H*-1,2,3-triazole derivatives of carbohydrates as alpha-glucosidases inhibitors. *J. Med. Chem.* **53** (2010) 2364–2375.
7. Pokhodylo, N.; Shyyka, O.; Matychuk, V.: Synthesis of 1,2,3-triazole derivatives and evaluation of their anticancer activity. *Sci. Pharm.* **81** (2013) 663–676.
8. Reddy, P. V. B.; Kamala Prasad, V.; Manjunath, G.; Venkata Ramana, P. V.: Synthesis, characterization and evaluation of antibacterial activity of (*E*)-*N*-(substituted benzylidene)-2-(2-fluorobenzyl)-5-ethyl-2*H*-1,2,3-triazole-4-carbohydrazides. *Ann. Pharm. Fr.* **74** (2016) 350–357.
9. Sadeghpour, H.; Khabnadideh, S.; Zomorodian, K.; Pakshir, K.; Hoseinpour, K.; Javid, N.; Faghih-Mirzaei, E.; Rezaei, Z.: Design, synthesis, and biological activity of new triazole and nitro-triazole derivatives as antifungal agents. *Molecules* **22** (2017) 1150.
10. Slámová, K.; Marhol, P.; Bezouska, K.; Lindkvist, L.; Hansen, S. G.; Kren, V.; Jensen, H. H.: Synthesis and biological activity of glycosyl-1*H*-1,2,3-triazoles. *Bioorg. Med. Chem. Lett.* **20** (2010) 4263–4265.
11. Geesi, M. H.; Mohamed, H. A.; Abdel-Wahab, B. F.; Hegazy, A. S.; Kariuki, B. M.; El-Hiti, G. A.: Crystal structure of ethyl 4-amino-5-(5-methyl-1-(4-tolyl)-1*H*-1,2,3-triazole-4-carbonyl)-2-(phenylamino)thiophene-3-carboxylate, C₂₄H₂₃N₅O₃S. *Z. Kristallogr. NCS* **233** (2018) 673–674.
12. El-Hiti, G. A.; Abdel-Wahab, B. F.; Alotaibi, M. H.; Yousif, E.; Hegazy, A. S.; Kariuki, B. M.: 5-Methyl-1-(4-methylphenyl)-*N'*-[1-(1*H*-pyrrol-2-yl)ethylidene]-1*H*-1,2,3-triazole-4-carbohydrazide monohydrate. *IUCrData* **3** (2018) x181162.