Yee Seng Tan and Edward R.T. Tiekink*

Crystal structure of *catena*-poly[(bis(*O,O'*-diethyl dithiophosphato- κ^2S , S')- μ_2 -1,2-bis(3-pyridylmethylene)hydrazine- $\kappa^2N:N'$)cadmium(II)], $\{C_{20}H_{30}CdN_4O_4P_2S_4\}_n$

Abstract

$C_{20}H_{30}CdN_4O_4P_2S_4$, triclinic, $P\bar{1}$ (no. 2), $a = 9.0764(3)$ Å, $b = 9.3969(3)$ Å, $c = 9.6964(3)$ Å, $\alpha = 86.516(3)^\circ$, $\beta = 63.362(3)^\circ$, $\gamma = 75.250(3)^\circ$, $V = 713.50(4)$ Å³, $Z = 1$, $R_{gt}(F) = 0.0166$, $wR_{ref}(F^2) = 0.0462$, $T = 100(2)$ K.

CCDC no.: 1959871

Table 1 contains crystallographic data and Table 2 contains the list of the atoms including atomic coordinates and displacement parameters.

Table 1: Data collection and handling.

Crystal:	Yellow prism
Size:	$0.22 \times 0.14 \times 0.11$ mm
Wavelength:	Cu $K\alpha$ radiation (1.54184 Å)
μ :	10.21 mm^{-1}
Diffractometer, scan mode:	XtaLAB Synergy, ω -scans
$\theta_{\max}$, completeness:	67.1° , >99%
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	15508, 2560, 0.021
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 2560
$N(\text{param})_{\text{refined}}$:	162
Programs:	CrysAlis ^{PRO} [1], SHELX [2, 3], WinGX and ORTEP [4]

Source of material

The $Cd[S_2P(OEt)_2]_2$ precursor was prepared in high yield from the in situ reaction of $Cd(NO_3)_2 \cdot 4H_2O$ (Acros Organic; 15.42 g, 0.05 mol), EtOH (Merck; 12.25 mL, 0.21 mol), P_2S_5 (Sigma-Aldrich; 11.11 g, 0.05 mol) and 50% w/w NaOH solution (Merck; 8.80 mL, 0.11 mol). The 1,2-bis(3-pyridylmethylene)hydrazine ligand was prepared in high yield from the 2:1 reaction (reflux) of 3-picolylamine (Aldrich; 2.03 mL, 0.02 mol) and hydrazinium hydroxide (Merck; 0.49 mL, 0.01 mol) in ethanol solution (Merck; 5 mL). The title compound was obtained by mixing a suspension of $Cd[S_2P(OEt)_2]_2$ (0.50 g, 1.04 mmol) and 1,2-bis(3-pyridylmethylene)hydrazine (0.22 g, 1.05 mmol) in dimethylformamide (Merck; 5 mL), followed by stirring for 30 min. at 373 K. The solution was filtered and the filtrate collected in a sample vial containing acetonitrile (Merck; 1 mL). Yellow block crystals formed after one day. Yield: 0.39 g, (54.1%, based on $Cd[S_2P(OEt)_2]_2$). **M. pt** (MelTemp Melting Point Apparatus): 426–428 K. **Elem. Anal.** (Leco TruSpec Micro CHN Elemental Analyser): Calc.

<https://doi.org/10.1515/ncrs-2019-0655>

Received September 5, 2019; accepted October 17, 2019; available online November 14, 2019

*Corresponding author: Edward R.T. Tiekink, Research Centre for Crystalline Materials, School of Science and Technology, Sunway University, 47500 Bandar Sunway, Selangor Darul Ehsan, Malaysia, e-mail: edwardt@sunway.edu.my. <https://orcid.org/0000-0003-1401-1520>

Yee Seng Tan: Research Centre for Crystalline Materials, School of Science and Technology, Sunway University, 47500 Bandar Sunway, Selangor Darul Ehsan, Malaysia

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$).

Atom	x	y	z	U_{iso}^*/U_{eq}
Cd	0.500000	0.500000	1.000000	0.01167(6)
S1	0.62710(5)	0.21727(4)	0.89421(5)	0.01518(9)
S2	0.82782(5)	0.48870(4)	0.80199(5)	0.01587(9)
P1	0.84648(5)	0.27358(4)	0.78453(5)	0.01265(10)
O1	0.98987(15)	0.17908(13)	0.82879(14)	0.0187(2)
O2	0.92541(14)	0.21281(13)	0.61007(13)	0.0154(2)
N1	0.41685(17)	0.56317(15)	0.79881(15)	0.0140(3)
N2	0.49294(19)	0.93956(16)	0.54745(17)	0.0203(3)
C1	1.0169(2)	0.2188(2)	0.9563(2)	0.0202(4)
H1A	0.927426	0.198654	1.055805	0.024*
H1B	1.012810	0.324990	0.957880	0.024*
C2	1.1894(2)	0.1272(2)	0.9325(2)	0.0236(4)
H2A	1.216210	0.157193	1.011919	0.035*
H2B	1.275656	0.141812	0.830011	0.035*
H2C	1.188779	0.022857	0.940202	0.035*
C3	1.0940(2)	0.2290(2)	0.4983(2)	0.0199(4)
H3A	1.179689	0.184106	0.535561	0.024*
H3B	1.091571	0.334694	0.484423	0.024*
C4	1.1388(2)	0.1523(2)	0.3478(2)	0.0220(4)
H4A	1.255519	0.152879	0.273481	0.033*
H4B	1.059711	0.203682	0.306990	0.033*
H4C	1.130265	0.050186	0.365003	0.033*
C5	0.4415(2)	0.68340(18)	0.72021(19)	0.0151(3)
H5	0.500008	0.743068	0.741411	0.018*
C6	0.3846(2)	0.72530(19)	0.60771(19)	0.0174(3)
C7	0.2985(2)	0.6367(2)	0.5781(2)	0.0206(4)
H7	0.258755	0.661239	0.502101	0.025*
C8	0.2717(2)	0.5126(2)	0.6608(2)	0.0209(4)
H8	0.212395	0.451349	0.643116	0.025*
C9	0.3327(2)	0.47946(19)	0.76937(19)	0.0170(3)
H9	0.314491	0.394049	0.825640	0.020*
C10	0.4139(2)	0.8573(2)	0.5218(2)	0.0203(4)
H10A	0.373770	0.882130	0.446030	0.024*

for $C_{20}H_{30}CdN_4O_4P_2S_4$: C, 34.66; H, 4.36; N, 8.09%. Found: C, 34.51; H, 4.08; N, 8.31%. IR (Bruker Vertex 70 V equipped with Platinum ATR from 400 to 80 cm^{-1}): 1192(w) ν (C—O); 1012(s) ν (P—O); 661(s) ν (P—S); 379(w) ν (Cd—N); 287(m) ν (Zn—S).

Experimental details

H atoms were geometrically placed (C—H = 0.95–0.99 $\text{\AA}$) and refined as riding with $U_{iso}(H) = 1.2\text{--}1.5U_{eq}(C)$.

Discussion

The potentially bidentate, bridging ligands, *n*-pyridylaldazines (*n*-PyAld), i.e. $n\text{-NC}_5\text{H}_4\text{C(H)=N—N=C(H)C}_5\text{H}_4\text{N-}n$, for $n = 3$ and 4, form one-dimensional coordination polymers in their 1:1 adducts with zinc(II) and cadmium(II) dithiophosphates $[\text{S}_2\text{P(OR)}_2]^-$ [5]. However, very different coordination geometries and topologies of the chains are found in these structures. Thus, for $\{\text{Zn}[\text{S}_2\text{P(OR)}_2](n\text{-PyAld})\}_2$, a distorted tetrahedral N_2S_2 coordination geometry and a twisted chain is found when $n = 3$

and $R = \text{Et}$ [6]. By contrast, when $n = 3$ and $R = \text{isopropyl}$ (iPr), a linear chain with a step-ladder topology and an octahedral *trans*- N_2S_4 donor set is found [7]. In the case when $n = 4$, and $R = \text{Et}$ [8] and cyclohexyl (Cy) [9], zigzag coordination polymers are found but with tetrahedral N_2S_2 and octahedral *cis*- N_2S_4 donor sets, respectively. A greater homogeneity in the coordination polymers formed by cadmium(II) is noted in that *trans*- N_2S_4 donor sets are found. Thus, for $\{\text{Cd}[\text{S}_2\text{P(OR)}_2](n\text{-PyAld})\}_2$, when $n = 3$ and $R = \text{Me}$ [10], iPr [11, 12] and Cy [13], linear coordination polymers, with a step-ladder topology, are found. When $n = 4$, and $R = \text{Et}$ [14] and iPr [11], linear coordination polymers are found. However, it should be noted that zero-dimensional species are sometimes isolated when polymeric species might otherwise be anticipated. An example is seen in the structure of binuclear $\{\text{Zn}[\text{S}_2\text{P(O-iPr)}_2]_2(4\text{-PyAld})\}_2$ [15]. More intriguing are monodentate (terminal) modes of coordination of the 4-PyAld ligand have been noted in some dithiocarbamate (S_2CNR_2) derivatives, namely $\text{Zn}[\text{S}_2\text{CN(Me)CH}_2\text{CH}_2\text{OH}]_2(4\text{-PyAld})$, with a square-pyramidal geometry based on a NS_4 donor set [16] and $\text{Cd}[\text{S}_2\text{CN(nPr)CH}_2\text{CH}_2\text{OH}]_2(4\text{-PyAld})_2$ [*n*-propyl = *n*Pr], with a *trans*- N_2S_4 donor set [17]. In continuation of systematic studies in this area, herein, the crystal and molecular structures of $\{\text{Cd}[\text{S}_2\text{P(OEt)}_2]_2(3\text{-PyAld})\}_n$, (I), are described.

The cadmium(II) atom in (I) lies on a centre of inversion (the symmetry related dithiophosphate ligand is generated by the symmetry operation (i) $1 - x, 1 - y, 2 - z$) and the 3-PyAld molecule is disposed about a centre of inversion (the other half of the 3-PyAld ligand is related by (ii) $1 - x, 2 - y, 1 - z$). A view of the coordination geometry is shown in the figure (70% probability displacement ellipsoids) which highlights the *trans*- N_2S_4 donor set. While to a first approximation, the dithiophosphate ligands are symmetrically chelating to define a plane defined by four S atoms, there is a disparity of 0.026 $\text{\AA}$ in the Cd—S1 [2.6741(4) $\text{\AA}$] and Cd—S2 [2.7003(4) $\text{\AA}$] bond lengths. Nevertheless, the P1—S1, S2 [1.9862(6) and 1.9963(6) $\text{\AA}$] differ by only 0.01 $\text{\AA}$. Significantly, more symmetric coordination modes are noted in the $R = \text{Me}$ [10] and iPr [12] analogues but, the situation in (I) matches that seen in the $R = \text{Cy}$ structure [13] where the difference in the Cd—S bond lengths amounts to 0.021 $\text{\AA}$. The Cd—N bond length is 2.3830(13) $\text{\AA}$. The lack of systematic trends in the Zn, Cd—S, N bond lengths is well-known for these systems, even if control of polymer formation and topology was achieved [18–20]. The coordination geometry in (I) is defined by the *trans*- N_2S_4 donor set and is based on an octahedron with the maximum deviation from the ideal *cis* angle manifested in the acute S1—Cd—S2 chelate angle of 77.236(12)°.

The linear chain in (I), with a step-ladder topology, is parallel to $[0\ 1\ -1]$. Chains are connected into a two-dimensional array by $\pi \cdots \pi$ stacking between pyridyl rings

[inter-centroid separation for $Cg(N1,C5-C9) \cdots Cg(N1,C5-C9)^{iii} = 3.6794(10)$ Å for (iii) $1 - x, 1 - y, 1 - z$] and pyridyl- $C-H \cdots O$ (alkoxy) [$C7-H7 \cdots O2^{iii}$: $H7 \cdots O2^{iii} = 2.42$ Å, $C7 \cdots O2^{iii} = 3.318(2)$ Å with angle at H7 = 158°] interactions. Layers stack along $[-2\ 2\ 1]$ and inter-digitate with successive layers but there are no directional interactions between them.

Acknowledgements: Sunway University Sdn Bhd is thanked for financial support of this work through Grant no. STR-RCTR-RCCM-001-2019.

References

1. Rigaku Oxford Diffraction: CrysAlis PRO. Rigaku Corporation, Oxford, UK (2018).
2. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr. A* **64** (2008) 112–122.
3. Sheldrick, G. M.: Crystal structure refinement with SHELXL. *Acta Crystallogr. C* **71** (2015) 3–8.
4. Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Cryst.* **45** (2012) 849–854.
5. Tiekink, E. R. T.: Perplexing coordination behaviour of potentially bridging bipyridyl-type ligands in the coordination chemistry of zinc and cadmium 1,1-dithiolate compounds. *Crystals* **8** (2018) article no. 18 (29 pages).
6. Tan, Y. S.; Tiekink, E. R. T.: Crystal structure of *catena*-[(bis(*O,O'*-diethyl dithiophosphato-*S,S'*)- μ_2 -1,2-bis(3-pyridylmethylene)hydrazine-*N,N'*)zinc(II)], $\{C_{20}H_{30}N_4O_4P_2S_4Zn\}_n$. *Z. Kristallogr. NCS* **235** (2020) 297–299.
7. Avila, V.; Tiekink, E. R. T.: *catena*-Poly[[bis(*O,O'*-diisopropyl dithiophosphato- κ^2S,S')zinc(II)]- μ -1,2-bis(3-pyridylmethylene)hydrazine- $\kappa^2N:N'$]. *Acta Crystallogr. E* **62** (2006) m3530–m3531.
8. Tan, Y. S.; Tiekink, E. R. T.: Crystal structure of *catena*-{poly[bis(*O,O'*-diethyldithiophosphato-*S*)-(μ_2 -1,2-bis(4-pyridylmethylene)hydrazine-*N,N'*)-zinc(II)] di-acetonitrile solvate}, $\{C_{20}H_{30}N_4O_4P_2S_4Zn \cdot 2C_2H_3N\}_n$. *Z. Kristallogr. NCS* **234** (2019) 895–897.
9. Chen, D.; Lai, C. S.; Tiekink, E. R. T.: *catena*-Poly[[bis(*O,O'*-dicyclohexyldithiophosphato- κ^2,S,S')zinc(II)]- μ -1,2-bis(4-pyridylmethylene)hydrazine- $\kappa N:N'$]. *Acta Crystallogr. E* **61** (2005) m2052–m2054.
10. Tan, Y. S.; Tiekink, E. R. T.: Crystal structure of *catena*-[(bis(*O,O'*-dimethyl dithiophosphato-*S,S'*)- μ_2 -1,2-bis(3-pyridylmethylene)hydrazine-*N,N'*)cadmium(II)], $\{C_{20}H_{30}CdN_4O_4P_2S_4\}_n$. *Z. Kristallogr. NCS* **235** (2020) 339–341.
11. Lai, C. S.; Tiekink, E. R. T.: Polymeric topologies in cadmium(II) dithiophosphate adducts of the isomeric *n*-pyridinealdazines, $n = 2, 3$ and 4. *Z. Kristallogr. C. M.* **221** (2006) 288–293.
12. Tan, Y. S.; Tiekink, E. R. T.: A redetermination of the crystal structure of *catena*-[(bis(*O,O'*-isopropyl dithiophosphato-*S,S'*)-(μ_2 -1,2-bis(3-pyridylmethylene)hydrazine-*N,N'*)cadmium(II)], $\{C_{24}H_{38}CdN_4O_4P_2S_4\}_n$. *Z. Kristallogr. NCS* **235** (2020) 253–255.
13. Lai, C. S.; Tiekink, E. R. T.: Delineating the principles controlling polymer formation and topology in zinc(II)- and cadmium(II)-dithiophosphate adducts of diimine-type ligands. *J. Molec. Struct.* **796** (2006) 114–118.
14. Tan, Y. S.; Tiekink, E. R. T.: Crystal structure of *catena*-[(bis(*O,O'*-diethyl dithiophosphato-*S,S'*)- μ_2 -1,2-bis(4-pyridylmethylene)hydrazine-*N,N'*)cadmium(II)], $\{C_{20}H_{30}CdN_4O_4P_2S_4\}_n$. *Z. Kristallogr. NCS* **235** (2020) 335–337.
15. Tan, Y. S.; Tiekink, E. R. T.: Crystal structure of *tetrakis*(*O,O'*-diisopropyldithiophosphato- κ^2S,S')-(μ_2 -1,2-bis(4-pyridylmethylene)hydrazine- $\kappa^2N:N'$)zinc(II), $C_{36}H_{66}N_4O_8P_4S_8Zn_2$. *Z. Kristallogr. NCS* **234** (2019) 827–829.
16. Broker, G. A.; Jotani, M. M.; Tiekink, E. R. T.: Bis[*N*-2-hydroxyethyl, *N*-methyldithiocarbamato- κ^2S,S']-4-[(pyridin-4-ylmethylidene)hydrazinylidene)methyl]pyridine- κN^1) zinc(II): crystal structure and Hirshfeld surface analysis. *Acta Crystallogr. E* **73** (2017) 1458–1464.
17. Broker, G. A.; Tiekink, E. R. T.: Bis[*N*-(2-hydroxyethyl)-*N*-propyldithiocarbamato- κ^2S,S']bis-(4-[(pyridin-4-ylmethylidene)hydrazinylidene)methyl]pyridine- κN^1)cadmium. *Acta Crystallogr. E* **67** (2011) m320–m321.
18. Lai, C. S.; Liu, S.; Tiekink, E. R. T.: Steric control over polymer formation and topology in adducts of zinc dithiophosphates formed with bridging bipyridine ligands. *CrystEngComm* **6** (2004) 221–226.
19. Lai, C. S.; Tiekink, E. R. T.: Engineering polymers with variable topology – bipyridine adducts of cadmium dithiophosphates. *CrystEngComm* **6** (2004) 593–605.
20. Chen, D.; Lai, C. S.; Tiekink, E. R. T.: Supramolecular aggregation in diimine adducts of zinc(II) dithiophosphates: controlling the formation of monomeric, dimeric, polymeric (zig-zag and helical), and 2-D motifs. *CrystEngComm* **8** (2006) 51–58.