The model has been extended to impurity diffusion^{2a}. In view of a discussion to follow in^{2a} the experimental values of A and B are listed in Table 1.

This work has received financial support from the Swedish Natural Science Research Council and the Swedish Board for Technical Development. We thank fil. lic. C. ROXBERGH, fil. mag. P.-E. ERIKSSON, and ing. H. OLSSON for skilful help with the experiments and Dr. A. LODDING for interest, encouragement and discussions.

Table 1. Arrhenius and linear parameters for diffusion in liquid Na and K.

System	$D_0 \cdot 10^3 (\text{cm}^2/\text{s})$	Q (kcal/mole)	$A \cdot 10^7 \text{cm}^2$ $\text{s}^{-1} \text{grad}^{-1}$	$B (^\circ\text{K})$
K in Na	0.46 ± 0.12	1.82 ± 0.12	2.72 ± 0.28	225 ± 20
Na in Na	0.86 ± 0.09	2.22 ± 0.08	3.90 ± 0.13	263 ± 9
Na in K	0.93 ± 0.19	2.11 ± 0.16	4.75 ± 0.30	261 ± 15
K in K	0.76 ± 0.06	2.02 ± 0.07	3.89 ± 0.11	244 ± 9

The ratio of the nuclear g_I -factors of ^{95}Mo and ^{97}Mo

H. KRÜGER, O. LUTZ, A. NOLLE and A. SCHWENK

Physikalisches Institut der Universität Tübingen, 74 Tübingen, Gmelinstraße 6, Germany

Physikalisches Institut der Universität Tübingen

(Z. Naturforsch. 28a, 119–120 [1973]; received 1 December 1972)

With a multi-nuclei Fourier transform NMR spectrometer the ratio of the Larmorfrequencies of ^{95}Mo and ^{97}Mo in a solution of K_2MoO_4 in H_2O has been measured with high accuracy. Using hyperfine interaction constants from literature, we have calculated the hyperfine structure anomaly $^{95}\Delta^{97} = -0.0102(14)\%$ of molybdenum in the atomic ground state $4d^{55}s^7S_3$.

Introduction

For many isotopic pairs values of the hyperfine structure anomaly (HFSA) have been quoted¹. For the two stable odd isotopes of molybdenum ^{95}Mo and ^{97}Mo recently PENDLEBURY and RING² have measured the magnetic hyperfine interaction constants in the $4d^{55}s^7S_3$ atomic ground state: $A(^{95}\text{Mo}) = -208.5790$ (20) MHz, $A(^{97}\text{Mo}) = -212.9781$ (20) MHz. Together with the ratio of the g_I -factors of PROCTOR and YU³, which has an accuracy of only $1 \cdot 10^{-4}$, PENDLEBURY and RING² gave a value of the HFSA in the atomic ground state: $^{95}\Delta^{97} = -0.01(1)\%$. The error results from the NMR measurement of PROCTOR and YU³. If one therefore knows the ratio of the g_I -factors, which is equal to the ratio of the Larmor-frequencies, with an accuracy of at least $1 \cdot 10^{-5}$, one can get a more meaningful value of the HFSA of the molybdenum isotopes ^{95}Mo and ^{97}Mo . A nonvanishing value of the HFSA can give information about both the electronic configuration and some nuclear properties.

The Hyperfine structure anomaly

The HFSA of two isotopes 1 and 2 is defined by

$$^1\Delta^2 = (A_1/A_2) \cdot (g_{I2}/g_{I1}) - 1 \quad (1)$$

A_1, A_2 are the experimental magnetic hyperfine interaction constants of the atomic state in question; g_{I1}, g_{I2} are the nuclear g -factors. The value of Δ is usually given in $\%$.

The magnetic hyperfine interaction constant, can be written as:

$$A = A_{\text{point}} (1 - \delta) (1 - \varepsilon), \quad (2)$$

where A_{point} is the hyperfine interaction constant, that one would get, if the observed nucleus were a point nucleus, δ is a correction considering the distribution of the electric charge over the volume of the nucleus^{4, 5, 6} and ε is a correction considering the distribution of the magnetization⁷⁻¹⁰. δ and ε take values of a few percents for $s_{1/2}$ and $p_{1/2}$ electrons; in other electronic states, if relativistic effects and effects of configuration mixing can be neglected, one finds: $\delta = \varepsilon = 0$. Taking into account the definition of A_{point} one gets the following equation for the HFSA

$$^1\Delta^2 = \frac{(1 - \delta_1)(1 - \varepsilon_1)}{(1 - \delta_2)(1 - \varepsilon_2)} - 1, \quad (3)$$

if reduced mass effects can be neglected. This is the case for the two molybdenum isotopes. $\varepsilon_1, \varepsilon_2, \delta_1, \delta_2$ are the corrections for the respective isotope 1 or 2.

Therefore Eqn. (1) can be written in a fairly good approximation as:

$$^1\Delta^2 \approx (\delta_2 - \delta_1) + (\varepsilon_2 - \varepsilon_1) = \Delta_\delta + \Delta_\varepsilon \quad (4)$$

Eqn. (4) shows the physical meaning of the HFSA.

¹ G. H. FULLER and V. W. COHEN, ORNL-4591 [1970].

² J. M. PENDLEBURY and D. B. RING, J. Phys. B5, 386 [1972].

³ W. G. PROCTOR and F. C. YU, Phys. Rev. 81, 20 [1951].

⁴ J. E. ROSENTHAL and G. BREIT, Phys. Rev. 41, 459 [1932].

⁵ M. F. CRAWFORD and A. L. SCHAWLOW, Phys. Rev. 76, 1310 [1949].

⁶ H. J. ROSENBERG and H. H. STROKE, Phys. Rev. A5, 1992 [1972].

⁷ A. BOHR and V. F. WEISSKOPF, Phys. Rev. 77, 94 [1950].

⁸ J. EISINGER and V. JACCARINO, Rev. Mod. Phys. 30, 528 [1958].

⁹ N. J. IONESCO-PALLAS, Phys. Rev. 117, 505 [1960].

¹⁰ H. H. STROKE, R. J. BLIN-STOYLE, and V. JACCARINO, Phys. Rev. 123, 1326 [1961].

Measurement of the ratio of the Larmor frequencies

The NMR signals of the molybdenum isotopes have been detected in a field of 1.807 tesla by a Bruker B-KR 322s pulse spectrometer. The signals were enhanced by a Hewlett-Packard signal analyzer 5480 A and the Fourier transformation was computed on the CDC 3300 of the Zentrum für Datenverarbeitung der Universität Tübingen. The magnetic field was held constant with the aid of a ^7Li NMR probe ¹¹.

^{97}Mo has relatively short relaxation times ¹². The Fourier transformation of the free induction decay yields the Larmorfrequency. The linewidth of the NMR signal of ^{97}Mo of 100 Hz is partly due to the inhomogeneity of the magnetic field and partly due to relaxation processes.

Though both isotopes have nuclear spin $I = 5/2$, ^{95}Mo has long relaxation times ¹². Therefore a special technique, the so-called Quadriga Fourier transform Spectroscopy of SCHWENK ¹³ was used for recording the NMR signal of ^{95}Mo . The linewidth of 60 Hz of this signal is due to the used technique.

The Larmor frequencies of ^{95}Mo (about 5.014 MHz) and of ^{97}Mo (about 5.119 MHz) were measured alternately in the same probe and sample (10 mm tube diameter) only by varying the excitation frequency, at a temperature of (301 ± 3) K. The sample was a 3 molal solution of K_2MoO_4 in H_2O . For the NMR signal of both molybdenum isotopes a signal to noise ratio of 50 was achieved, within summation times of about 6 minutes.

We have measured 67 ratios of the Larmor-frequencies of ^{95}Mo and ^{97}Mo at four different days. The result is:

$$\nu(^{95}\text{Mo}) / \nu(^{97}\text{Mo}) = 0.979\,445\,(1)$$

All the measured ratios lie within the given limits of error.

PROCTOR and YU ³ gave the following value of that ratio:

$$\nu(^{95}\text{Mo}) / \nu(^{97}\text{Mo}) = 0.9794(1)$$

These values are in agreement within their limits of error.

¹¹ J. KAUFMANN and A. SCHWENK, Z. Angew. Phys. 21, 527 [1966].

The HFSA of ^{95}Mo and ^{97}Mo

Using the A-factors of PENDLEBURY and RING ² of the atomic ground state $4d^5 5s\,^7\text{S}_3$ and our ratio of the Larmor frequencies of ^{95}Mo and ^{97}Mo we get for the HFSA:

$$^{95}\Delta^{97}(4d^5 5s\,^7\text{S}_3) = -0.0102(14)\%.$$

The HFSA is relatively large because of the dominant contribution of the unpaired s-electron to the A-factor. The error results from the measurement of PENDLEBURY and RING ². ROSENTHAL and BREIT ⁴ and CRAWFORD and SCHAWLOW ⁵ have given a formula for δ based on the assumption, that the nuclear radius varies proportional to $(\text{number of nucleons})^{1/3}$ and that the charge distribution is constant over the nuclear volume. From this formula one gets for a $s_{1/2}$ electron in the case of molybdenum:

$$\Delta_\delta = 0.016\%.$$

ROSENBERG and STROKE ⁶ have used a more realistic model of the distribution of the electric charge. They get for a $s_{1/2}$ electron in the case of molybdenum:

$$\Delta_\delta = 0.007\%.$$

If one assumes, that the HFSA of molybdenum is wholly due to the 5s electron, one can calculate from Eqn. (4) the contribution to the HFSA, which is due to the distribution of the magnetization over the volume of the nuclei. We get:

$$\Delta_\epsilon = -0.017\%.$$

One can see, that we get a relatively large contribution to the HFSA, though the magnetic moments of both isotopes are only 2 % different from each other.

Acknowledgement: We like to thank Dr. J. KAUFMANN for helpful remarks and the Deutsche Forschungsgemeinschaft for the financial support of this work.

¹² J. KAUFMANN, Z. Phys. 182, 217 [1964].

¹³ A. SCHWENK, J. Mag. Res. 5, 376 [1971].