On a T_1 Study of Molecular Motion in Some Tetra Substituted Ammonium Halides

Fevzi Köksal

Department of Physics, Faculty of Sciences, Ankara University, Turkey

Z. Naturforsch. **34a**, 1296–1300 (1979); received August 9, 1979

The spin-lattice relaxation times of tetramethylammonium bromide $(\text{CH}_3)_4\text{NBr}$, iodide $(\text{CH}_3)_4\text{NI}$, tetraethylammonium chloride $(\text{CH}_3\text{CH}_2)_4\text{NCl}$, bromide $(\text{CH}_3\text{CH}_2)_4\text{NBr}$, iodide $(\text{CH}_3\text{CH}_2)_4\text{NI}$, tetrapropylammonium iodide $[\text{CH}_3(\text{CH}_2)_3]_4\text{NI}$, tetrabutylammonium bromide $[\text{CH}_3(\text{CH}_2)_4]_4\text{NBr}$, iodide $[\text{CH}_3(\text{CH}_2)_4]_4\text{NI}$, tetraheptylammonium bromide $[\text{CH}_3(\text{CH}_2)_6]_4\text{NBr}$ and iodide $[\text{CH}_3(\text{CH}_2)_6]_4\text{NI}$ were measured at 60 MHz in the temperature interval 100 K to near the melting points. Two minima of T_1 were observed for tetramethylammonium halides, for tetraethylammonium bromide and iodide and for tetrapropylammonium iodide. The minima in T_1 were attributed to CH_3 group reorientations and reorientations of the tetramethyl, ethyl and propylammonium cations. The activation energies of the processes were determined.

1. Introduction

Motional properties of compounds of the type $(\text{CH}_3)_4\text{X}$ where X is C, Si, Ge, Sn, Pb and $(\text{CH}_3)_4\text{AsZ}$ where Z is Cl, Br and I were studied by broad line nmr and pulsed nmr techniques [1–7]. In addition, $(\text{CH}_3)_4\text{NZ}$ type compounds were studied [8] at 25.3 MHz and two minima were observed at T_1 . One of these minima was attributed to CH_3 group reorientations and the other to reorientations of the cations in the compounds. Therefore it seemed interesting to study larger molecules and it is the purpose of this study to investigate the motional properties of tetramethylammonium (TMA) bromide, iodide; tetraethylammonium (TEA) chloride, bromide iodide; tetrapropylammonium (TPA) iodide, tetrabutylammonium (TBA) bromide, iodide; tetraheptylammonium (THA) bromide and iodide by pulsed nmr at 60 MHz.

2. Experimental Considerations and Results

The spin-lattice relaxation time measurements were performed at 60 MHz using an SXP type 4-100 MHz Bruker pulse spectrometer employing $180^\circ\text{--}\tau\text{--}90^\circ$ pulse sequence. The compounds were obtained from commercial sources, and before use they were dried under 10^{-3} torr vacuum for two days by keeping the samples at 323 K. The samples were 0.8 cm in diameter and 0.3 cm in height and the measurements were made between 100 and

490 K. The temperature control system used was a Bruker B-ST 100/700 controller. Most of the measurements were repeated three times and found to be well reproducible.

The experimental values of T_1 were plotted against $10^3 \text{ K}/T$ as shown in Figures 1–10. For TMA bromide the spin-lattice relaxation time passes through two minimum at 323 and 385 K with $T_1 = 29$ ms and $T_1 = 38$ ms, respectively. The T_1 of TMA iodide also passes through two minima at 308 K with $T_1 = 29$ ms and at 355 K with $T_1 = 36$ ms.

For TEA chloride we observed only a minimum at 130 K with $T_1 = 80$ ms; because of its low melting point we did not go to higher temperatures. However for TEA bromide the experiments were made between 485 and 120 K, and T_1 passes through two minima at 166 K with $T_1 = 86$ ms and at 425 K with $T_1 = 1.6$ s. Similarly, for TEA iodide we observed two minima at 113 K with $T_1 = 42$ ms and at 444 K with $T_1 = 2.3$ s.

For TPA iodide the experiments were made between 485 and 120 K and two minima in T_1 were observed. One of the minima occurs at 158 K with $T_1 = 86$ ms and the other at 400 K with $T_1 = 0.28$ s.

For TBA bromide and iodide the experiments were performed between 335 and 100 K. Due to their low melting points we did not go to higher temperatures, and only one minimum was observed for each of the compounds. The T_1 minimum for TBA bromide occurs at 125 K with $T_1 = 80$ ms and for TBA iodide the minimum occurs at 154 K with $T_1 = 80$ ms.

Reprint requests to Dr. F. Köksal, A. Ü. Fen Fakültesi Fizik Bölümü, Beşevler-Ankara, Turkey.

Fig. 1. Spin-lattice relaxation time of TMA bromide against 10^3 K/T.

Fig. 2. Spin-lattice relaxation time of TMA iodide against 10^3 K/T.

Fig. 3. Spin-lattice relaxation time of TEA chloride against 10^3 K/T.

Fig. 4. Spin-lattice relaxation time of TEA bromide against 10^3 K/T.

Fig. 5. Spin-lattice relaxation time of TEA iodide against 10^3 K/T.

Fig. 6. Spin-lattice relaxation time of TPA iodide against 10^3 K/T.

Fig. 7. Spin-lattice relaxation time of TBA bromide against 10^3 K/T.

Fig. 8. Spin-lattice relaxation time of TBA iodide against 10^3 K/T.

Fig. 9. Spin-lattice relaxation time of THA bromide against 10^3 K/T.

Fig. 10. Spin-lattice relaxation time of THA iodide against 10^3 K/T.

For THA bromide and iodide the T_1 minima occur respectively at 130 and 138 K and the minimum values of T_1 are equal and $T_1 = 0.16$ s.

3. Analysis of Results

The two minima in T_1 for TMA bromide and iodide indicate two types of motion as explained by Albert et al. [8]. The low temperature minima were attributed to methyl group reorientations, and the high temperature minimum were attributed to reorientational motions of the $(\text{CH}_3)_4\text{N}$ cation. The calculated spin-lattice relaxation rate including the proton-proton dipolar interactions in the methyl groups and the intra methyl group dipolar inter-

actions within the same molecule is given by [8]

$$T_1^{-1} = A g(\omega_0, \tau_{c2}) + B g(\omega_0, \tau_{c1}), \quad (1)$$

where $g(\omega_0, \tau_{ci})$ are the spectral density functions and

$$A = \frac{9}{20} \frac{\gamma^4 \hbar^2}{r^6}, \quad B = \frac{3}{20} \frac{\gamma^4 \hbar^2}{r^6} + \frac{27}{10} \frac{\gamma^4 \hbar^2}{r_1^6}.$$

Furthermore,

$$\tau_{c2}^{-1} = \tau_c^{-1} + \tau_{c1}^{-1}. \quad (2)$$

In these equations τ_c is the correlation time for methyl group reorientations, τ_{c1} the correlations time for the reorientations of the cation, ω_0 is the Larmor frequency, r is the proton-proton distance within the methyl group and r_1 is the distance between the centers of the methyl groups within the same molecule. When $\tau_c \ll \tau_{c1}$ the minimum value of T_1 can be written [6]

$$T_{1\min}^{-1} = \frac{9}{20} \frac{\gamma^4 \hbar^2}{r^6} \frac{1.43}{\omega_0}. \quad (3)$$

Incorporation of the usual constants and $r = 1.79 \text{ \AA}$ in (3) gives $T_{1\min} = 34 \text{ ms}$. The low temperature experimental T_1 minima for TMA bromide and iodide are equal and 29 ms. This is somewhat shorter than the calculated value, however the discrepancy is not great and comes from the dipolar interactions between the methyl group within the same molecule. The correlation times where supposed to satisfy the Arrhenius equation

$$\tau_{ci} = \tau_{ci}^0 \exp(E_a/RT), \quad (4)$$

and the activation energy E_a corresponding to the linear portion above the low temperature minimum in Fig. 1 was found to be $2.1 \text{ kcal mol}^{-1}$. The activation energy corresponding to the linear portion above the second minimum is $5.7 \text{ kcal mol}^{-1}$. The activation energy for methyl group reorientations in TMA iodide cannot be obtained from the experimental results due to the short temperature interval between the minima. But the activation energy above the high temperature minimum was found to be $5.6 \text{ kcal mol}^{-1}$. These activation energies are smaller than the value obtained by Albert et al. [8], but the frequency dependence of the low temperature experimental T_1 minima for both compounds agree with the frequency ratio to 25.3 MHz at which the experiments were performed. Furthermore our $E_a = 2.1 \text{ kcal mol}^{-1}$ for methyl

group reorientations is in agreement with the E_a 's of methyl groups reported by several authors [2, 3, 5, 6, 7, 9].

For TEA chloride we have observed only one minimum in T_1 and the $T_{1\min} = 89 \text{ ms}$ does not agree with the calculated $T_{1\min} = 34 \text{ ms}$. The activation energy corresponding to the linear portion in Fig. 3 is found to be $2.0 \text{ kcal mol}^{-1}$. However for TEA bromide and iodide we have observed two minima in T_1 which again can be attributed to methyl group reorientations and cation reorientations. The activation energies above the low temperature minima are 2.7 and $2.2 \text{ kcal mol}^{-1}$ and the activation energies above the high temperature minima are 10.2 and $10.3 \text{ kcal mol}^{-1}$, respectively, for TEA bromide and iodide. Although the low temperature minimum of TEA bromide does not agree with the calculated result, the T_1 minimum for TEA iodide is 42 ms, which is close to the calculated 34 ms. This discrepancy between the observed and calculated minimum values of T_1 may be due to a nonrandomness in the reorientations of the methyl group about its threefold axis at low temperatures. A discrepancy like here was also encountered by Reilly et al. [9] at the reorientations of the methyl group in methyl ammonium aluminum sulfate.

For TPA iodide there are two minima in T_1 , and again the low temperature minimum does not agree with the calculated result. The activation energy corresponding to the linear portion above the low temperature minimum was found to be $2.3 \text{ kcal mol}^{-1}$ and that above the high temperature minimum 11 kcal mol^{-1} .

For TBA bromide and iodide the activation energies were found to be 2.5 and $2.8 \text{ kcal mol}^{-1}$, respectively. The activation energies for THA bromide and iodide in Figs. 9 and 10 are 2.2 and $2.0 \text{ kcal mol}^{-1}$. The experimental T_1 minima, 160 ms, for THA halides are much longer than the predicted 34 ms. This difference and the somewhat low activation energies for the relaxation seem to suggest quantum mechanical torsional state effects [10] rather than classical reorientation of the methyl groups at low temperatures. On the other hand CH_2 protons certainly have longer T_1 's than CH_3 protons, and due to spin-exchange averaging the observed single T_1 's for TEA, TPA, TBA and THA halides can be expected to be longer than those for TMA halides, which include no CH_2 groups. Also

this may be the reason for the discrepancies between the experimental and the calculated T_1 minima for TEA, TPA, TBA, and THA halides.

This study was supported in part by the Magnetic Resonance Research Unit of TBTAk (Turkish Scientific and Technical Research Council).

- [1] J. G. Powles and H. S. Gutowsky, *J. Chem. Phys.* **21**, 1695 (1953).
- [2] E. O. Stejskal, D. E. Woessner, T. C. Farrar, and H. S. Gutowsky, *J. Chem. Phys.* **31**, 55 (1959).
- [3] G. W. Smith, *J. Chem. Phys.* **42**, 4229 (1965).
- [4] S. Albert and J. A. Ripmeester, *J. Chem. Phys.* **57**, 2642 (1972).
- [5] T. T. Ang and B. A. Dunell, *Can. J. Chem.* **54**, 1985 (1976).
- [6] F. Köksal, *J. Chem. Soc.*, to be published.
- [7] T. T. Ang and B. A. Dunell, *J. Chem. Soc. Faraday Trans. 2*, **75**, 169 (1979).
- [8] S. Albert, H. S. Gutowsky, and J. A. Ripmeester, *J. Chem. Phys.* **56**, 3672 (1972).
- [9] D. E. O'Reilly and T. Tsang, *Phys. Rev.* **157**, 417 (1967).
- [10] P. S. Allen and S. Clough, *Phys. Rev. Letters* **22**, 1351 (1969).